Recurrent Fever SYPHILIS

Frambœsia Sleeping Sickness

Practical work with GALYL in the shape of

INTRAVENOUS (Dilute and Concentrated) and INTRAMUSCULAR INJECTIONS administered in
Military, Naval. and the principal General Henricals throughout the United Vision of the Principal General Henricals throughout the United Vision of the Principal General Henricals throughout the United Vision of the Principal General Henricals throughout the United Vision of the Principal General Henricals and Theorem 1981 (1981) and Theorem 2011 (1981) and Military, Naval, and the principal General Hospitals throughout the United Kingdom, has demonstrated that this preparation is more rapid and less toxic in action than any compound of the "606" group, which accounts for the consistently excellent clinical results without any undesirable by-effects.


FOR INTRAVENOUS INJECTIONS:—
(1) DILUTE.—GALYL is supplied in neutral glass ampoules containing the necessary dose of Sodium Carbonate, sterile distilled water only being stead for the discouler. used for the dissolution.

(2) CONCENTRATED:—A special outfit containing one dose GALYL, one ampoule sterilised solution, and one small filter is supplied.

Doses: 0·10—0·15—0·20—0·25—0·30—0·35—0·40.

(3) FOR INTRAMUSCULAR INJECTIONS: GALYL is supplied in OILY EMULSION.

Doses: 0.10-0.15-0.20-0.30-0.46.




HECTINE is an anti-syphilitic of low arsenic percentage, it is employed as a treatment following Galyl or ab initio in all stages of the disease. HECTINE is valuable in cases requiring prolonged treatment. FORMS.—The preparation is put up in hermetically sealed ampoules for Intramuscutar Injection and in pills for Internal administration. HECTINE is Sodii Benzo-sulpho-p-amnio-phenyl arsonas. Containing 21 per cent. of arsenic.

HECTARGYRE Mercurial Salt of Hectine, a combined Arsenico-Mercurial treatment of Syphilis.

of For the **Medical Treatment** CANCER.

Extract from Proceedings of the Royal Society of Medicine, February, 1918:—
Dr. George Herschell, comparing electric and chemical colloid copper, attated: Chemical colloid appears to give the best results. In the following cases chemically prepared colloid was used. Experimentally it has been proved: (1) That particles of colloid copper can be demonstrated in the granulations of cancerous growths after two or three injections of the colloid; and clinically (2) There is invariably great relief from pain; (3) Appetite and strength return and the patient puts on flesh; (4) There is in many cases a diminution in the size of the tumour; (5) The injections are absolutely non-toxic, although in many cases an inflammatory reaction is manifested in the cancerous growths.

"As regards my own personal experience in the use of this method of treatment in addition to my own cases, I have had an opportunity of observing the effects in the practice of some of my medical friends, and in nearly all cases the progress of the disease appeared to be arrested."



CUPRASE is a Colloidal Copper hydroxide which is obtained chemically by the reduction of Salts of Copper in the presence of Albumosic Acid.

This treatment has passed far beyond the experimental stage. The numerous clinical reports from Doctors of repute in various countries give cases of a great variety. In all of these pain has been eliminated, and in a good percentage cures are claimed, and in no instance any undesirable effects.

LITERATURE, CLINICAL REPORTS, AND PRICE LIST ON REQUEST.

THE ANGLO-FRENCH DRUG CO., LTD., (late M. Bresillon & Co.), GAMAGE BUILDING, HOLBORN, LONDON, E.C. Telephone: Holborn 1311.