Recurrent Fever SYPHILIS

4,000 injections (intrayenous) were given in one of our Service Hospitals.

"No case has given the slightest cause for anxiety, and the clinical results are satisfactory."

Staff Surgeon's Report.

GALYL

Frambœsia
Sleeping Sickness

GALYL is as effective as SALYARSAN or NEOSALYARSAN on Spirochætes and Trypanosomes, more rapid in action, and free from

the neurotropic and congestive action of these preparations.

FOR INTRAVENOUS INJECTIONS:-

(1) **DILUTE.**—GALYL is supplied in neutral glass ampoules containing the necessary dose of So-lium Carbonate, sterile distilled water only being used for the dissolution.

(2) **CONCENTRATED.** — A special outfit containing one dose GALYL, one ampoule sterilised solution, and one small filter is supplied.

Doses. 0·10-0·15-0·20-0·25-0·30 -0·35-0·40, (3) FOR INTRAMUSCULAR INJECTIONS:-

GALYL is supplied in GLU-COSE SOLUTION.

Doses. 0·10-0·15-0·20-0·30..

THE IMPROVED INTRAMUSCULAR INJECTION.
A STABLE SOLUTION NOT an Emulsion.

The GALYL is in the form of a Sodium Salt in a weak solution of Glucose. This is much easier to use and more rapidly absorbed than the oily emulsion. It is in every way identical with the intravenous,—efficacy, dose, price, etc.,—only differing in administration.

Thorough and exhaustive tests in various parts of the Kingdom have proved it to be equal in every way with the intravenous; moreover, patients may resume and continue their occupations without inconvenience after injection of intramuscular GALYL.



HYPERTHERMAL THERAPY.

AMBRINE

Ambrine is the only Authorised preparation of Dr. Barthe de Sandfort. We are compelled to state this, as there are various imitations.



May be applied with camel-ha'r brush, but our specially designed spray greatly facilitates the application of Ambrine, especially in the case of deep wounds or extensive surfaces.

Ambrine forms a dressing which INSTANTLY alleviates pain—promotes rapid healing—forms a sure protection against infection of the wound—can be removed without pain, hemorrhage, or injury to the newly-formed tissues. The healing leaves no scars or contractions.

The most efficient means and methods of ensuring active Hyperthermal-Therapy to the practical treatment of Rheumatism, Gout, Arthritis, Yaricose Ulcers, and similar affections.

Sins,—I thank you for sending me the Ambrine; I found it very effective in a c monic and painful varicose ulceration of the leg. Please send another 5s, worth (Signed) A. G. W—, M.D.

The treatment of CARBUNCLES, BOILS, ANTHRAX, ACNE, STYES, and Diseases arising from Staphylococcus.

STANNOXYL

(An Oxide of Tin and Tin Metal free from Lead).

A scientific production the value of which has been studied very closely. The effect is very prompt; from the second day of treatment the pain is relieved, the carbuncles begin to dry up, and those which are just opening are stopped in their course; the core is not expelled but reabsorbed.

In the majority of cases definite results are obtained within five or six days; it is seldom necessary to take the full ten days' treatment, and relapses are unknown.

The daily dose for Adults is 4 to 8 tablets; Children 2 to 4 tablets.

Supplied in vials of 80 tablets.

Further Literature sent on Application to Medical Practitioners and Chemists by the Sole Agents for the British Empire-

THE ANGLO-FRENCH DRUG CO., Ltd., Gamage Building, Holborn, London, E.C. 1.

Telephone: Holionn 1311.

(late M. BRESILLON & Co.).

Telegrams: "AMPSALVAS, LONDON."