The Treatment of CARBUNGLES, BOILS, ANTHRAX, AGNE, STYES, and diseases arising from Staphylococcus.

STANNOXYL

(An Oxide of Tin and Tin Metal Free from Lead).

A scientific production the value of which has been studied very closely. The effect is very prompt; from the second day of treatment the pain is relieved, the carbuncles begin to dry up, and those which are just opening are stopped in their course; the core is not expelled but reabsorbed.

In the majority of cases definite results are obtained within five or six days; it is seldom necessary to take the full ten days' treatment, and relapses are unknown.

The daily dose for Adults is 4 to 8 tablets; Children 2 to 4 tablets.

Supplied in via's of 80 tablets.

The SCIENTIFIC TREATMENT of INFLUENZA and Allied Ailments.

KINECTINE

According to Dr. Mouneyrat, the discoverer of Galyl (the widely adopted Salvarsan Substitute).

FORMULA:—Chlorhydrate of Quinine c. Hectins—i.e., Benzo-sulfone-para-amino-phenyl-arsenate of Quinine.

Non-toxic, produces no ill results. Easily taken (tablets) and well tolerated.

Highly satisfactory clinical results.

Not only a prophylactic against, but a specific in, INFLUENZA,

Catarrh, Coryza, NEURALGIA, Malaria, &c.

Sir,-I enclose P.O. for the tube of Kinectine. The drug has given every satisfaction. The Angle-French Drug Co., Ltd., Holborn, London, E.C.

H.M.S. ——, 27-7-17.
W. B. H. W. Surgeon, R.N.

RATIONAL TREATMENT OF CONSTIPATION

By the double action of secretions and peristalsis.

OPOLAXYL

Opolaxyl is a combination of the secretions of the liver (biliary), pancreas, and intestines, with vegetable extract of a non-drastic nature.

It combines all the secretions to correspond to nature's therapy and promotes a flow of bile and glycogen with their hamatopoietic and antitoxic properties.

It is a normal regulator of the gastro-intestinal functions, consequently it improves the metabolic exchanges in the entire organism.

Opolaxyl does not lose its efficacy by prolonged use; its effect is gentle and constant, with the least irritation. It is not merely purgative; it stimulates the defective organs and revives the normal equilibrium; it has a lasting effect by reason of it rebuilding the organs, thus correcting diminutional function.

Opolaxyl is put up in small size tabular form easy to swallow and should be swallowed without crunching at bedtime or before breakfast.

Dose-For obstinate constipation 2 or 3 tablets, afterwards 1 tablet every 3 or 4 days for a month

DEAR SIRS.—Having derived considerable benefit from the use of Opolaxyl Tablets in my own personal case, I write as a practising physician to ask if you would make any reduction of price in the event of my sending you an order for a fair quantity—say one dozen bottles.

Yours faithfully,

C. A———, M.D.

LITERATURE ON APPLICATION.

The ANGLO-FRENCH DRUG Co., Ltd., Gamage Building, Holborn, London, E.C. 1.

Telephone: Holborn 1311.

(Late M. Bresillon & Co.)

Telegrams: "AMPSALVAS, LONDON."