

*Iodine in its Most Reliable and Palatable Form.*

# Iodogénol

**IODINE**

in its

**MOST RELIABLE FORM.**

**PÉPIN**

**IODINE**

in its

**MOST PALATABLE FORM.**

**IODOGENOL** is a preparation containing Iodine in an organic, assimilable and one might almost say "living" form. **IODOGENOL** possesses about **38** times the physiological activity of that of iodide of potassium, and although this preparation has, according to the clinical reports of eminent professors, succeeded where the usual iodide treatment had failed, no undesirable by-effects are produced.

**IODOGENOL does not produce iodism.**

**IODOGENOL** is an undoubted digestive stimulant and promotes appetite and has a markedly beneficial effect on the general nutrition. It has been tried in many of the large hospitals, and given highly satisfactory clinical results in cases of **Syphilis, Rheumatism, the various phases of Tuberculosis, General Debility, etc.**

*20 minims of IODOGENOL are equivalent to 8 grs. Iodide of Potassium.*

*The Combined Treatment of SYPHILIS.*

# SUPSALVS




Fig. 1.

**STABLE SUPPOSITORIES OF "606" (OF FRENCH MANUFACTURE)**

At the International Congress of Medicine Ehrlich stated that the biochemical action of "606" on spirochaetes is not direct but indirect, a third factor found in the body fluids being necessary.

This success is explained by the well-known experiment of Levaditi: "If living treponemas be placed in a solution of Arsenobenzol (Ehrlich's 606) they continue to live in it. But if a trace of extract of liver be added to the mixture the treponemas are destroyed."

"If 606 has to be taken up and transformed by the liver in order to become toxic to the treponema, there is no better mode of absorption of the drug than by way of the intestine, since all the veins of the intestines join the portal vein. If this be the case no route could be more indirect and more unsatisfactory for active treatment than one that is not intestinal or not intravenous (i.e. prehepatic), since some of the drug must necessarily become fixed everywhere before the passage through the liver has activated it."—Dr. SAFOURAUD, La Clinique (13-4-1913).

As a result of numerous clinical experiments, Dr. Bagrov, of Moscow, has arrived at the same conclusion, and recommends the rectal method of administration of 606.

**Extremely Simple in use.**

**No Ill-effects.**

**Most Satisfactory Clinical Results.**

**Rapid Absorption.**

Fig. 1 represents the special patent metallic envelope containing a Supsalvs for hot climates. The projecting edge in the middle is cut by scissors—the two ends are then easily pulled apart. The envelope should first be immersed in cool or iced water for 10 minutes.

**200** cases were treated by the combined treatment in one of the London hospitals. In each case a negative reaction was attained.

# MERSALV

**FOR MERCURIAL INJECTION IN CONNECTION WITH SUPSALVS TREATMENT**

**CHEMISTRY.**—"Mersalv" contains 10 per cent. metallic mercury, which by a special mechanical process exists in an exceedingly minute state of subdivision. It is a non-greasy preparation, and, in contra-distinction to other mercurial preparations, contains no organic fat or oils. "Mersalv" is of a white creamy consistence, of pleasant odour, and cleanly in application.

**IN SPECIAL GLASS-STOPPERED BOTTLES FOR HOT CLIMATES.**

*Further Literature sent on Application to Medical Practitioners by the Sole Agents for the British Empire—*

**The ANGLO-FRENCH DRUG Co., Ltd., Gamage Building, Holborn, London, E.C. 1.**

Telephone: HOLBORN 1311.

(Late M. BRILLON & Co.)

Telegrams: "AMPSALVS, LONDON."