

THE MOST EFFECTIVE TREATMENT OF VENEREAL DISEASES.

GALYL

INTRAVENOUS.

INTRAMUSCULAR.

Identical in Dose and Efficacy.

60,000 Injections administered in **Naval, Military and General Hospitals** have demonstrated that Galyl is **efficient, rapid and well-tolerated**, which accounts for the consistently excellent results obtained without undesirable after-effects.

Intravenous GALYL.

After 5,000 Injections administered in one of our Service Hospitals,

Staff Surgeon reports:

"No case has given the slightest cause for anxiety, and the clinical results are very satisfactory."—*Staff Surgeon's Report.*

Intramuscular GALYL

is supplied in a Solution of Glucose and is therefore much easier to use and more rapidly absorbed than an oily emulsion. Exhaustive tests have proved it to be as effective as the Intravenous Injection.

The treatment of **CARBUNCLES, BOILS, ANTHRAX, ACNE, STYES**, and Diseases arising from **Staphylococcus**.

STANNOXYL

ROBERT & CARRIÈRE, PARIS.

(A Oxide of Tin and Tin Metal Free from Lead).

A scientific production the value of which has been studied very closely. The effect is very prompt; from the second day of treatment the pain is relieved, the carbuncles begin to dry up, and those which are just opening are stopped in their course; the core is not expelled but reabsorbed.

In the majority of cases definite results are obtained within five or six days; it is seldom necessary to take the full ten days' treatment, and relapses are unknown.

Vide THE LANCET, Jan. 19th, 1918, pp. 99 and 109.

"GENTLEMAN.—I have not found anything to equal Stannoxyl in the treatment of Acne Vulgaris. In the cases I have used it, the results have been excellent. I have tried it in a few cases of Skin infection with encouraging results.—Yours faithfully, ———, F.R.C.S." "March, 1918."

The daily dose for Adults is 4 to 8 tablets;

Children 2 to 4 tablets.

Supplied in vials of 80 tablets.

Rational Treatment of Constipation

By the double action of secretions and peristalsis.

OPOLAXYL

Opolaxyl is a combination of the secretions of the liver (biliary), pancreas, and intestines, with vegetable extract of a non-drastring nature.

It combines all the secretions to correspond to nature's therapy and promotes a flow of bile and glycogen.

It is a normal regulator of the gastro-intestinal functions, consequently it improves the metabolic changes in the entire organism.

Opolaxyl does not lose its efficacy by prolonged use; its effect is gentle and constant, with the least irritation. It is not merely purgative; it stimulates the defective organs and revives the normal equilibrium; it has a lasting effect by reason of it rebuilding the organs, thus correcting diminutional function.

Opolaxyl is put up in small size tabular form and should be swallowed without crunching at bedtime or before breakfast.

DOSE—For obstinate constipation 2 or 3 tablets, afterwards 1 tablet every 3 or 4 days for a month.

Further Literature sent to the Medical Profession on application to the Sole Agents for the British Empire—

The ANGLO-FRENCH DRUG Co., Ltd., Gamage Building, Holborn, London, E.C. 1.

Telephones: HOLBORN 1311.

Telegrams: "AMPSALVAS, LONDON."