BRITISH MEDICAL JOURNAL

MARCH 28 1942

TABLE OF CONTENTS

ADDRESSES AND PAPERS	LEADING ARTICLES	CORRESPONDENCE
ADDRESSES AND PAPERS Rehabilitation in the Royal Air Force. R. WATSON-JONES, M.Ch., F.R.C.S	ANNOTATIONS Tropical Hygiene in India	War Neurosis. ELIOT SLATER, M.D. 421 How Much Vitamin D? HELEN M. M. MACKAY, M.D
Sterilization of Surgical Rubber Gloves. D. ASTLEY SANFORD, M.B., F.R.C.S., and H. A. COOKSON, M.B., F.R.C.P.Ed	C. H. Usher, F.R.C.S.Ed	Preserving the Race. Sir Francis Fremantle, M.D., M.P
GENERAL ARTICLES AND NEWS Preparations and Appliances: Splint for Use after Torek's Operation for Undescended Testis416 GALLANTRY IN CIVIL DEFENCE	REVIEWS The Hormonal System	MEDICAL MEMORANDA Objective Tinnitus. Neville Young, F.R.C.S.Ed

ANACARDONE

(Nikethamide B.D.H.)

Anacardone has won universal recognition as a potent cardiac and respiratory stimulant.

In extreme cases in which an immediate response is essential, Anacardone should be given intravenously and in adequate dosage (5 c.c. or more).

In less urgent states of collapse, cardiac failure, respiratory depression during anæsthesia, surgical shock or narcotic poisoning, Anacardone may be administered intramuscularly or even orally.

Anacardone is available at the following net prices to members of the medical profession:

SOLUTION (INJECTION OF	F NIKE	THAM	IDE B	.P.)			
Box of 6 x 2 c.c. ampoules	•••	•••		•••	4s. od.		
Box of 12 x 2 c.c. ampoules	•••	•••	•••	•••	7s. od.		
Box of 3 x 5 c.c. ampoules			•••		5s. od.		
Box of 12 x 5 c.c. ampoules	•••	•••	•••	•••	16s. 9 d.		
SOLUTION FOR ORAL ADMINISTRATION							
Bottle of 15 c.c	•••	•••		•••	3s. 6d.		
Bottle of 100 c.c	•••	•••	•••		18s. 3d		

THE BRITISH DRUG HOUSES LTD. LONDON N.1

Telephone : Clerkenwell 3000 Telegrams : Tetradome Telex London

Ancd/E/15

ISOFLAV

SOLUTION TABLETS

Buffered Proflavine Sulphate Isotonic

For the Prevention and Control of Wound Infection



Isoflav has given excellent results as an antiseptic application for wounds of the brain. It is effective against bacteria and practically innocuous to brain tissue.

★ Extract from British Medical Journal, 1941, March 8th, p. 378:-

In five cases solutions of 0.1 per cent. proflavine sulphate at pH 6.3 (made from Boets Isoflav tablets) were used freely as irrigating lotions to the exposed brain and skull wounds after débridement had been carried out and before the wounds were closed by primary suture. In another case of small penetrating shot-gun wounds where débridement was technically impossible the various puncture wounds were treated by a single irrigation with proflavine solution. Only one out of the six patients died, the remainder, including the case of shot-gun wounds, healing by first intention without any signs of wound infection or edema of the brain. The one death occurred after operation from shock due to multiple injuries to other parts of the body; histological examination of the brain showed no damage in excess of the trauma inflicted.

Bottle of 50 tablets ... $5/7\frac{1}{2}$ Nett

Obtainable through all branches of



Literature sent upon request

BOOTS PURE DRUG CO. LTD NOTTINGHAM

GENTIAN VIOLET JELLY

with

'MERTHIOLATE' Brand

SODIUM ETHYL MERCURI THIOSALICYLATE

FOR THE TREATMENT OF BURNS

This jelly, containing 1% Gentian Violet, provides a convenient method of applying the dye to burned areas. Particularly adapted for use in burns about the face and hands, areas where the skin is thin, and in body flexures. The addition of 'Merthiolate' which is an excellent general germicide with a low toxicity for tissues, enhances the value of the product and further lessens the possibility of infection.

Supplied in one-ounce and five-ounce collapsible tubes.

References: British Medical Journal, November 16th, 1940, p. 679; The LANCET, November 16th, 1940, p. 627; Practitioner, January, 1941, p. 27.

ELI LILLY AND COMPANY LIMITED BASINGSTOKE AND LONDON