



## Prescribe

## KAYLENE-OL for

Toxaemia and Constipation during the Antenatal and Sostnatal Periods

Formula:

Kaylene (Colloidal Kaolin) 7.5%, Liquid Faraffin 25%

DOSE.—Kaylene-ol, 1 to 2 dessertspoonfuls half an hour before breakfast and at bedtime. If constipation is marked, substitute KAYLENE-OL & PHENOL PHTHALEIN (0.5%).

Samples and literature on request.

## KAYLENE, LIMITED

Sole Distributors: ADSORBENTS, LTD. WATERLOO ROAD LONDON N.W 2



## "FRUIT SALT"

for normal bowel activity

With good reason, ENO's "Fruit Salt" might be regarded as the ideal laxative. It is free from undesirable side-actions. It is suitable for the young, the aged, the invalid, the convalescent and the case of pregnancy. ENO's entails no risk of systemic dehydration, and because of its freedom from sugar it is also safe for diabetic cases. By its purity, by its palatable, refreshing taste it has established itself all the world over as an ideal send-off for the day.

ENO'S "Fuit Salt" is prepared by a special process and contains approximately: Sod. Bicarb 53.25%, Acid Tart. 43.50%, Sod Bitart. 1.50%, Sod Tart. 1.75%

J. C. ENO, Ltd., Medical Dept. Grear West Road, Brentford Mx

