

## Urolucosil<sup>★</sup>

THE SULPHONAMIDE OF CHOICE  
IN B. COLI INFECTIONS OF THE  
URINARY TRACT


The recommended dose for adults is 0.10—0.20 G. five to six times daily: this gives a total daily dose of under 2 G. in 24 hours. Because of this small daily dose and its remarkably high solubility, and because of the small degree of acetylation, Urolucosil is the drug of choice in the treatment of urinary-tract infections.

*The characteristics of Urolucosil may be summarized as follows:—*

- 1 Extremely well suited for the treatment of infections of the urinary tract, especially in cases of uncomplicated infections with B. coli. Not advised for infections due to Staph. Aureus, B. pyocyaneus, B. proteus, Strept. faecalis.
- 2 Easily and completely absorbed in from 1 to 2 hours.
- 3 Rapidly excreted in the urine, almost exclusively in an active non-acetylated form.
- 4 Side effects seldom observed; no disturbance of intestinal flora; no changes in the blood picture.
- 5 Treatment is very safe in view of the small amount (1.2 G.) administered over 24 hours, even though it is repeated at frequent intervals.

*Contra indicated, as with other sulphonamides, where there is organic or mechanical obstruction.*

*Descriptive literature on Urolucosil and trial quantities will be sent to registered medical practitioners on written request.*



*William R. WARNER and Co., Ltd., Power Road, London W. 4.*