

PHILIPS

what are you missing on TV tonight?

If you have ever missed a favourite programme through working too late, being on call or away from home, you will appreciate the Philips N1700 VCR television recorder.

The N1700 will record $2\frac{1}{2}$ hours of what you want to watch, and play it back when you want to watch it. Further, a 3-day clock allows you to set the machine to switch itself on in your absence, and record something you'd otherwise have missed.

And remarkably, if you are at home and there's something worth watching on both sides, you can watch one and record the other.

The Philips N1700 television recorder allows you to be really selective about your viewing—now, even more than you may have imagined—with a new channel of medical programmes, sent to you on video tape cassette . . . called . . .




the video journal of medicine!

Britain's new and unique television journal has been created by doctors for doctors. It will now be published on the Philips N1700 format. Each hour-long edition of the Video Journal of Medicine brings the very best in medicine to its subscribers.

Contributions from eminent specialists around the world ensure a balanced, often critical, but always authoritative opinion on the most recent treatment methods in addition to detailed reviews of existing therapies.

The VJM can go much further than any conventional reading matter. Through the medium of television it can condense a day's heavy reading into an hour's effortless and convenient viewing.


That's the advantage of The Video Journal of Medicine and the Philips N1700 television recorder. Let us send you a brochure and the name of your local Philips television recorder specialist. Contact us and we can also advise you of the professionally related savings, including tax benefits you can obtain on purchasing a television recorder when subscribing to the VJM.



Recording and playback of material may require consent—see the Copyright Act 1956; also the Performers' Protection Acts 1958 to 1972.

**Write to: The Sales Director,
The Video Journal of Medicine, 13/14 Bateman St, London W1V 6EB or Telephone 01-437 9611**

Will your practice stand up if your car breaks down?



Car Finance for Doctors and Dentists

Get details of this scheme for Car purchase offered to the profession by

Medical Sickness Finance Corporation Limited

Company registration number 464750-London

- Minimum deposit
- Maximum period of repayment
- Payments waived on death

For particulars write to us mentioning this advertisement at

7-10 Chandos Street, Cavendish Square, London W1A 2LN or Telephone: 01-636 1686

Registered Office



A MEMBER OF THE MEDICAL SICKNESS GROUP

ADVERTISEMENTS

PHARMACEUTICAL PRODUCTS

Readers are reminded that prescribers' literature and clinical information are always available from advertisers on request.



ILONIUM

a powerful antiseptic

Isonium can be used in the treatment of most pus-bearing wounds. It will be found to effectively assist in clearing up such conditions as sloughing bedsores, suppurative varicose ulcers, boils, whitlows and carbuncles.

ILONIUM is an antiseptic ointment with some analgesic properties. It contains no antibiotics and apart from phenol its constituents are mainly derived from natural oils and oleoresins.

Isonium is available in sizes of 450g, 100g and 30g.

COMPOSITION: Venice turpentine Colophony, Phenol, Chlorophyll, Pinene, Camphene, Cineole, Borneol, Thymol, Dipentene.

Full information by return from
ILON LABORATORIES (Hamilton) LTD., Lorne Street, Hamilton.

PUVA

is now available
for private patients at

PSORIASIS UVA (Medical Referrals Only) Ltd.,
2 HARMONT HOUSE,
20 HARLEY STREET, London W.1.

Telephone 01-580 0875

For details please contact
the Physician-in-Charge.

THE RETREAT, YORK for Psychiatric Illnesses

Founded in 1792 by the Quaker, William Tuke, who established the tradition of concern for patients as individuals, The Retreat is a 250-bedded private registered nursing home surrounded by extensive grounds on the outskirts of the historic City of York. It is easily reached by rail and motorways.

Care and treatment is offered for most types of psychiatric illness on the short or long term in a sympathetic and friendly atmosphere. Patients suffering from neuroses, psychoses, alcoholism and dementia are treated in surroundings suitable for their individual needs.

The Nursing Home is a registered charity and is able to offer inclusive care in shared accommodation from £12.65 per day or in single rooms at slightly increased charge. The Nursing Home is recognised by the main private patient schemes.

For further details apply to The Medical Director, The Retreat, York. YO1 5BN (Tel. 0904-54551).

You know they're in safe hands with a trained nurse.

There are always times when a trained British Nursing Association nurse would be a great help.

And it need not be full-time.

Because we can provide a nurse for as long or as short a time as you need.

We are the largest nursing agency in the U.K. With qualified nurses always available. 24 hours a day and 7 days a week.

They're experienced, dedicated and of the highest professional standard.

If you feel that a trained B.N.A. nurse would be a help in your life, you have only to phone or write.

We will send you a detailed brochure.

BNA

British Nursing Association,
270 Oxford Street,
London W1N 0HQ.
Tel: 01-409 1921

or see our local address in Yellow Pages.



For a beast of a cough



ACTIFED COMPOUND LINCTUS

**The cough-stopper for
adults and children alike**

Actifed* Compound Linctus contains triprolidine hydrochloride, pseudoephedrine hydrochloride, and codeine phosphate in a palatable fruit-flavoured syrup.

*Trade Mark

Full prescribing information is available on request.
Wellcome Medical Division
The Wellcome Foundation Ltd.
Berkhamsted, Herts



Wellcome

Prothiaden

What's the significance of the new 75mg tablet?

Presentation Prothiaden is dothiepin hydrochloride, an antidepressant of the tricyclic group.

Prothiaden is available as sugar-coated tablets, each containing 75mg of dothiepin hydrochloride. The tablets are red in colour and bear the overprint 'P 75' in white. It is also available in hard gelatin capsules, each containing 25mg of dothiepin hydrochloride. The capsules are red, brown in colour with the overprint 'P25' in white.

Uses Prothiaden is indicated in the treatment of depression and the anxiety frequently associated with depressive illness.

Dosage and Administration Prothiaden should be given in a dosage of 75 to 150mg daily. The following dosage schemes are suggested.

Mild to moderate depression, 25mg three times daily or 75mg at night.
Moderate to severe depression, 150mg daily in divided doses, or as a single dose at night. (If the latter regimen is adopted, it is preferable to use a smaller dose for the first few days).

In certain circumstances, i.e. in hospital use, Prothiaden has been given at dosages up to 225mg daily. No dietary restrictions are necessary during treatment with Prothiaden.

Contra-indications, Warnings, etc. Prothiaden has anticholinergic properties; therefore, it may precipitate urinary retention in susceptible individuals and its use should be avoided in patients with existing or potential urinary retention. Patients with closed-angle glaucoma should not be given Prothiaden and the occurrence of a painful red eye in a patient receiving the drug may indicate acute closed-angle glaucoma; this requires urgent treatment. In patients with chronic simple glaucoma, the risk of Prothiaden causing a rise in intraocular pressure is relatively small provided adequate anti-glaucoma therapy is being used. Caution is advised when treating epileptic patients and those with cardiovascular disorders. From studies in animals, it was concluded that Prothiaden had no teratogenic effects in the species tested. Nevertheless, as with any relatively new drug, the use of Prothiaden during pregnancy should be avoided if possible.

Use with other drugs: Prothiaden should not be given concurrently with MAO inhibitors; nor should it be given within 14 days of ceasing treatment with an MAO inhibitor. Prothiaden may alter the pharmacological effects of some concurrently administered drugs: CNS depressants, including alcohol and narcotic analgesics, will be potentiated, as will the effects of adrenaline and noradrenaline (it should be borne in mind that some local anaesthetic preparations contain these sympathomimetics). The hypotensive effect of certain antihypertensive agents (e.g. bethanidine, debrisoquine, guanethidine) may be reduced.

Side-effects: Generally, the side-effects associated with Prothiaden have been mild and controlled by reduced dosage. The following have been reported - dryness of mouth, constipation, disturbed accommodation, lassitude, dizziness, orthostatic hypotension, palpitations, somnolence, tremor, headache.

Overdosage: The symptoms of overdosage with Prothiaden may include sedation, dry mouth, blurring of vision, tachycardia, tremor, sweating, nausea, vomiting, confusion. The main dangers from overdosage arise from unconsciousness, convulsions, abnormal cardiac rhythms, hypotension and depression of respiration. The smallest dose of Prothiaden alone which resulted in the death of an adult was reported to be 0.75-1.0g (30 to 40 x 25mg capsules). The largest dose from which recovery took place was reported to be 5.0g (200 x 25mg capsules). In view of the many factors which influence the outcome of an overdose, these figures should not be considered in isolation.

Treatment of overdosage: Gastric lavage, in the unconscious patient or where the cough reflex is depressed, the lungs should be protected by a cuffed endotracheal tube. Repeated gastric/intestinal aspiration may remove drug and metabolites excreted into the gut via the bile. General support of the circulation with continuous ECG monitoring is advised. Abnormalities of cardiac rhythm and epileptic convulsions may occur and should be treated accordingly. Forced diuresis and haemodialysis are not recommended. Bed-rest is advisable, even after clinical recovery.

Pharmaceutical Precautions - Recommended storage conditions 5°C to 20°C.

Legal Category POM.

Package Quantities - Prothiaden Tablets (75mg) 100, 500. Prothiaden Capsules (25mg) 100, 600.

Basic NHS Prices: 100 x 25mg £1.57 600 x 25mg £9.03
100 x 75mg £4.52 500 x 75mg £21.70

Product Licence Number Prothiaden capsules PL 0096/5007. Prothiaden Tablets PL 0096/0046.

The Crookes Laboratories Ltd., Basingstoke, Hants.

References: 1. Pearce, J. B. & Linford Rees, W. J., *Int. Med. Res.* 1974, **2**, 12.
2. Rees, J. A. & Cryer, P. C. *Curr. Med. Res. Opin.* 1976, **4**, 416.

Prothiaden 75mg

more effective, better tolerated
treatment for depression and anxiety.

... dothiepin in a single daily dose is better tolerated and produces a greater overall response than a divided dose regimen of amitriptyline.

2

... A single daily dose of dothiepin (Prothiaden) has been shown to produce a better clinical response and to be better tolerated than a divided daily dose regimen. 1

... with dothiepin both the incidence and severity of side-effects were much less than with amitriptyline.

2

H. K. LEWIS of GOWER STREET, LONDON

BRITAIN'S SPECIALIST MEDICAL BOOK CENTRE

BOOKSHOP

Large stocks on all branches of Medicine, Surgery and allied subjects

Catalogues sent free on request

(Please state special interests)

LIBRARY

(Medical and Scientific)

Annual Subscription from £7.00

Reduced rates for multiple subscriptions

(Available in U.K. only)

Prospectus post free on request

H. K. LEWIS & Co. Ltd., 136 GOWER ST., LONDON WC1E 6BS

Telephone: 01-387 4282

Telegrams: Publicavit, London WC1

ERRATUM

Human Growth and its Disorders

W. A. Marshall

Published December 1977

Page 166, paragraph 4, line 2 reads
"Testosterone oenanthate 250mg.

intravenously"

This should, of course, read
intramuscularly as intravenous injection
of this substance could be highly
dangerous.



Academic Press

London New York San Francisco

A Subsidiary of Harcourt Brace Jovanovich, Publishers

24-28 Oval Road, London NW1, England

111 Fifth Avenue, New York, NY 10003, USA

BONE AND JOINT DISEASES

Arthritis used to be the Cinderella of diseases, but improved diagnostic techniques, advances in joint replacement surgery, and a whole range of non-steroidal drugs have led to a much more active approach to the management of the common bone and joint diseases. Detailed advice on their diagnosis and treatment was given last year in a series of 16 articles in the *BMJ*, and these have now been gathered together in the latest volume based on the popular "Today's Treatment" series.

Demy Octavo

Paperback

131 pages

ISBN 0 7279 0026 9

**Price: Inland £3.00; Abroad US \$7.50,
including postage**

Payment must be enclosed with order or a surcharge of 30p will be made for rendering invoices and statements

Order your copy now from

The Publisher, BRITISH MEDICAL JOURNAL,

BMA House, Tavistock Square, London WC1H 9JR

or through any leading bookseller

Caved-S


**heals peptic ulcers
SAFELY**

Caved-S

**heals peptic ulcers
EFFECTIVELY**

Caved-S

**heals peptic ulcers
ECONOMICALLY**



More action, less impetigo

More action because Cicatrin carries the double antibiotic action of neomycin and bacitracin, a combination that rapidly destroys staphylococci and streptococci the principal pathogens in impetigo.

Cicatrin cream contains neomycin, bacitracin, l-cysteine, glycine and dl-threonine. Available in 15g and 30g sizes.


Full prescribing information is available on request.
Calmic Medical Division, The Wellcome Foundation Ltd.
Berkhamsted, Herts



CICATRIN

cream

TRADE MARK



**Sleep disturbance
is one of the first
symptoms of depression**

brand of doxepin

*Trade Mark

Improves sleep...even before it lifts the patient's depression

Sinequan is **not** an hypnotic but a sedative-antidepressant. Helps to relieve sleep disturbances due to depression while alleviating the underlying depression itself. Improves the quality and quantity of sleep...patients experience fewer awakenings... feel more rested in the morning.¹ Can obviate the need for hypnotics.²

Now in a 75mg. capsule for once daily dosage at bedtime

Indications: depression with or without anxiety. **Contraindications:** glaucoma, urinary retention, hypersensitivity to the drug. **Side effects:** dry mouth and drowsiness are most commonly reported. **Precautions:** Sinequan may potentiate other compounds - e.g. monoamine oxidase inhibitors; not recommended in pregnancy or children under 12 years of age. **Packs and Basic N.H.S. Cost:** 10mg capsules (P.L. 57/5032), pack of 100, £2 42; 25mg capsules (P.L. 57/5033), pack of 100, £3 44; 50mg capsules (P.L. 57/5034), pack of 100, £5 70; 75mg capsules (P.L. 57/0133), pack of 60, £5.39. **References:** 1. Scientific Exhibit, Annual Meeting of Massachusetts Medical Society, Boston, Mass. May 30th - June 1st 1972. 2. **Modern Medicine** (G.B.), September 1976.

Further information on request to the Company.


PFIZER LIMITED
SANDWICH, KENT

Introducing

ICE for BIC

Tanderil® Chloramphenicol eye ointment.

An effective non-corticosteroid anti-inflammatory/antibacterial eye preparation indicated for inflammatory conditions of the eye such as:-

Blepharitis



Tanderil® Chloramphenicol eye ointment will relieve the chronic inflammation and treat the primary or secondary bacterial infection.

Trauma



Tanderil® Chloramphenicol eye ointment will soothe the eye, control the inflammatory response and prevent bacterial infection.

Conjunctivitis



Tanderil® Chloramphenicol eye ointment will relieve the inflammation and resolve bacterial infection without enhancing virus replication.

Tanderil® Chloramphenicol eye ointment is available as a white to yellowish white soft ointment containing 10% oxyphenbutazone B.P. and 1% chloramphenicol B.P. as active ingredients and phenyl ethyl alcohol 0.5% as a preservative.

Geigy

Full prescribing information is available.
Geigy Pharmaceuticals, Macclesfield, Cheshire SK10 2LY Telephone (0625) 21933