

Migrave fights migraine effectively



with minimal toxicity

2 PINK MIGRALEVE taken at first warning, will often completely abort impending migraine attacks. Residual symptoms, if any occur, can be controlled with YELLOW MIGRALEVE. Thus, Migrave offers:

- **Proven effectiveness**
A clinical trial showed Migrave effective in 79% of migraine attacks.²
- **Minimal contra-indications**
Relating only to paracetamol and small buclizine content.
- **Less vomiting than with ergotamine¹**
- **Proven patient preference²**

Migrave[®]

NON-ERGOTAMINE

THE MIGRAINE TREATMENT PATIENTS PREFER

Full information available by post only from

International Laboratories Ltd. Sunbury-on-Thames, Middlesex TW16 7HN

Recommended prescribing

On first presentation –
Migrave Duo 24 o.p.
2 Pink stat, 2 Yellow q.q.h.
(The duo pack contains
16 Pink and 8 Yellow tablets.
Basic NHS price 88p.)
Subsequently – taking patient
experience into account,
either Pink or Yellow
Migrave may be prescribed
separately in multiples
of 12, so avoiding a double
prescription charge.
Pink alone will often
completely abort attacks.

References 1) *Medical Digest* (1972) 7, 17

2) *Practitioner* (1973), 211 357.

MIGRALEVE PINK tablets
contain buclizine, paracetamol,
codeine. Migrave Yellow tablets
contain paracetamol, codeine.

The Newington Unit


Ticehurst House, Wadhurst, Sussex TN5 7JA

This newly formed Unit aims to help those with alcohol-related disorders and to tackle the underlying problem of dependence.

General medical and psychological measures will be supplemented by an educative approach, both during the initial period of stay and subsequently after discharge.

Further information from the Medical Director of the Unit. Telephone: 0580 200391.

A Nestor Nursing Home



ADVERTISEMENTS

PHARMACEUTICAL PRODUCTS

Readers are reminded that prescribers' literature and clinical information are always available from advertisers on request.

suicide? despair?



Talk to someone who cares

**TELEPHONES
- COUNTRYWIDE -**

ARE MANNED—DAY AND NIGHT—BY

the samaritans

Are you ordinary enough to be one of them?

You need to be a good listener, who understands how other people feel and wants to help them as a friend.

If you are found suitable you will be given the necessary preparation and supervision.

If you feel you might not be able to cope but would like to try, you are probably the type we are looking for.

The Samaritans, too, need friends. To support what they do you can collect or subscribe money and do other administrative duties which they have not time to carry out themselves.

If you would like to join the Samaritans or are interested in helping please write for details to The Hon. Secretary, The Samaritans, 17 Uxbridge Road, Slough, Berks.



And REMEMBER, if you are suicidal or despairing, you can talk to the Samaritans in complete confidence, any hour—day or night.

-just what the doctor ordered

136 piece cutlery set including pastry forks



Genuine silver plated cutlery produced by Sheffield craftsmen in Kings, Dubarry or Rattail designs and set into a fully lined, solid or inlaid leather mahogany table.

An investment at the price and a beautiful piece of furniture to grace any home.

Special price £295 (usual retail price £425)

* Sterling silver or gold plate – special prices on request.

Place settings of 124, 120, 62, 50 & 44 pieces also available and accessories include carving sets, grapefruit spoons and tea knives.

Telephone Maureen Raymond on 01-359 0520 or contact

Crown Silverware Ltd.

279 UPPER STREET, ISLINGTON, LONDON N1 2TZ

Our application has been accepted. Here's why:

administration direct to the scalp without disturbing even the most elaborate hair-do.

Dermovate Scalp Application
(0.05% w/w clobetasol propionate)
For severe steroid-responsive scalp conditions including psoriasis.

25ml (£1.90*)
Also available in 100ml plastic squeeze bottles.

25 ml
Glaxo Ltd
Greenford

Betnovate
Trade mark
scalp application
30 ml
Glaxo

Betnovate (0.1% w/w betamethasone valerate)
The effective treatment for steroid-responsive scalp conditions such as seborrhoeic dermatitis and the inflammation associated with severe dandruff. Gives prompt relief from itching followed by a highly satisfactory clearance of lesions.

Scalp Application
A specially formulated alcohol/water base slightly gelled to spread across the scalp without running down the face or neck. Cooling. Soothing. Saves all the inconvenience of frequent shampoos or massage with messy gels.

leaders in topical steroid therapy

Prescribing information

Dosage and administration Apply sparingly night and morning until improvement occurs. With Betnovate Scalp Application improvement may be sustained by applying once a day or less frequently.

With Dermovate Scalp Application, once control is achieved, therapy should be discontinued. Repeated short courses may be used to control exacerbations. If continuous steroid treatment is necessary, a less potent preparation should be used.

Side effects Betnovate and Dermovate Scalp Applications are usually well tolerated, but if signs of hypersensitivity appear, application should be stopped immediately. With all topical corticosteroids, when extensive areas are treated, sufficient systemic absorption may

occur to produce the features of hypercorticism. This is more likely to occur in infants and children, with occlusion, or when treatment is prolonged. Local atrophy may occur after prolonged treatment under occlusion. Rarely, treatment of psoriasis with corticosteroids (or its withdrawal) is thought to have provoked the pustular form of the disease.

Precautions Care must be taken to keep the preparation away from the eyes. Do not use near a naked flame. In infants and children avoid long term continuous topical therapy where possible, as adrenal suppression can occur even without occlusion. Select the least potent corticosteroid which will control the disease. Topical administration of corticosteroids to pregnant animals can cause foetal abnormali-

ties. The relevance of this finding to humans has not been established; however, topical steroids should not be used extensively in pregnancy, i.e. in large amounts or for prolonged periods.

Contra-indications Viral and dermatophyte infections of the scalp.

Product licence numbers
Betnovate Scalp Application: PL/0004/5133
Dermovate Scalp Application: PL/0004/0242

Further information on Betnovate and Dermovate (trade marks) Scalp Applications is available from:
Pharmaceuticals Division
Glaxo Laboratories Ltd
Greenford, Middlesex UB6 0HE

*Basic NHS cost (at time of going to press)



Prothiaden

What's the significance of the new 75mg tablet?

Presentation Prothiaden is dothiepin hydrochloride, an antidepressant of the tricyclic group.

Prothiaden is available as sugar-coated tablets, each containing 75mg of dothiepin hydrochloride. The tablets are red in colour and bear the overprint 'P 75' in white. It is also available in hard gelatin capsules, each containing 25mg of dothiepin hydrochloride. The capsules are red/brown in colour with the overprint 'P25' in white.

Uses Prothiaden is indicated in the treatment of depression and the anxiety frequently associated with depressive illness.

Dosage and Administration Prothiaden should be given in a dosage of 75 to 150mg daily. The following dosage schemes are suggested:

Mild to moderate depression, 25mg three times daily or 75mg at night.
Moderate to severe depression, 150mg daily in divided doses, or as a single dose at night. (If the latter regimen is adopted, it is preferable to use a smaller dose for the first few days).

*In certain circumstances, i.e. in hospital use, Prothiaden has been given at dosages up to 225mg daily. No dietary restrictions are necessary during treatment with Prothiaden.

Contra-indications. Warnings, etc. Prothiaden has anticholinergic properties; therefore, it may precipitate urinary retention in susceptible individuals and its use should be avoided in patients with existing or potential urinary retention. Patients with closed-angle glaucoma should not be given Prothiaden and the occurrence of a painful red eye in a patient receiving the drug may indicate acute closed-angle glaucoma; this requires urgent treatment. In patients with chronic simple glaucoma, the risk of Prothiaden causing a rise in intraocular pressure is relatively small provided adequate anti-glaucoma therapy is being used. Caution is advised when treating epileptic patients and those with cardiovascular disorders. From studies in animals, it was concluded that Prothiaden had no teratogenic effects in the species tested. Nevertheless, as with any relatively new drug, the use of Prothiaden during pregnancy should be avoided if possible.

Use with other drugs. Prothiaden should not be given concurrently with MAO inhibitors; nor should it be given within 14 days of ceasing treatment with an MAO inhibitor. Prothiaden may alter the pharmacological effects of some concurrently administered drugs: CNS depressants, including alcohol and narcotic analgesics, will be potentiated, as will the effects of adrenaline and noradrenaline (it should be borne in mind that some local anaesthetic preparations contain these sympathomimetics). The hypotensive effect of certain antihypertensive agents (e.g. bethanidine, debrisoquine, guanethidine) may be reduced.

Side-effects. Generally, the side-effects associated with Prothiaden have been mild and controlled by reduced dosage. The following have been reported—dryness of mouth, constipation, disturbed accommodation, lassitude, dizziness, orthostatic hypotension, palpitations, somnolence, tremor, headache.

Overdosage: The symptoms of overdosage with Prothiaden may include sedation, dry mouth, blurring of vision, tachycardia, tremor, sweating, nausea, vomiting, confusion. The main dangers from overdosage arise from unconsciousness, convulsions, abnormal cardiac rhythms, hypotension and depression of respiration. The smallest dose of Prothiaden alone which resulted in the death of an adult was reported to be 0.75–1.0g (30 to 40 x 25mg capsules). The largest dose from which recovery took place was reported to be 5.0g (200 x 25mg capsules). In view of the many factors which influence the outcome of an overdose, these figures should not be considered in isolation.

Treatment of overdosage: Gastric lavage; in the unconscious patient or where the cough reflex is depressed, the lungs should be protected by a cuffed endotracheal tube. Repeated gastric/intestinal aspiration may remove drug and metabolites excreted into the gut via the bile. General support of the circulation with continuous ECG monitoring is advised. Abnormalities of cardiac rhythm and epileptic convulsions may occur and should be treated accordingly. Forced diuresis and haemodialysis are not recommended. Bed-rest is advisable, even after clinical recovery.

Pharmaceutical Precautions—Recommended storage conditions 5°C to 20°C.

Legal Category POM.

Package Quantities—Prothiaden Tablets (75mg) 100, 500. Prothiaden Capsules (25mg) 100, 600.

Basic NHS Prices: 100 x 25mg £1.57 600 x 25mg £9.03
100 x 75mg £4.52 500 x 75mg £21.70

Product Licence Number Prothiaden capsules PL 0096/5007. Prothiaden Tablets PL 0096/0046.

The Crookes Laboratories Ltd., Basingstoke, Hants. 

References: 1. Pearce, J. B. & Linford Rees, W. J., *Int. Med. Res.* 1974, **2**, 12.
2. Rees, J. A. & Cryer, P. C. *Curr. Med. Res. Opin.* 1976, **4**, 416.

'... dothiepin in a single daily dose is better tolerated and produces a greater overall response than a divided dose regimen of amitriptyline.' 2

'... with dothiepin both the incidence and severity of side-effects were much less than with amitriptyline.' 2

A single daily dose of dothiepin (Prothiaden) has been shown to produce a better clinical response and to be better tolerated than a divided daily dose regimen. 1

Prothiaden 75mg

more effective, better tolerated
treatment for depression and anxiety.

Spot the difference.



With **Quinoderm** the answer is clear. Our **Quinoderm** hydrocortisone cream is invaluable for the initial treatment of acne lesions – the rapid improvement in appearance is of positive benefit to the patient.

Quinoderm Cream provides logical follow-on treatment. By controlling the condition, it reduces the risk of tissue damage and residual scarring. *Clinical reports indicate that acne responds to **Quinoderm**.

Quinoderm
HITS THE SPOT.

Full prescribing information and clinical reports on request
Quinoderm Ltd, Manchester Road, Oldham OL8 4BR

*The Practitioner Vol 153 p. 192
*The Practitioner Vol 202 p. 405
*The Practitioner Vol 208 p. 541

KIRBY LESTER PORTABLE CARDIOGRAPH

Performance at a price
that's hard
to beat!



A single channel
E.C.G. built into
a portable attache
case, with sufficient
battery power for over
100 complete electrocardiograms.

- Max. Frequency Response 100Hz.
- Accepts most paper sizes.
- 3 Sensitivity settings.
- Solid State electronics.
- Serviced on your premises.
- Built in recharge unit.
- 2 paper speeds.
- Fully guaranteed.

**KIRBY
LESTER
LTD**

Osborne Industrial Estate, Waddington Street,
Oldham, OL9 6QQ. Tel: 061-620 1421



The New Ear Plug that Breathes Bilsom Propp-o-Plast

NEW

Propp-o-Plast is a new disposable ear plug with excellent acoustical properties that breathes. Preformed from finely spun mineral fibres, within a thin porous plastic film covering.

HYGIENE

Propp-o-Plast is 100% disposable, which guarantees that the toughest hygiene requirements are met.

The preformed shape is maintained during insertion into the ear, thereby eliminating the need to mould and shape the plug with subsequent contamination of the fibres.


COMFORT

Propp-o-Plast, unlike many types of earplug which merely block the ear canal, breathes. Minute pores in the thin plastic covering allow air and moisture to pass freely. That is why the feeling of pressure or vacuum often experienced with other earplugs is eliminated.

SAFETY

Propp-o-Plast offers improved attenuation in the lower frequency range.

Propp-o-Plast plugs average attenuation is 25dB (A) providing protection for noise situations up to 110dB (A).



Bilsom
for your comfort

Bilsom

BILSOM INTERNATIONAL LIMITED, 47 BELL STREET BMJ
HENLEY-ON-THAMES, OXFORDSHIRE RG9 2AH

Please send me further details: ☐ Propp-o-Plast (Disposable ear plug)
☐ Propp (Disposable ear plug) ☐ Earuown

Name

Company

Position

Address

Is your business protected against the most frightening loss of all?

As an intelligent businessman you've probably tried to provide for every eventuality. Fire and burglary. Bad debts and public liability. Fidelity bonding and professional indemnity.

But what about the most vital part of your business? Probably the biggest contributory

factor in your success. Human beings. Key personnel. Your partners and co-directors.

If someone of immense importance to your business suddenly dies, you may take months—even years—to replace such a person.


Could your business stand the long term effects of such a loss?

You might find that the bank wants to curtail the overdraft. Or you can't meet all-important delivery dates. Or you lose major sales contracts. Meanwhile, what happens to the business?

There is an answer. You can insure against such a loss of vital people. You can be helped over the financial hurdles that result from such losses. Our 134 years of successful money management can provide just the protection you want.

Take steps immediately to look into Equity & Law's Business Assurance schemes. They are more than an investment. They are essential to your business.

Talk today to your financial adviser. Or contact us direct. This is something so important, it could prove costly to delay.



Equity&Law



Equity & Law Life Assurance Society Limited, 20 Lincoln's Inn Fields, London WC2A 3ES.

ears with otitis externa need

Neo-Cortef

EYE-EAR DROPS AND OINTMENT

Supplied:

Neo-Cortef Eye-Ear Drops

In 5 ml. and 10 ml. dropper bottles.

Neo-Cortef Eye-Ear Ointment In 3.9 g. tubes.

Each ml. or g. contains: hydrocortisone acetate, 15 mg., and neomycin sulphate, 5 mg. (equiv. to 3.5 mg. neomycin-base).

Warning: Topical administration of corticosteroids to pregnant animals can cause abnormalities of foetal development. The relevance of this finding to human beings has not been established; however, topical steroids should not be used extensively in the first trimester of pregnancy, i.e. in large amounts or for prolonged periods.

Full product information available on request



Upjohn

738 REGISTERED TRADEMARK NEO-CORTEF 1478UK
UPJOHN LIMITED · CRAWLEY · SUSSEX



SMITH & NEPHEW PHARMACEUTICALS LTD. RANGE OF OPHTHALMOLOGICAL PRODUCTS

Ganda®

The effective non-miotic therapy for primary and secondary open angle glaucoma, available in three strengths:

Ganda 3 - 0.5 guanethidine monosulphate B.P. 5% w/v and adrenaline B.P. 1.0% w/v
Ganda 4 - 1.0% guanethidine monosulphate B.P. 5% w/v and adrenaline B.P. 0.5% w/v
Ganda 5 - 1.0% guanethidine monosulphate B.P. 5% w/v and adrenaline B.P. 1% w/v
P.L. 33/53 PL 33/71 Ganda 4+0.5 PL 33/70 Ganda 5 - 1 PL 33/69

Simplene®

For the reduction of intra-ocular pressure in primary and secondary, open angle glaucoma.

Simplene is supplied as a sterile, ophthalmic solution of adrenaline B.P. and is now available in two strengths:
Simplene 0.5% (Adrenaline B.P. 0.5% w/v)
Simplene 1% (Adrenaline B.P. 1% w/v)
P.L. 33/57,72

Eppy®

For the reduction of intra-ocular pressure in primary and secondary, open angle glaucoma.

Eppy is supplied as a sterile ophthalmic solution of adrenaline B.P. 1% w/v presented in a 7.5ml glass bottle with separate sterile dropper.
P.L. 33/5022

Fluorets®

Sterile, individually wrapped paper strips, each impregnated with fluorescein sodium B.P., designed to permit application of fluorescein to the eye without the risk of bacterial contamination and the subsequent development of potentially dangerous infection.
P.L. 33/5055


Sno®pilo

For the treatment of glaucoma.

Sno pilo is supplied as a sterile ophthalmic solution of pilocarpine hydrochloride B.P. and is presented in a 10ml plastic dropper bottle. It is available in five strengths: 0.5% w/v, 1.0% w/v, 2.0% w/v, 3.0% w/v and 4.0% w/v.
P.L. 33/54-59


Minims®

The comprehensive range of sterile, single-dose, disposable eye drops.
P.L. 33/5000-5021, 33/48, 33/50, 33/55



Full prescribing information is available from: SMITH & NEPHEW PHARMACEUTICALS LTD. Welwyn Garden City · Herts · England

On Call



Call Sheerness 3371—Hollister Stoma Advisory Service

Often it's difficult for a doctor to devote enough time to helping a stoma patient select the most suitable type and size of appliance.

That's why we employ fully qualified and specially trained Stoma Advisory Nurses who can see you or your patient, analyse the problem with expert knowledge and understanding—and recommend the right answer.

For both ileostomy and colostomy patients, Hollister have the most comprehensive range of bags and accessories. So the Stoma Advisory Nurse can ensure that the patient uses equipment that is absolutely right.

Whenever there's a need for specialised help, please remember—we're on call, Sheerness 3371 or write to Abbott Laboratories Ltd., Freepost, Queenborough, Kent. ME11 5BR—no stamp needed.



The Team That Cares



a product of

HOLLISTER

TRADEMARK OF HOLLISTER INCORPORATED, CHICAGO, ILLINOIS, U.S.A.

available from

ABBOTT

Abbott Laboratories Ltd., Queenborough, Kent. ME11 5BR.

Tegretol®

Tablets and syrup

Epilepsy

Tegretol is a most effective anti-epileptic treatment, indicated for the management of temporal lobe or grand mal epilepsy.

Temporal Lobe

Tegretol is increasingly being used as the drug of first choice for controlling the many facets of this complex form of epilepsy.

Grand Mal

Because there is more to the successful management of epilepsy than just seizure control. Tegretol is consolidating its position as the drug of first or second choice.

Mood and Behaviour

can be distinctly improved by using Tegretol, resulting in a smoothing out of behaviour patterns, and an improvement in interpersonal relationships.

Tolerability

There is increasing agreement that Tegretol is a well tolerated therapy without the drawbacks of causing over-sedation, irritability, hirsutism or gum hypertrophy, all of which tend to undermine patient compliance.