

Debrisan beads
magnified x300
blue background

Indication

Small sterile red spherical beads of diameter of 0.1-0.3mm diameter packed in plastic castors or single use sachets.

Uses:

A dressing for moist, small area burns, wounds and indolent ulcers, whether clean or infected.

Preparation and Administration

A thin layer of Debrisan should be applied onto the wound. The beads will absorb the exudate and the tissue exudate is drawn up into the layer. The Debrisan should be renewed before saturation occurs. Depending on the rate of exudation this may be necessary from one to five times a day. Twice daily is usually adequate. When Debrisan is changed the old material is readily rinsed off with water or saline and new material then be sprinkled onto the wound.

Instructions for use are enclosed with each sachet. It should be used in this manner and will remain effective and pliable.

Contra-indications, Precautions, etc.

- 1) Do not leave Debrisan for more than 24 hours on wounds with a very low exudation rate as it may dry and form a crust which may be difficult to wash off.
- 2) Occlusive dressings may lead to maceration of skin round the wound under treatment.
- 3) When deep infected wounds are treated, care must be taken to wash Debrisan from the depths of the wound.
- 4) No side effects have been reported.

Warning:

Debrisan spillage can render surfaces very slippery. Clear spillages promptly.

Precautions:

- 1) Not to be used on dry wounds.
- 2) When exudate has been markedly removed by Debrisan alternative treatment should be substituted.
- 3) In order to avoid cross-infection it is recommended that the contents of a castor be confined to the treatment of a single patient for one day.

Pharmaceutical Precautions:

Keep in a dry place in well-closed containers.

Packs and Price:

4g sachets in packs of 7 - Basic NHS price £9.75.
Also available in 60g castors.

Legal Category: P.O.M.

Product Licence No.: 0009/0021



Pharmacia

Debrisan (regd) dextranomer is a product of Pharmacia (Great Britain) Ltd Prince Regent Road Hounslow Middlesex TW3 1NE Telephone: 01-572 7321 Further information is available on request to the Company


Creating the healing environment of the future.

By a unique mechanical action Debrisan removes from the wound those substances which normally complicate wound healing, such as bacteria and wound debris.

Pain, oedema and malodour are quickly reduced. Healing can now occur in an ideal environment. Fast.

Use Debrisan on all moist suppurating wounds such as pressure sores and leg ulcers.

Patients can enjoy the benefits of the healing environment of the future. Today.



DEBRISAN BEADS

dextranomer

The molecular way to cleanse wounds; promote healing.

The combination of
simple language and straightforward
line drawings has made the

ABC OF OPHTHALMOLOGY

one of the most popular series
of articles in the Medical Practice section
of the British Medical Journal.


All the common problems are covered –
visual defects, squints, glaucoma and
cataracts – together with detailed
instruction on examination, diagnosis
and management.

The articles
have now been
collected into book
form – in exactly
the same format
as they appeared in
the BMJ –
price: Inland £2.50;
Overseas US\$6.25
(Inland £2.00;
Overseas US\$5.00
for BMA members).


AVAILABLE NOW

Order your copy now

From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller



*Lift the burden
of arthritic pain*



PARAHYPON

Paracetamol B.P. • Codeine Phosphate B.P. • Caffeine B.P.

the pink pain-killer

PRESCRIBING INFORMATION

Indications For relief of pain associated with arthritis, rheumatism and other diseases of the skeletal system, headache, toothache, colds and influenza, and other conditions where simple analgesia is required. **Dosage** Adults and children over 12 years: 1-2 tablets up to four times daily. Children 6 to 12 years: ½-1 tablet up to four times daily. **Contra-indications and Precautions** Contra-indicated in patients with a known sensitivity to any of the constituents. Use with caution in the presence of renal or hepatic dysfunction. Side-effects with PARAHYPON* are rare in therapeutic doses. Paracetamol has been widely used and reports of adverse reactions are rare. Overdosage with paracetamol may cause liver damage. **Presentation** Each PARAHYPON tablet contains: Paracetamol B.P. 500 mg; Codeine Phosphate B.P. 5 mg; Caffeine B.P. 10 mg. The tablets are pink, scored on one side and impressed with the name 'PARAHYPON' on the other. Basic NHS cost: £0.51 for 25 tablets. Additional information is available on request.



DOCTOR AT SEA.

As a qualified doctor you can join the Royal Navy on a 5-year Short Career Commission.

You will have the opportunity of serving in ships, in submarines, or with the Royal Marines Commandos, and in a wide variety of Naval Establishments.

Career counselling will help you plan your future.

There are opportunities for approved General and Higher Professional Training in preparation for careers in general practice and the hospital disciplines. Similar opportunities also exist for training in Naval Occupational and Community Medicine which includes aviation, underwater, submarine, nuclear, preventive and industrial medicine.

If you join immediately after registration your salary will be £7,153 as a Surgeon Lieutenant.

You can, however, join at any age up to 39 when your professional experience is taken into account and you could join as a Surgeon Lieutenant Commander earning £9,183 a year.

There is extra pay for certain recognised post-graduate qualifications and for Specialist and Consultant status.

There is a generous Boarding School Allowance for your children.


If you leave at the end of your 5-year Commission you will receive a tax-free gratuity. You may prefer to apply for extension to eight years or transfer to a pensionable Medium or Full Career Commission.

For more information write to: Surgeon Commander D.J. McKay, LM, LS, MRCP, RN (724M014), Medical Directorate General (Naval), Ministry of Defence, First Avenue House, High Holborn, London WC1V 6HE.



ROYAL NAVY MEDICAL OFFICER

All theory and no practice?



We can help you put that right

Our bank loan scheme is now available to General Practitioners to assist with the purchase of a share in practice premises and assets or the acquisition of new surgery premises. Loans are repayable over periods up to 15 years at an interest of $2\frac{1}{2}\%$ above bank base rate *on the reducing balance*.

For further details apply to:

Sleath

J. W. Sleath & Co. Ltd.,
58 Theobalds Road,
London, WC1X 8SG
Telephone 01-242 4375

Alcohol Treatment Unit

This private unit has been established in the grounds of Ticehurst House to provide comprehensive and enlightened treatment for those with drinking problems. Treatment is structured to individual needs and is undertaken by doctors, nurses and therapists with wide experience in this specialty.

Subscribers to the main Private Contributory Schemes may claim benefits within the terms of these schemes.

Further information is available from the Medical Director,

The Newington Unit

Ticehurst House, Ticehurst, Wadhurst, Sussex.
Telephone: Ticehurst (0580) 200391.



Caved-S[®]

Deglycyrrhizinated Liquorice, Bism.Subnit.
Alum.Hydrox.,Mag.Carb.,Sod.Bic.,Frangula.

heals peptic ulcers

EFFECTIVELY

AND

ECONOMICALLY

Full prescribing information on request from
TILLOTTS LABORATORIES, Henlow Trading Estate, Henlow, Beds.

79004

BONE AND JOINT DISEASES

Arthritis used to be the Cinderella of diseases, but improved diagnostic techniques, advances in joint replacement surgery, and a whole range of non-steroidal drugs have led to a much more active approach to the management of the common bone and joint diseases. Detailed advice on their diagnosis and treatment was given last year in a series of 16 articles in the *BMJ*, and these have now been gathered together in the latest volume based on the popular "Today's Treatment" series.

Demy Octavo

Paperback

131 pages

ISBN 0 7279 0026 9

**Price: Inland £3.00; Abroad US \$7.50,
including postage**

Payment must be enclosed with order or a surcharge of 30p will be made for rendering invoices and statements

Order your copy now from

The Publisher, BRITISH MEDICAL JOURNAL,
BMA House, Tavistock Square, London WC1H 9JR
or through any leading bookseller



**In the only double
blind in vivo trial
CERUMOL®**

An arachis oil base containing turpentine, paradichlorobenzene and chlorbutol.

**was shown to be the
most successful
product in facilitating
ear wax removal[†]**

†J. Laryngology & Otology (1970) LXXXIV No. 10: 1055

The trial included sodium bicarbonate and olive oil.



Further information is available on request from the Company. Laboratories for Applied Biology Ltd, 91 Amhurst Park, London N16 5DR
Cerumol is a trade mark

Do you care for elderly people?

As part of the Scottish Health Education Unit's continuing education programme, the following aids are now available:

Caring with Confidence

A guidance manual for those caring for the elderly. Price outwith Scotland £1.30. Free in Scotland.

Mental Confusion in the Elderly

The first in a series available both on film and VCR designed to give a deeper understanding of the special needs and problems of the elderly. This film is intended primarily for medical undergraduates. Available on free loan or purchase.


The Work of Home-Helps

This training programme consisting of 35 mm slides with accompanying tape or script, is designed for home-helps, voluntary workers and paramedics caring for the elderly. Available on free loan or purchase.

The above aids are available from your area health education officer or write to:

ii SCOTTISH HEALTH EDUCATION UNIT, Dept E, 21 Lansdowne Crescent, Edinburgh EH12 5EH.

Join the search...



...to help us find
the 'lost ostomists'

Abbott Laboratories are holding a series of 'Stoma Care Information Days' around the country. Those meetings arranged for the near future are:-

Stockton – Monday June 11th 10am-8pm
The Conference Room, School of Nursing,
North Tees General Hospital,
Hardwick Estate, Stockton-on-Tees.

Middlesbrough – Wednesday June 13th 10am-5pm
The Cleveland Health Centre,
Cleveland Centre, Middlesbrough.

Ipswich – Wednesday June 13th 10am-8pm
The Health Education Centre,
46A St. Matthews Street,
Ipswich.

If you live in one of these areas you should have already received full information. Other meetings will be announced in future issues of this publication. For information, please contact the Lost Ostomist Campaign Office at Abbott Laboratories – Sheerness (079 56) 3371.

Abbott Laboratories Ltd., Queenborough, Kent ME11 5EL


NOW A RANGE OF THREE




NIHON KOHDEN

cardiofax


ECG's



The ECG – 2101 is a single-channel high performance professional Electrocardiograph designed for mains and re-chargeable battery operation. Floating inputs ensure patient isolation and safety. Ideally suited for the ECG Department, or for carrying around the wards and/or Patient consultations. Proven over the last seven years in use in the United Kingdom.



NEW



NEW, the ECG 5303, a 3-channel Electro-cardiograph designed to meet the regulations for class 11 CF type equipment. Photoc isolation, leakage currents are less than 10 uA.

NEW, the ECG 5101, a one-channel ECG using disposable dry cell batteries, freely available from stores the ECG offers more than 1,100 full 12 lead recordings

Fighting disease with Electronics

HOLSTAR INSTRUMENTS LIMITED

21, West Street, Epsom, Surrey KT17 1UZ

Tel Epsom (037 27) 28315

Published September 1978

Supplement to BRITISH HEART JOURNAL

PREVENTION OF ARRHYTHMIAS

The Proceedings of a Symposium held at Freeman Hospital, Newcastle upon Tyne, on 12 and 13 September 1977, arranged by Professor D G Julian

Contents

Foreword *Dennis Krikler*

Anatomy of conducting tissues revisited *Robert H Anderson and Anton E Becker*

Effects of cardiac denervation on cardiac arrhythmias and electrophysiology *Donald C Harrison, Jay W Mason, John S Schroeder and Edward B Stinson*

Ventricular arrest: a possible mechanism of sudden death *Jan Pool, Anja Roodenburg, Boy Glaser and Jacobus Lubsen*

Catecholamine-induced severe ventricular arrhythmias with Adams-Stokes syndrome in children: report of four cases *Philippe Coumel, Jean Fidelle, Vincent Lucet, Patrick Attuel and Yves Bouvrain*

Surgical treatment of ventricular tachycardia *R A J Spurrell and A J Camm*

Electrophysiology of early ventricular arrhythmias in acute myocardial ischaemia in the pig's heart *Michiel J Janse, André G Kléber, Juan Cinca and Dirk Durrer*

Some factors that influence the activity of antiarrhythmic drugs *E M Vaughan Williams*

Adrenergic influences on arrhythmias due to coronary occlusion and reperfusion *Peter B Corr, Patricia A Penkoske and Burton E Sobel*

Electrophysiological basis for the contrasting prophylactic efficacy of acute and prolonged beta-blockade *A E G Raine and E M Vaughan Williams*

Management in hospital of ventricular fibrillation complicating acute myocardial infarction *K I Lie, K L Liem, and D Durrer*

Clinical pharmacology and antiarrhythmic actions of tocainide *Donald C Harrison, Peter J Meffin, and Roger A Winkle*

Tribulations in trials: experience gained from Nottingham study of beta-adrenergic blockade in acute myocardial infarction *J R A Mitchell*

Review of the present status of drug treatment for preventing sudden death *R H Lovell*

PRICE: Inland £1.25; Abroad U.S. \$3.50, including postage

Payment must be enclosed with order or a surcharge of 30p will be made for rendering invoices and statements

ORDER FROM: The Publisher, BRITISH HEART JOURNAL,
BMA House, Tavistock Square, London WC1H 9JR

Imodium

TRADE MARK

loperamide hydrochloride

made to
measure
treatment for
acute and
chronic
diarrhoea



The tailored dosage means that patients only take Imodium when they need it with no interruption to normal routine. Imodium is easy to take and convenient to carry.

Designed for the specific control of diarrhoea, Imodium relieves pain and abdominal cramps rapidly and effectively.

Imodium
simply
stops diarrhoea fast.

Further information is available from :



Janssen Pharmaceutica
2340 Beerse, Belgium

or from :


Janssen Pharmaceutical Limited,
Janssen House, Marlow, Bucks.

PL 0242/0028 PL 0242/0040

WHY NOT SUBSCRIBE TO THE SPECIALIST JOURNALS PUBLISHED BY THE BMA:—A SELECTION INCLUDES

Archives of Disease in Childhood, British Heart Journal, British Journal of Ophthalmology, Gut, Journal of Clinical Pathology & Journal of Neurology, Neurosurgery & Psychiatry.

FULL DETAILS OF SUBSCRIPTION RATES FROM BMA SPECIALIST PUBLICATIONS: BMA HOUSE, TAVISTOCK SQUARE, LONDON WC1H 9JR
TELEPHONE 01-387 4499 EXT. 309.



Simply right for the job


Lopresor[®]

metoprolol
tartrate for hypertension

Efficacy—Tolerability— β_1 Selectivity—Simple dosage regime:
the essentials of Lopresor simplicity in hypertension



Lopresor prescribing data: Presentation: Lopresor (metoprolol tartrate) as 50mg and 100mg tablets. Indications: Hypertension and angina pectoris. Dosage: *Hypertension*: Initially one 100mg tablet morning and evening. Doses of 100-200mg twice daily may subsequently be prescribed depending upon response. The total daily dose of Lopresor may be given as a single dose if required. Where there has been lack of response to previous therapy, Lopresor may be added to the regimen with the appropriate adjustment of the previous therapy where necessary. *Angina pectoris*: One or two 50mg tablets, twice or three times daily. *Contraindications*: Atrioventricular block, digitalis-refractory heart failure, severe bradycardia, cardiogenic shock. *Precautions*: Lopresor has proved safe in a large number of asthmatic patients; although it is a selective beta-blocker it is prudent to exercise care in the treatment of patients with chronic obstructive pulmonary disease. The dosage of any adrenergic bronchodilators may require adjustment. As with other beta-blockers, Lopresor should not be given to patients with cardiac decompensation unless concomitantly treated with digitalis and/or diuretics. It may be necessary to adjust the dose of the hypoglycaemic agent in labile and insulin-dependent diabetes. Lopresor therapy should be brought to the attention of the anaesthetist prior to general anaesthesia. As a result of animal studies in the rat and rabbit there is no reason to believe that Lopresor is, or maybe, teratogenic in humans. However, the administration of Lopresor during pregnancy, as with other drugs, is advised only if there are compelling reasons. *Side-effects*: Slight gastro-intestinal discomfort and disturbance of sleep pattern occasionally occur. In most cases these effects have been transient, or have disappeared after a reduction in dosage. *Further information*: Because of its cardioselectivity, Lopresor can be administered even to patients with obstructive respiratory disorders, provided adequate precautions are taken. *Legal category*: Prescription only. *Basic NHS price*: 50mg. Packs of 100 tablets £4.20. 100mg. Packs of 100 tablets £7.80. 100mg Calendar Packs of 56 tablets £4.37. *Product licence numbers*: Lopresor 50mg tablets PL0001/0065. Lopresor 100mg tablets PL0001/0066 *Product licence holder*: Geigy Pharmaceuticals, Macclesfield, Cheshire.



**now available in
calendar packs**