

"Since the commencement of hostilities we employ **Neocaine-Surrenine** Corbiere which has given us every satisfaction."

Pouchet and  
Sourdut:  
Regional Anæsthesia.

# NEOCAINE SURRENINE

"Neocaine - Surrenine gives marvellous results in the operation of extraction, dental anæsthesia, in pulpctomy, and cases of periodontitis."

Odontology:  
Professor Choquet

**A PERFECT COCAINE SUBSTITUTE OF LOW TOXICITY.**

**Effective and Safe for LOCAL ANÆSTHESIA, &c.**




FIG. 1.

**NEOCAINE** is a synthetic product of French manufacture. It is in the form of a white powder, belongs to the Benzoyl Dimethyl Ethyl series, and is easily soluble in cold water. **Neocaine possesses all the therapeutical qualities of Cocaine** (except as an exhilarant) **but only one-sixth—one-seventh of its toxicity.**

The anæsthetic power of Neocaine is equal to that of Cocaine; also its duration and rapidity of action.

## COMPOSITION OF NEOCAINE-SURRENINE

Pure Neocaine ... 1 gramme.  
Borated Adrenalin Chlor. (Takamine) 2 milligrammes.

## APPLICATIONS—

Local Anæsthesia by Infiltration—Regional Anæsthesia—  
Rachianæsthesia—Anæsthesia by external application.

**NEOCAINE-SURRENINE may be obtained in:—**

\*5 centigramme doses, each in sterilised double capsule; easily pulled apart (Fig. 1 E).

\*Tube containing 20 5 cg. doses of the powder in capsules.

Box containing 12 tubes of 20 doses each (240 doses).

\* Not suitable for export to hot humid climates; for this purpose powder in tubes or solution in ampoules is recommended.

### TUBES containing—

5 grammes of Powder.  
 $\frac{1}{2}$  ounce " "  
1 " " "

### AMPOULES containing—

Prepared sterilised Solutions ready  
for use—2 %,  $2\frac{1}{2}$  %, 5 %.

**AMPOULES** of 1, 5, 10, 20, 50 and 100 c.c., containing Isotonic Sterile Serum for making extemporaneous solutions.

# ENDOCRISINES

**Complete Extracts of Internal Secretory Glands.**

Prepared at freezing point and *in vacuo*.

Owing to the non-chemical treatment of the glands **the Extracts retain their full activity** and dessication by heat is also avoided.

By drying the organs that have been removed under the best possible conditions of asepsis, *in vacuo* and at the temperature of melting ice, we obtain products that are free from all microbial contamination, and do not differ from fresh glands except in the fact that they have lost their water, their entire activity being preserved because neither heat nor chemical reagents have destroyed these fragile elements.

**THYROIDINE** **OVARINE** **ORCHITINE** **HYPOPHYSIS** **SUPRARENAL**  
(Corpus luteum) (Testicle) (Pituitary)  
**KIDNEY** **MAMMA** **LIVER** **SPLEEN** **THYMUS**

(Full indications for these extracts are contained in the literature.)

# BILÉYL

**A Pure and Stable Extract of Fresh Ox-gall in the Form of Keratin-coated Globules.**

These dissolve in the intestine and then become active.

### INDICATIONS:

As a corrective in Intestinal Putrefaction. Stimulating action on the Biliary Secretion. For Icterus, Biliary Calculus or Colic.

Hereditary Simple Cholæmia, Entero-colitis of the muco-membranous type with Constipation.

Further Literature on application to—

**THE ANGLO-FRENCH DRUG CO., Ltd.** (late M. BRESILLON & Co.),  
GAMAGE BUILDING, HOLBORN, LONDON, E.C.

Telegrams: "AMPSALVAS, LONDON."

Phone: 1311 HOLBORN,