Rational Treatment of CONSTIPATION.

By the double action of secretions and peristalsis.

OPOLAXYL

Opolaxyl is a combination of the secretions of the liver (biliary), pancreas, and intestines, with vegetable extracts of a non-drastic nature.

It combines all the secretions to correspond to nature's therapy, and promotes a flow of bile and glycogen.

It is a normal regulator of the gastro-intestinal functions, consequently it improves the metabolic changes of the entire organism.

Opolaxyl does not lose its efficacy by prolonged use, its effect is gentle and constant, with the least irritation. It is not merely purgative, it stimulates the defective organs and revives the normal equilibrium it has a lasting effect by reason of it rebuilding the organs, thus correcting diminutional function.

Opolaxyl is put up in small size tabular form and should be swallowed without crunching at bedtime or before breakfast.

Dose—For obstinate constipation 2 or 3 tablets, afterwards 1 tablet every 3 or 4 days for a month.

KINECTINE

According to Dr. MOUNEYRAT, the discoverer of Galyl (the safe Salvarsan Substitute).

The SCIENTIFIC TREATMENT of HAY FEVER and allied ailments.

FORMULA:

Chlorhydrate of Quinine c. Hectine—i.e., Benzosulfone-para-amino-phenyl-arsenate of Quinine.

Non-toxic, produces no ill-effects. Easily taken (tablets) and well tolerated.

Not only a Prophylactic against, but, a specific in

HAY FEVER, Catarrh, INFLUENZA, &c.

Rapid Abortive Action in the Initial Stages.

Chronic Conditions

Chronic Conditions.

DEAR SIRS,

For the first time for 18 years I have escaped my annual attack of Influenza, thanks to Kinectine. Please send me further supply. I enclose cheque 30s."—Yours faithfully,

Adult Dose: 1 tablet three times a day for three days, then 2 tablets every other day until 18 tablets have been taken. Half doses for Children.

The Most Powerful and Effective Urinary Antiseptiq,

URASEPTINE

URASEPTINE is a granulated product entirely soluble in water, its bases being Piperazine, Urotropine, Helmitol, Benzoates of Sodium and Lithium. It contains 60 centigrams (10 grs.) of active matter to each teaspoonful.

It purifies the Urine, and this action is due to its three principal properties:

- I. It is a URINARY ANTISEPTIC
- 2. A SOLVENT of URIC ACID and of PHOSPHATES.
- 3. A MILD NON-TOXIC DIURETC.

INDICATIONS. — Arthritism, Gout, Giavel, Hepatic and Renal Colic, Rheumatism, Calculus, &c., Phosphaturia, Urinary Antisepsis, Pyelitis, Bacteriuria, Cystitis, Prostatitis, Urethritis, Puyuria, Urinary Abscess, Vesical Catarrh, &c.

Dose: 2 to 6 teaspoonfuls daily.

STANNOXYL

ROBERT & CARRIÈRE, PARIS.

(An Oxide of Tin and Tin Metal Free from Lead.)
The treatment of CARBUNCLES, BOILS,
ANTHRAX, ACNE, STYES, and Diseases
arising from Staphylococcus.

A scientific production the value of which has been studied very closely. The effect is very prompt; from the second day of treatment the pain is relieved, the carbuncles begin to dry up, and those which are just opening are stopped in their course; the core is not expelled but reabsorbed.

In the majority of cases definite results are obtained within five or six days; it is seldom necessary to take the full ten days' treatment, and relapses are unknown.

Vide THE LANCET, Jan. 19th, 1918, pp. 99 and 109.

"March, 1918.
GENTLEMEN, I have not found anything to equal Stannoxyl in the treatment of Acne Vulgaris. In the cases I have used it, the results have been excellent. I have tried it in a few cases of Skin infection with encouraging results.—Yours faithfully,

The daily dose for Adults is 4 to 8 tablets; Children 2 to 4 tablets.

Supplied in vials of 80 tablets.

LITERATURE, CLINICAL REPORTS and PRICE LISTS on REQUEST.

The ANGLO-FRENCH DRUG Co., Ltd., Gamage Building, Holborn, London, E.C. 1.

Telephone: Holborn 1311.

Telegrams: "AMPSALVAS, LONDON."

WEST END DEPOT:-MODERN PHARMACALS, 48, Mortimer Street, W. 1. Telephone: Museum 564.