

URASEPTINE

A
URINARY
ANTISEPTIC
AND MILD
NON-TOXIC
DIURETIC.

A granular compound of
Piperazine, Hexamine, Helmitol,
etc.

Dose.—Three to six teaspoonfuls daily.



In response to numerous enquiries, we have pleasure in informing the Medical Profession that we can now supply

STANNOXYL in Liquid Form.



EXTERNAL

APPLICATION.

This is a solution of Protochloride of Tin in Glycerine, to be diluted (1 in 40) with water for use as a lotion preferably, in conjunction with the Tablets.

In bottles containing 2 ozs. of solution.

Tested and Approved in Accordance with
L.G.B. requirements.

INTRAVENOUS.

GALYL

IS THE
SAFEST OF
ALL ARSENICAL
COMPOUNDS

IN

SYPHILIS

INTRAMUSCULAR.

Vide Presidential Address, British Pharmaceutical
Congress, 10th July, 1918.

This preparation of

CRYSTALLISED DIGITALINE ÉCALLE

is standardised according to the Keller process,
modified by Écalle, and gives the best guarantee
to the Practitioner.



PURITY.

SECURITY.

In bottles of 60 granules of 1/10 mgr. of
Digitaline.

LATAPIE REACTION

*Simplified Wassermann Test according
to the Latapie Method of the
Pasteur Institute.*

LITERATURE, CLINICAL REPORTS and PRICE LISTS on REQUEST.

The **ANGLO-FRENCH DRUG Co., Ltd.**, Gamage Building, Holborn. London, E.C. 1.

Telephone: HOLBORN 1311.

NEW YORK: 1270 BROADWAY.

MONTREAL: 330a ST. ANTOINE ST.

Telegrams: "AMPSALVAS, LONDON."

PARIS: 5, RUE CLAUZEL.

WEST END DEPOT:—MODERN PHARMACALS, 48, Mortimer Street, W. 1. Telephone: MUSEUM 564.