

"OXOID"

The word "OXOID" is the trade mark of OXO Ltd., and is used in connection with their therapeutical preparations both in tablet and fluid extract form.

ORGANO-THERAPEUTICAL PRODUCTS

OXO LABORATORY PREPARATIONS

VITAMINS

"OXOID" Brand

VITAMINS 'A' & 'D'

Capsules

A safeguard against infection and a prophylactic against the Common Cold. Also prescribed for rapid adaptation of vision in semi-darkness. Supplies anti-rachitic or "Sunshine" vitamin during Pregnancy and Lactation, or for Infants.

In tins of 25 Capsules. Also in bottles of 250, 500 and 1,000.

"OXOID" Brand

VITAMIN 'B'

Tablets and Ampoules

For the treatment of Polyneuritis, Neuralgia, Sciatica, Nervous Debility, Loss of Appetite, Gastric Intestinal Disorders, &c. Specially useful in cases of B₁ deficiency in Pregnancy and Lactation.

In bottles of 25, 100, 250 and 500 tablets of 1 mg. and 3 mg. Also supplied in ampoules of 5 mg. and 25 mg.

Literature supplied on application.

OXO LIMITED, Thames House, London. E.C.4


OXO LABORATORY PREPARATIONS

The rapid enzymic action of BENGER'S FOOD


The instant hot milk is added to Benger's Food, self-digestive action by natural pancreatic enzymes begins.




As supplied in the tin, Benger's Food is a special wheaten powder, containing 77% of insoluble starch.



So quick is the enzymic action of Benger's Food that within 2 minutes of adding the hot milk insoluble starch has been reduced to 0.5%.



The milk proteins have simultaneously been modified, so that when prepared Benger's Food comes in contact with the gastric juices it separates into fine flocculi instead of the heavy tough curds formed by milk itself. All this change in two minutes! And by the time the milk used to make Benger's Food is sufficiently cool to drink, the self-digestion of Benger's Food is carried as far as it need be for all cases where digestion is slightly impaired or where light diet is indicated.



For Infants, Invalids and those suffering from severe digestive weakness, the time for pre-digesting Benger's Food should be extended to 15 minutes and upwards to 45 minutes. At the end of this period the wheaten powder is converted to dextrins, dextri-maltoses and maltose, and the milk proteins modified so that the finest possible curd is formed on contact with the gastric juices.

Sole Manufacturers: **BENGER'S LTD.**,

Holmes Chapel, Cheshire, Eng.

The Benger Laboratories are always at the disposal of the medical profession in any of their dietetic problems. Physicians' samples of Benger's Food are always available.