

'ROCHE' News in Brief

Vitamin C Saturation in Hospital Children

From the results of an extensive investigation recently completed at the Royal Hospital for Sick Children, Glasgow, the authors concluded that "while full saturation may not be necessary for the maintenance of health, it appears to us that children living in slums where acute infections are so prevalent should have ample supplies of vitamin C, because with the onset of infection their reserves tend to become exhausted. Similarly, in the medical wards of children's hospitals, precautions should be taken to ensure that ample vitamin C in one form or another is given to all patients." (Glasgow Med. Journ., 1944, 24, p. 55.)

'Prostigmin' and Blood Clotting

A heparin tolerance test has recently been devised which is considered to be valuable in showing delicate changes in the blood clotting mechanism. By means of this test the author has shown that autonomic stimulation shortens the coagulation time whilst vagal stimulation prolongs it. 'Prostigmin' was found to raise the heparin curve, and when given post-operatively in gynaecological cases prevented the flattening of the curve seen during the first 2 to 4 days after operation when the danger from thrombosis is greatest. "Prostigmin" in our experience has improved the clotting tendency of the post-operative state and is useful to combat post-operative atony of the intestines and bladder." (Arch. Surg., 1944, 48, p. 105.)

Biological Activity of the Tocopherols

The vitamin E fraction of wheat-germ oil presents a mixture of alpha, beta and gamma tocopherols while the synthetic tocopherol preparation consists only of the alpha isomer. In a recent paper in the *Journal of Nutrition* (25 (5), 433-440, 1943) entitled "Biological Determination of Vitamin E," it is reported that—according to feeding tests—synthetic alpha, beta and gamma tocopherols and α -tocopheryl acetate have the relative biological activities of 100, 25, 19 and 100 respectively. Rats on a low vitamin E diet gained weight during pregnancy in direct proportion to the amount of α -tocopherol given. These tests point to the superior action of α -tocopherol over the beta and gamma isomers respectively.

Sulphonamide Sensitivity

Striking results from the use of ascorbic acid to relieve the effects of sensitivity to sulphapyridine are reported by Schropp. In a case of pneumonia in a child the first dose of sulphapyridine produced herpes labialis, soreness of the mouth and eyes, soreness and bleeding of the gums and vesicles on the mucous membrane of the lip. A 50-mg. dose of ascorbic acid given with each subsequent dose relieved all these side-reactions, and 18 hours after the first dose of ascorbic acid the temperature had fallen, and the soreness of the mouth and gums had disappeared. (Canad. Med. Assoc. Journ., 1943, 49, p. 515.)

'Allonal'
CONTROLS FEAR...
Relieves pain and counteracts
toxic effects of local anaesthetics

Each 'Allonal' tablet contains allyl-isopropyl-barbiturate of amidopyrine 0.126 gm. (approx. gr. 2), amidopyrine B.P. 0.034 gm. (approx. gr. $\frac{1}{2}$). Issued in bottles of 12, 100, 500, and larger packings.

ROCHE PRODUCTS LTD.
WELWYN GARDEN CITY, ENGLAND

THE THERAPY OF ASTHMA



The treatment of asthma resolves itself into a consideration of underlying factors and causes. Often in ASTHMA the underlying cause is not discoverable or changes from time to time—now irritant dusts, now bacterial infection, etc. The underlying factor is fortunately always the same—bronchospasm.


Thus sometimes causative agents can be removed or mitigated but always the underlying factor—bronchospasm—can be treated, successfully, with FELSOL.

Most cases of Asthma are chronic and demand patience in treatment—persistence with FELSOL will yield the highest possible percentage of successes.

NO MORPHIA . . . NO NARCOTICS

Phenazone 0.47 Iodpyrine 0.03
Antipyrine Aceto-salicylas 0.40
Caffeine 0.10
Ext. Viscum Album 0.01
Ext. Brachycardia 0.01
Physicians' samples and literature willingly sent on request

BRITISH FEL SOL COMPANY LTD., 206/212 St. John St., London, E.C.1
Telephone: Clerkenwell 5862. Telegrams: Felsol, Smith, London



MYALGIA AND FEBRILE CONDITIONS

MUSCULAR aches and pains and the feverish conditions which frequently accompany the common cold can be effectively eased by the administration of 'ANADIN.' Because of its balanced formula of aspirin, phenacetin and caffeine, 'ANADIN' can be prescribed with the assurance that it is well-tolerated over prolonged periods.

Dose: Two tablets at four-hourly intervals.
Formula: Acetophenetidin 2 gr.; Ac. Acetyl salicylic. 4 gr.; Caffeine Alkaloid. $\frac{1}{2}$ gr.

ANADIN Tablets

ANADIN LIMITED, 12 CHENIES STREET, LONDON, W.C.1