First Impression 1974 Reprinted 1978

## MEDICINE IN OLD AGE

The series of twenty two articles published in the *British Medical Journal* have been collected in book form as a convenient source of reference for general practitioners. The emphasis is on remedial mechanisms, and the use that can be made of psychogeriatric and social services in keeping the elderly healthy and relatively independent members of the community.

Demy Octavo

Paperback
ISBN 0 900221 93 3

175 pages

Price: Inland £2.00; Abroad US\$5.00 including postage

Payment must be enclosed with order or a surcharge of 30p will be made for rendering invoices and statements

## ORDER YOUR COPY NOW

From: The Publisher, British Medical Journal, BMA House, Tavistock Square, London WC1H 9JR or through any leading bookseller

## Inocium TRADE MARK TRADE MARK

loperamide hydrochloride

made to measure treatment for acute and chronic diarrhoea

The tailored dosage means that patients only take **Imodium** when they need it with no interruption to normal routine. **Imodium** is easy to take and convenient to carry.

Designed for the specific control of diarrhoea, **Imodium** relieves pain and abdominal cramps rapidly and effectively.

Further information is available from :



Janssen Pharmaceutica 2340 Beerse, Belgium Imodium simply stops diarrhoea fast.

or from:


Janssen Pharmaceutical Limited, Janssen House, Marlow, Bucks.

PL 0242/0028 PL 0242/0040

WHY NOT SUBSCRIBE TO THE SPECIALIST JOURNALS PUBLISHED BY THE BMA:—A SELECTION INCLUDES

Archives of Disease in Childhood, British Heart Journal, British Journal of Ophthalmology, Gut, Journal of Clinical Pathology & Journal of Neurology, Neurosurgery & Psychiatry.

FULL DETAILS OF SUBSCRIPTION RATES FROM BMA SPECIALIST PUBLICATIONS: BMA HOUSE, TAVISTOCK SQUARE, LONDON WC1H 9JR TELEPHONE 01-387 4499 EXT. 309.



Injection (4 ml and 25 ml ampoules; each ml contains 0.25 mg bumetanide) and Burinex K (0.5 mg bumetanide and 7.7 mmol slow release KCI). Dosage: oral and parenteral normally 1 mg. Higher doses may be used in resistant oedema, for long-term maintenance therapy of CCF, Burinex K is usually given as 2 tablets O.M. Contra-indications: Hepatic coma and severe electrolyte disturbances. Burinex K should not be used with potassium sparing agents. Adverse reactions: Occasional skin rashes and minor gastric upset. Precautions: Burinex may potentiate antihypertensive therapy and rarely, may affect carbohydrate metabolism. Periodic electrolyte checks are advisable on long-term therapy, Burinex K should be stopped immediately if signs or symptoms of bowel ulceration appear. Product Licence No: 0043/0027B

