


ANOTHER INNOVATION FROM JANSSEN IN THE TREATMENT OF VAGINAL THRUSH.

Rapid release of
active fungicidal
Gyno-Daktarin
emulsion coupled
with ...



... absorption of
discharge and no leakage
of medication.

Gyno-Daktarin* *tampons* (miconazole)

Highly acceptable and effective short course therapy for
candidal vaginitis.



Also available : Gyno-Daktarin cream, ovules and pessaries.

N.B. Not all product forms available in all countries.

Further information available on request from :

JANSSEN PHARMACEUTICAL Ltd.  JANSSEN PHARMACEUTICA N.V.
Janssen House, Marlow, Bucks. SL7 1ET 2340 Beerse, Belgium

* Trademark

© Janssen Pharmaceutica 1980

PL 460/80

WHY NOT SUBSCRIBE TO THE SPECIALIST JOURNALS PUBLISHED BY THE BMA:—A SELECTION INCLUDES

Archives of Disease in Childhood, British Heart Journal, British Journal of Ophthalmology, Gut, Journal of Clinical Pathology & Journal of Neurology, Neurosurgery & Psychiatry.

FULL DETAILS OF SUBSCRIPTION RATES FROM BMA SPECIALIST PUBLICATIONS: BMA HOUSE, TAVISTOCK SQUARE, LONDON WC1H 9JR
TELEPHONE 01-387 4499 EXT. 309.

PARDALE FOR PAIN RELIEF
PARDALE FOR PAIN RELIEF
PARDALE FOR PAIN RELIEF
PARDALE FOR PAIN RELIEF
PARDALE FOR PAIN RELIEF

Paracetamol 400mg Codeine Phosphate 9mg
Caffeine Hydrate 10mg

Full product information available on request



Dales Pharmaceuticals Limited

Barrows Lane, Steeton, Keighley, Yorkshire BD20 6PP
 (Steeton 53222) PL 0123 5015



CARDIO-MINI
MICRO-CONSTRUCTION

- * All the features you ever wanted in a Electrocardiograph.
- * Demonstrations & Brochures on request.

Cardiacare Instruments (UK) Ltd.
 48 HALL ROAD, ROMFORD, ESSEX RM6 4LJ
 TEL: 01-597 3099 TELEX: 24924 REF: 2801

Cut out car tax.

You could save hundreds of pounds by buying a tax-free Audi or Volkswagen if you're a visitor to Britain, a British resident shortly going abroad, a NATO Serviceman, a member of British Forces or a Diplomat.

Just talk to our Personal Export experts.

Fill in the coupon or ring 01-486 8411.



Personal Export Sales

Personal Export Sales Volkswagen (GB) Limited
 95 Baker Street London W1M 1FB

Name

Profession

Address

Tel. No. Home Business

I am interested in:

For use in: (Country)

BMJ/6

FREE Radio/Cassette on every car delivered before 28th April 1980

Notice to Subscribers

"BMJ"—Issues Published out of Sequence

Owing to recent industrial action taken by a section of the printing industry subscribers will have experienced problems in receiving their *British Medical Journal*.

Unfortunately the issue dated 5 April 1980 has not yet been published, but it is hoped to produce this by mid-June. The issues dated April 12 to May 31, 1980 were published outside the United Kingdom and therefore considerable delay and irregularity in despatch occurred. The June 7 issue was unaffected by delays.

We are pleased to announce that the dispute is now settled and normal despatch has been resumed.

We apologise for the trouble and inconvenience you have been caused.

The combination of simple language and straightforward line drawings has made the

ABC OF OPHTHALMOLOGY


one of the most popular series of articles in the Medical Practice section of the British Medical Journal.

All the common problems are covered - visual defects, squints, glaucoma and cataracts - together with detailed instruction on examination, diagnosis and management.

The articles have now been collected into book form - in exactly the same format as they appeared in the BMJ - price: Inland £2.50; Overseas US\$6.25 (Inland £2.00; Overseas US\$5.00 for BMA members).

AVAILABLE NOW

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller



FIGHT BACK AGAINST CANCER

It is good to remember that most people live their lives untouched by any form of cancer.

But as all too many are aware, cancer is something that casts its shadow far beyond those it directly affects. That is why so many people think it right to help the urgent work of the Imperial Cancer Research Fund.

From our discoveries in the past has come much of today's hope for sufferers. To go forward with our research for future alleviation, we ask *your* help in the present.



The main laboratories at Lincoln's Inn Fields

IMPERIAL CANCER RESEARCH FUND



*Donations will be most gratefully received by the Appeals Secretary,
Room 23 P.O. Box 123, Lincoln's Inn Fields, London WC2A 3PX*


Halcion[®]

(triazolam)

3010 Patient UK Study

Treatment for Insomnia Under Monitored Release Study Protocol


HALCION PERFORMANCE



Performance

Halcion was shown to be as effective at the end of therapy as at the beginning with an 85% level of efficacy, thus demonstrating no significant development of tolerance. 90% of 'new' patients found Halcion effective. Dosage recommendation was 0.25mg (normal adult dose) or 0.125mg (geriatric patients) taken at bedtime.


HALCION PREFERENCE



Preference

Of 1299 patients who took a hypnotic previously, 811 preferred Halcion. Halcion was significantly preferred to nitrazepam, flurazepam, temazepam and diazepam.

HALCION SAFETY



Safety

Halcion was well tolerated and side effects generally were not severe. Most commonly reported side effects included drowsiness 2.2%, headache 1.8% and dizziness 1.7%.

Prescribing Information: Indications: For treatment of insomnia including anxiety-induced insomnia. **Dosage and Administration:** Usual adult dose one 0.25mg tablet at bed-time. For geriatric patients initial dose is 0.125mg which may be increased to 0.25mg if necessary. Prolonged administration is not necessary or recommended. **Contraindication:** Known hypersensitivity to benzodiazepines. **Precautions, Warnings:** Concomitant administration with CNS depressants including alcohol may result in accentuation of their effect. Safety for use during pregnancy has not yet been established. Caution should be exercised in treating depressed patients. **Side Effects:** Drowsiness, dizziness, amnesia and impaired co-ordination may occur. Side effects are usually dose-related. **UK Availability:** 0.25mg powder blue tablets and 0.125mg lavender tablets in bottles of 100. **Product Licence Nos:** 0.25mg 0032/0058. 0.125mg 0032/0063. Information adapted from Triazolam Monitored Release Study (UK) January to September 1979. Data available from Upjohn Limited.



REGISTERED TRADEMARK: HALCION

UPJOHN LIMITED · CRAWLEY · WEST SUSSEX

Data Sheet available on request 1800 UK

NATRILIX[®]

indapamide

a primary antihypertensive agent

Natrilix offers a simple, effective means of controlling blood pressure both in newly diagnosed and in poorly controlled patients with mild to moderate hypertension

Prescribing information

Presentation

Pink biconvex sugar-coated tablets each containing 2.5 mg indapamide hemihydrate.

Uses

For the treatment of hypertension.

Dosage and administration

Adults


The dosage is one tablet daily to be taken in the morning.

Children

There is no experience of the use of this drug in children.

Contra-indications, warning, etc.

There are no absolute contra-indications to the use of Natrilix but caution should be exercised when prescribing it in cases of severe renal or hepatic impairment. As with all new drugs, the administration of Natrilix should be avoided during pregnancy although no teratological effects have been seen in animals. At doses higher than that recommended, Natrilix has a diuretic effect, therefore it is not recommended to prescribe it with a diuretic agent which may cause hypokalaemia. Also, slight weight loss has been reported in some patients taking Natrilix. Reported side effects have included nausea and headache,



but they are generally uncommon and mild in nature.

Serum urate levels may rise slightly but there is no evidence that glucose tolerance is adversely affected.

Box of 30 tablets.

Product licence number 0093/0022.

NATRILIX is available as:
FLUDEX[®] in Africa, Belgium, France, Holland, Portugal, Switzerland.
NATRILIX[®] in English speaking Countries of Africa, Brazil, Central America, Germany, Ireland, Italy, Middle East, South East Asia, United Kingdom.
TERTENSIF[®] in Spain.
NATRIX[®] in Korea.

Further information is available on request from
Servier Laboratories Ltd.,
Servier House, Horsenden Lane
South, Greenford, Middx.
UB6 7PW.



Les Laboratoires Servier,
22, rue Garnier - 92201 Neuilly -
France.

one tablet daily

WHY NOT SUBSCRIBE TO THE SPECIALIST JOURNALS PUBLISHED BY THE BMA:—A SELECTION INCLUDES

Annals of the Rheumatic Diseases, British Journal of Venereal Diseases, Thorax, Journal of Epidemiology and Community Health, British Journal of Industrial Medicine.

FULL DETAILS OF SUBSCRIPTION RATES FROM BMA SPECIALIST PUBLICATIONS: BMA HOUSE, TAVISTOCK SQUARE, LONDON WC1H 9JR
TELEPHONE 01-387 4499 EXT. 309.



Selective

*
*'Very few lower gastrointestinal disorders occurred. This, in all probability, reflects its selective antibacterial activity which limits drug disturbance of the normal colonic and vaginal flora.'*³

Effective

*'In our clinical trials mecillinam achieved a cure-rate of over 90% at 2 weeks in patients with urinary-tract infections.'*¹

*'The original bacterial strain in the urine was eradicated in 100% of the cases.'*²

Low Toxicity

*'Mecillinam has a novel mode of action but shares the penicillins' low toxicity.'*⁴

*'... the data establish that pivmecillinam is a safe and potent drug in the treatment of urinary tract infection.'*⁵

References

1. Rumfitt, W., and Hamilton-Miller, J.M.T., *Lancet*, 1, 445, 1978.

2. Ekberg, M. et al., *Scand. J. Infect. Dis.*, 10, 127, 1978.

3. Bresky, B., *J. Antimicrob. Chemother.*, 3 (Suppl. B), 121, 1977.

4. Geddes, A.M. and Wise R., *J. Antimicrob. Chemother.*

(Suppl. B), iii, 1977.

5. Verrier Jones, E.R. and Asscher, A.W., *J. Antimicrob.*

Chemother., 1, 193, 1975.

SELEXID Prescribing Information Available as Selexid tablets and Suspension. Each tablet contains 200mg pivmecillinam hydrochloride; each 5ml dose sachet of suspension contains 100mg pivmecillinam as white granules to be dissolved in a small quantity of water. Tablets and suspension should be taken with a meal. **Dose for Urinary Tract Infections:** Tablets and suspension: From 200mg 8 hourly - 400mg 6 hourly depending on severity. For children weighing less than 40kg: 20-40mg/kg bodyweight daily in 3-4 divided doses. **Contra-indications:** Penicillin or cephalosporin hypersensitivity. **Precautions/Adverse Reactions:** In accordance with normal practice Selexid should be avoided in the 1st trimester of pregnancy.

Dosage may need modification when renal function is impaired. Routine monitoring of liver and renal function advisable with prolonged therapy. Gastro-intestinal upset, rash, and anaphylaxis may occur. Full information is available on request. **Price:** £2.82 for a five day course.

Product Licence No.: 0043/0048.

 Leo Laboratories Limited, Hayes Gate House, Hayes, Middlesex

*Selexid is a trade mark