ANOTHER INNOVATION FROM JANSSEN

IN THE TREATMENT OF VAGINAL THRUSH.

Rapid release of active fungicidal Gyno-Daktarin emulsion coupled with ...



... absorption of discharge and no leakage of medication.

Gyno-Daktarin tampons (miconazole)

Highly acceptable and effective short course therapy for candidal vaginitis.



Also available: Gyno-Daktarin cream, ovules and pessaries. N.B. Not all product forms available in all countries.

Further information available on request from :

JANSSEN PHARMACEUTICAL Ltd. 💢 JANSSEN PHARMACEUTICA N.V.



Janssen House, Marlow, Bucks. SL7 1ET 🥏 2340 Beerse, Belgium

* Trademark

© Janssen Pharmaceutica 1980

WHY NOT SUBSCRIBE TO THE SPECIALIST JOURNALS PUBLISHED BY THE BMA:—A SELECTION INCLUDES Archives of Disease in Childhood. British Heart Journal, British Journal of Ophthalmology, Gut, Journal of Clinical Pathology & Journal of

Neurology, Neurosurgery & Psychiatry FULL DETAILS OF SUBSCRIPTION RATES FROM BMA SPECIALIST PUBLICATIONS: BMA HOUSE, TAVISTOCK SQUARE, LONDON WC1H 9JR

TELEPHONE 01-387 4499 EXT. 309.



Test strip for the semiquantitative determination of blood glucose in the range of 20–800 mg/100 ml (1–44 mmol/l)

Main features:

- The combination of two test patches on one test strip ensures a wide range of response and reliable results – no meter required
- Readily identifiable comparison colours for
 20 40 80 120 180 240 400 800 mg/l00 ml
 1 2 4 7 10 13 22 44 mmol/l
- Simple to use no need to rinse with water
- Stable at room temperature
- A long shelf-life even after vial has been opened Stable reaction colour – small timing errors do not lead to large reading errors

For more information please contact:
UK and Ireland
BCL, Bell Lane, Lewes
East Sussex BN 71 LG
Telefon: Lewes (07916) 71611

other countries your Boehringer Mannheim agent or



Boehringer Mannheim GmbH B-8800 Mannheim 31 W. Germany



Out of the shadows of depression and irrational fear with **Anafranil*10 mg**

(clomipramine hydrochloride)

Geigy

Anafranil (clomipramine hydrochloride). Dosage: Depression – 30–50mg per day. In severe cases increase to 75mg or higher if necessary. Phobic/Obsessional Statesgenerally higher dosage—build up to 100–150mg per day. Contra-indications: 1) Anafranil is contra-indicated in conjunction with, or within 14 days of treatment with a monoamine oxidase inhibitor. 2) In existing liver damage. 3) In cardiac/circulatory failure or following a recent myocardial infarction. Precautions: Caution is necessary in schaemic heart disease, glaucoma, and retention of urine. Great caution in patients with epilepsy. Anafranil may antagonise the action of certain antihypertensive drugs, and may potentiate the cardiovascular action of noradrenaline and adrenaline. As with other drugs, avoid using during the first trimester of pregnancy if possible. Side effects: Dry mouth, sweating, difficulty with accommodation, constipation, disturbance of micturition, tachycardia. Less commonly tremor, ataxia, orthostatic hypotension and vertigo. More rarely anxiety, agitation, fatigue, drowsiness, confusion, insomnia and headache. Epileptiform seizures have occurred in a small number of patients. Symptomless hypothermia and allergic skin reactions are extremely rare. Interference with sexual function, particularly ejaculation, may be encountered. Agranulocytosis has occasionally occurred in patients on tricyclic antidepressants – routine blood counts are advised during treatment with Anafranil. Availability: Two-tone yellow/caramel capsules of 10mg (PL0001/0037) £3.10 per 100. Two-tone orange/caramel capsules of 25mg (PL0001/5000) £6.11 per 100. Two-tone blue/caramel capsules of 50mg (PL0001/0068) £11.63 per 100. Orange flavoured and coloured syrup containing 25mg of Anafranil in 5ml (PL0001/5001) £5.55 per 150ml bottle. Full prescribing information is available on request from Geigy Pharmaceuticals, Horsham, West Sussex.