


Imodium

TRADEMARK

(loperamide hydrochloride)

symbol of
accuracy
in acute diarrhoea



Imodium's specific action
controls diarrhoea, relieving pain and
abdominal cramps rapidly and effectively.

A tailored dosage regime means that patients only take Imodium when they really need it, minimising the risk of constipation and encouraging economy of medication. Patients will find Imodium capsules easy to take and the blister pack of just twelve capsules is convenient to carry throughout the working day.

Imodium
Simply Stops Diarrhoea-fast
2 stat and 1
at each loose stool

Further information is available from :



JANSSEN PHARMACEUTICA
2340 Beerse, Belgium

JANSSEN PHARMACEUTICAL LTD.
Marlow, Bucks. SL7 1ET.

Presentation :

Hard gelatin capsules containing 2 mg loperamide hydrochloride and syrup containing 1 mg loperamide hydrochloride per 5 ml.

Indications :

Imodium* is indicated for the symptomatic control of acute diarrhoea of any aetiology.

Contra-indications and warnings etc. :

There are no specific contraindications to Imodium*. Studies in animals have shown no abnormal teratogenic effects, however the use of loperamide during pregnancy is subject to the usual precautions. In trials, no side effects have been reported that can reliably be distinguished from the symptoms of the gastro-intestinal disorder being treated. As persistent diarrhoea can be an indicator of potentially more serious conditions, Imodium* should not be used for prolonged periods until the underlying cause of the diarrhoea has been investigated.

Dosage and Administration :

Acute Diarrhoea : *Adults* : Two capsules initially, followed by one capsule after every loose stool. The usual dosage is 3 to 4 capsules a day; the maximum daily dose should not exceed 8 capsules. *Children* : 4 to 8 years : Syrup 5 ml four times daily until diarrhoea is controlled. In children under 8 years, further investigation may be necessary if diarrhoea has not responded to three days' treatment. 9 to 12 years : Syrup 10 ml (or 1 capsule) four times daily until diarrhoea is controlled.

Basic NHS Cost :

12 capsules (ex 25 pack) 104p. (correct at time of printing)

Product Licence Numbers :

Capsules 0242/0028 · Syrup 0242/0040

NATRILIX®

indapamide

a primary antihypertensive agent

Natrilex offers a simple, effective means of controlling blood pressure both in newly diagnosed and in poorly controlled patients with mild to moderate hypertension

Prescribing information

Presentation

Pink biconvex sugar-coated tablets each containing 2.5 mg indapamide hemihydrate.

Uses

For the treatment of hypertension.

Dosage and administration

Adults

The dosage is one tablet daily to be taken in the morning.

Children

There is no experience of the use of this drug in children.

Contra-indications, warning, etc.

There are no absolute contra-indications to the use of Natrilex but caution should be exercised when prescribing it in cases of severe renal or hepatic impairment. As with all new drugs, the administration of Natrilex should be avoided during pregnancy although no teratological effects have been seen in animals. At doses higher than that recommended, Natrilex has a diuretic effect, therefore it is not recommended to prescribe it with a diuretic agent which may cause hypokalaemia. Also, slight weight loss has been reported in some patients taking Natrilex. Reported side effects have included nausea and headache.



but they are generally uncommon and mild in nature.

Serum urate levels may rise slightly but there is no evidence that glucose tolerance is adversely affected.

Box of 30 tablets.
Product licence number
0093/0022.

NATRILIX is available as:
FLUDEX® in Africa, Belgium,
France, Holland, Portugal,
Switzerland.
NATRILIX® in English
speaking Countries of Africa,
Brazil, Central America,
Germany, Ireland, Italy,
Middle East, South East Asia,
United Kingdom.
TERTENSIF® in Spain.
NATRIX® in Korea.

Further information is available
on request from
Servier Laboratories Ltd.,
Servier House, Horsenden Lane
South, Greenford, Middx.
UB6 7PW.



Les Laboratoires Servier,
22, rue Garnier - 92201 Neuilly -
France.

one tablet daily



WHY NOT SUBSCRIBE TO THE SPECIALIST JOURNALS PUBLISHED BY THE BMA:—A SELECTION INCLUDES

Annals of the Rheumatic Diseases, British Journal of Venereal Diseases, Thorax, Journal of Epidemiology and Community Health, British Journal of Industrial Medicine.

FULL DETAILS OF SUBSCRIPTION RATES FROM BMA SPECIALIST PUBLICATIONS: BMA HOUSE, TAVISTOCK SQUARE, LONDON WC1H 9JR
TELEPHONE 01-387 4499 EXT. 309.



VOLTAROL® 50

diclofenac sodium

The symbol of a new flexibility

Eases rheumatic pain and stiffness

Improves mobility

Flexible dosage

Geigy

Presentation Yellow, biconvex enteric coated tablets of 25mg diclofenac sodium, imprinted Geigy on one side. Light brown, biconvex enteric coated tablets of 50mg diclofenac sodium, imprinted Geigy on one side. Yellowish-white suppositories of 100mg diclofenac sodium. **Mode of action** Voltarol is a non-steroidal agent with marked analgesic/anti-inflammatory properties. **Indications** Rheumatoid arthritis, osteoarthritis. **Dosage** Oral: 25-50mg three times daily initially. Later, 50mg twice daily or 25mg three times daily may be sufficient. Rectal: One suppository daily which may be combined with tablets. The daily dose should not exceed 150mg. **Contra-indications** Peptic ulceration. Patients in whom attacks of asthma, urticaria or acute rhinitis are precipitated by aspirin or other non-steroidal anti-inflammatory agents. **Precautions** Voltarol should not be prescribed during pregnancy, unless there are compelling reasons for doing so. Patients with a history of peptic ulcer, haematemesis, or melena, or with severe hepatic or renal insufficiency, should be kept under close surveillance. Voltarol has been reported to depress salicylate levels, and vice versa; the clinical relevance of this phenomenon is not yet clear. **Warnings and side-effects** Transient epigastric pain, nausea and diarrhoea, headache and slight dizziness have been reported as occasionally have skin rash, peripheral oedema and abnormalities of serum transaminases. **NHS price** Tablets 25mg: pack of 100 - £9.00; 50mg: pack of 100 - £17.50; Suppositories: pack of 10 - £2.98. **Product licence numbers** Tablets 25mg PL0001/0036, 50mg PL0001/0082, Suppositories 100mg PL0001/0083. **Product licence holder** Geigy Pharmaceuticals, Horsham, West Sussex. Full prescribing information is available from Geigy Pharmaceuticals, Horsham, West Sussex.