



Opticrom lets her forget her allergic conjunctivitis.


Opticrom has already proved invaluable to patients suffering from hay fever conjunctivitis and vernal kerato-conjunctivitis. But it doesn't stop there.

Opticrom eye drops will relieve and prevent red allergic itching eyes at any time of the year, whatever the cause.

And the unique action of sodium cromoglycate has proved extremely effective with none of the troublesome side-effects associated with steroids, anti-histamines, or decongestants.

All through the year Opticrom lets your patients lead normal active lives.

Opticrom[®]
SODIUM CROMOGLYCAT E B.P.



Fisons Limited, Pharmaceutical Division, Derby Road, Loughborough, Leicestershire LE11 0BB. Opticrom is for the treatment of acute allergic conjunctivitis such as hay fever and chronic allergic conjunctivitis. Vernal kerato-conjunctivitis. **Dosage and Administration:** Opticrom is a clear colourless aqueous solution of Sodium Cromoglycate B.P. 2% w/v with benzalkonium chloride 0.01% w/v. 2-Phenylethanol 0.4% w/v. The normal dose is one or two drops into each eye four times daily. Transient stinging may occur in some patients. **Contraindications:** Known hypersensitivity to benzalkonium chloride. **Overdosage:** No action other than medical observation should be necessary. **Basic NHS cost: £4.50 PL: C113/0039.** Registered Trade Mark.

Developments in Immunopathology

*The Proceedings of a Symposium organized by
The Royal College of Pathologists*

Edited by E. J. Holborow

Immunodeficiency—Maturation of the immune system and immunodeficiency • Pathogenesis of hypogammaglobulinaemia with thymoma and late-onset hypogammaglobulinaemia • Combined immunodeficiency and thymic abnormalities • Mechanisms in secondary hypogammaglobulinaemia • Diagnosis of defects of antibody production • Infection of immunosuppressed patients • **Lymphocytes. 1 Function**—Genetic restrictions in the immune response • Lymphocyte function and virus infections • **Lymphocytes. 2 Differentiation**—Differentiation of lymphoid precursor cells • Functional subsets of lymphocytes • **Lymphocytes. 3 Distribution**—Distribution of lymphocytes in health • Distribution of malignant lymphoid cells • **Roy Cameron Lecture**—Control of antibody formation: certain uncertainties • **Antibody-mediated tissue damage**—Hyperviscosity and other complications of paraproteinaemia • Tissue-specific autoantibodies in haemolytic anaemia • Tissue-specific antibodies in myasthenia gravis • Antinuclear acid antibodies • Lymphocytotoxic antibodies in systemic lupus erythematosus: their clinical significance • Complement activation profiles in disease • Antibody affinity: immunogenetic aspects and relationship to immune complex disease • **Therapeutic aspects**—Immunosuppression in established humoral responses

Price: Inland £7.50; Abroad US\$18.75, including postage

Payment must be enclosed with order or a surcharge of 50p will be made for rendering invoices and statements

The Publisher, *Journal of Clinical Pathology*, B.M.A. House, Tavistock Square, London WC1H 9JR

THE BRITISH JOURNAL OF FAMILY PLANNING

The Official Journal of the
National Association of Family Planning Doctors

CO-EDITORS: Dr. Ruth Skrine and Dr. Mary Pollock

The British Journal of Family Planning is a quarterly journal which publishes original papers and review articles by leading authorities on a wide range of relevant subjects. Clinical meetings are reported, and papers of particular interest are often published in full. Each issue contains notices of training courses and future meetings, reviews of important publications and a lively correspondence section. There is often a column contributed by the Institute of Psycho-Sexual Medicine.

The journal serves as a channel of communication between the council of the National Association of Family Planning Doctors and the members. These include doctors working in hospitals, general practice and N.H.S. and private clinics. Consultants from many of the medical specialities are among the membership which now totals nearly 2,000 and is growing rapidly. Health authorities and postgraduate medical centres are subscribing to the journal in increasing numbers.

The journal is published in January, April, July and October.

**Annual Subscription: Inland £12.00
Overseas US \$30.00**

Order from Professional & Scientific Publications,
Tavistock House East, Tavistock Square, London WC1H 9JR
or through any leading subscription agent.

Examination Candidates

Revise with Success
use **PASTEST PRACTICE EXAMS**

**MRCP I & II MRCGP FRCS
PLAB (UK entry exam) MRCOG**

Write for details: Dept J, PasTest Service,
P.O. Box 81,
Hemel Hempstead,
Herts, U.K.
Tel: (0442) 52113

PASTEST

ADVERTISEMENTS

PHARMACEUTICAL PRODUCTS

Readers are reminded that prescribers' literature and clinical information are always available from advertisers on request.



The nasal-block buster

FOR COLDS, RHINITIS AND SINUSITIS

ACTIFED TABLETS

triprolidine hydrochloride BP/pseudoephedrine hydrochloride BP

Indications Symptomatic relief of upper respiratory congestion in: the common cold, hay fever, vasomotor and allergic rhinitis, acute sinusitis, otitis barotrauma.

Dosage Three times a day: Adults and children over 12 years: 1 tablet or 10 ml. 6-12 years: 7.5 ml. 1-6 years: 5 ml. 3-12 months: 2.5 ml.

Contra-indications Actifed* is contraindicated in persons under treatment with MAO inhibitors and within two weeks of stopping such treatment.

Precautions Although at recommended dosage pseudoephedrine has virtually no pressor effects in normotensive subjects, Actifed should be used with caution in patients with cardiovascular disorders. As with other antihistamine-containing

preparations, drowsiness may occur. In some patients the action of antihistamines may be potentiated by alcohol.

Presentation Each Actifed tablet contains 2.5 mg triprolidine hydrochloride and 60 mg pseudoephedrine hydrochloride. PL3/5003.

Each 5 ml of Actifed Syrup contains 1.25 mg triprolidine hydrochloride and 30 mg pseudoephedrine hydrochloride. PL3/5004.

Additional information is available on request.

Wellcome Medical Division
The Wellcome Foundation Ltd., Crewe, Cheshire.



*Trade Mark

An invitation to view London's newest private hospital.

The Directors, Management and Staff of The Cromwell Hospital are proud to inform you that the Hospital opened on 29th April 1981.

We are pleased to extend a cordial invitation to Consultants and Specialists, Nursing and Supplementary Professions to visit the Hospital on any of the following days between the hours of 4.00p.m. and 7.00p.m.

Tuesday 16th June	Tuesday 23rd June
Wednesday 17th June	Wednesday 24th June
Thursday 18th June	Thursday 25th June

Our senior members of staff will be happy to show you round the Hospital and its facilities and to discuss any matters of interest to you.

Bed reservations can be made by telephoning the Medical Reservations Department on 01-370 4233 extension 133 (24 hour service) or directly on 01-370 2297.

Out-patient and Diagnostic Services appointments can be made by

telephoning the Out-Patient Desk on 01-370 4233 extensions 557/8.

Arrangements for further information or visits at other times can be made by contacting either Miss Geraldine McHugh, Director of Nursing or Frank Atkinson, Chief Executive Officer.



THE CROMWELL HOSPITAL. OPENED APRIL 1981

CROMWELL ROAD LONDON SW5 0TU TELEPHONE 01-370 4233 A MEDICAL SERVICES INTERNATIONAL HOSPITAL

Just published

PROCEDURES IN PRACTICE

An illustrated step-by-step guide for doctors making their first attempts at the procedures commonly used in hospital practice. The articles explain the techniques in detail and give indications and contraindications for a range of tasks from setting up a drip to aspirating bone marrow.

The articles have been collected into book form in the same format as they appeared in the BMJ.


Price: Inland £4.00;
Overseas US\$19.00*
(Inland £3.50;
Overseas US\$17.50* for
BMA members)

**including air mail postage*

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller

PROCEDURES IN PRACTICE

ARTICLES FROM THE BRITISH MEDICAL JOURNAL



THE FIRST YEAR OF LIFE

Recent research has vastly improved our understanding of a child's development in the first year of life.

In particular, assessment of normal progress is now much more systematic, so that deviations from the normal are recognised earlier

Dr. Bernard Valman's illustrated account of current methods of examination, assessment and management of the medical problems of early infancy will prove invaluable to all concerned with small babies.

The articles have now been collected into book form in exactly the same format as they appeared in the BMJ.

Price: Inland £5.00;

Overseas US\$12.50


(Inland £4.50;

Overseas US\$11.25 for BMA members)

Order your copy now

From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR

or any leading bookseller




Imodium

TRADEMARK

(loperamide hydrochloride)

symbol of
accuracy
in acute diarrhoea



Imodium's specific action controls diarrhoea, relieving pain and abdominal cramps rapidly and effectively.

A tailored dosage regime means that patients only take Imodium when they really need it, minimising the risk of constipation and encouraging economy of medication.

Patients will find Imodium capsules easy to take and the blister pack of just twelve capsules is convenient to carry throughout the working day.

Imodium

Simply Stops Diarrhoea - fast

**2 // stat and 1 /
at each loose stool**

Further information is available from :



JANSSEN PHARMACEUTICA
2340 Beerse, Belgium

or

JANSSEN PHARMACEUTICAL LTD.
Marlow, Bucks. SL7 1ET.

Presentation :

Hard gelatin capsules containing 2 mg loperamide hydrochloride and syrup containing 1 mg loperamide hydrochloride per 5 ml.

Indications :

Imodium* is indicated for the symptomatic control of acute diarrhoea of any aetiology.

Contra-indications and warnings etc. :

There are no specific contraindications to Imodium*. Studies in animals have shown no abnormal teratogenic effects, however the use of loperamide during pregnancy is subject to the usual precautions. In trials, no side effects have been reported that can reliably be distinguished from the symptoms of the gastro-intestinal disorder being treated. As persistent diarrhoea can be an indicator of potentially more serious conditions, Imodium* should not be used for prolonged periods until the underlying cause of the diarrhoea has been investigated.

Dosage and Administration :

Acute Diarrhoea : Adults : Two capsules initially, followed by one capsule after every loose stool. The usual dosage is 3 to 4 capsules a day; the maximum daily dose should not exceed 8 capsules. **Children :** 4 to 8 years : Syrup 5 ml four times daily until diarrhoea is controlled. In children under 8 years, further investigation may be necessary if diarrhoea has not responded to three days' treatment. 9 to 12 years : Syrup 10 ml (or 1 capsule) four times daily until diarrhoea is controlled.

Basic NHS Cost :

o 12 capsules (ex 250 pack) 104p. (correct at time of printing)


Product Licence Numbers :

Capsules 0242/0028 - Syrup 0242/0040

© JPL/037/80

* Trademark

Geigy



anturan[®]
200

Now calendar packed

**reduces
cardiac mortality
following recent
myocardial infarction**