

Imodium

TRADEMARK

(loperamide hydrochloride)

symbol of
accuracy
in acute diarrhoea



Imodium's specific action
controls diarrhoea, relieving pain and
abdominal cramps rapidly and effectively.

A tailored dosage regime means that patients only take
Imodium when they really need it, minimising the risk of
constipation and encouraging economy of medication.
Patients will find Imodium capsules easy to take and the blister
pack of just twelve capsules is convenient to carry throughout
the working day.

Imodium
Simply Stops Diarrhoea-fast
2 stat and 1
at each loose stool

Further information is available from :



JANSSEN PHARMACEUTICA
2340 Beerse, Belgium

or

JANSSEN PHARMACEUTICAL LTD.
Marlow, Bucks. SL7 1ET.

Presentation :

Hard gelatin capsules containing 2 mg loperamide hydrochloride and syrup containing 1 mg loperamide hydrochloride per 5 ml.

Indications :

Imodium* is indicated for the symptomatic control of acute diarrhoea of any aetiology.

Contra-contraindications and warnings etc. :

There are no specific contraindications to Imodium*. Studies in animals have shown no abnormal teratogenic effects, however the use of loperamide during pregnancy is subject to the usual precautions. In trials, no side effects have been reported that can reliably be distinguished from the symptoms of the gastro-intestinal disorder being treated. As persistent diarrhoea can be an indicator of potentially more serious conditions, Imodium* should not be used for prolonged periods until the underlying cause of the diarrhoea has been investigated.

Dosage and Administration :

Acute Diarrhoea . **Adults** : Two capsules initially, followed by one capsule after every loose stool. The usual dosage is 3 to 4 capsules a day, the maximum daily dose should not exceed 8 capsules. **Children** : 4 to 8 years : Syrup 5 ml four times daily until diarrhoea is controlled. In children under 8 years, further investigation may be necessary if diarrhoea has not responded to three days' treatment. 9 to 12 years : Syrup 10 ml (or 1 capsule) four times daily until diarrhoea is controlled.

Basic NHS Cost :

12 capsules (ex 250 pack) 104p (correct at time of printing)

Product Licence Numbers :

Capsules 0242/0028 · Syrup 0242/0040

Just published

ABC OF EAR NOSE AND THROAT

Diseases of the ear, nose and throat account for a substantial proportion of all consultations in general practice; they cause time off work and, particularly in children, a great deal of misery and discomfort.

Mr. Harold Ludman's illustrated articles on the common problems of ENT provide the busy clinician with a practical and reliable guide to their assessment and management.

These articles have been collected together in book form in the same format as they appeared in the BMJ.

Price: Inland £3.50; Overseas US\$16.00*
(Inland £3.00; Overseas US\$14.50* for
BMA members)

*including air mail postage


Order your copy now

From: The Publisher, British Medical Journal, BMA House,
Tavistock Square, London WC1H 9JR
or any leading bookseller

ABC OF EAR NOSE AND THROAT

ARTICLES FROM THE BRITISH MEDICAL JOURNAL

H LUDMAN



NEW BOOKS AND EDITIONS

ESSENTIALS OF ORAL BIOLOGY

David Adams

152 pages 15 halftone + 53 line illus
paperback £3.95
A brief guide to the biological basis of dental surgery

GYNECOLOGIC ONCOLOGY

Edited by *Malcolm Coppleson*

2 Vols 1140 pages 259 line + 552 halftone +
51 line and tone + 62 colour illus hardback
£75.00

A comprehensive, authoritative and particularly detailed treatment of the whole field of gynecologic oncology.

AIDS TO PATHOLOGY

Michael F. Dixon

Second Edition 280 pages paperback £4.95
Concise revision aid to general and systemic pathology

REHABILITATION AFTER SEVERE HEAD INJURY

C. D. Evans

228 pages 33 line + 33 halftone illus
paperback £5.95
Essential reading for all those involved in the rehabilitation of brain-damaged patients

COMMON FOOT DISORDERS: DIAGNOSIS AND MANAGEMENT

A clinical guide

Edited by *Donald Neale*

264 pages 36 line + 52 halftone illus
paperback £10.95
Surveys the main areas of foot problems and indicates the modern approach to their management


FUNDAMENTALS OF RESPIRATORY MEDICINE

B. J. Sproule P. Lynne-Davies E. Garner King

224 pages approx. 20 line illus 19 tables
paperback £11.00

Provides the necessary data for an adequate, working knowledge of the fundamentals of respiratory disease

**Churchill
Livingstone**



Robert Stevenson House, 1-3 Baxter's Place, Leith Walk,
Edinburgh EH1 3AF, U.K.

Today's antibacterial



Success 9 times out of ten in chest infections would be hard to beat. It is this order of success, confirmed in worldwide clinical studies involving over 8,500 patients, that has helped thousands of doctors to decide that Seprin is today's antibacterial. With over 2,500 published reports and more than 250 million prescriptions written worldwide, is it any wonder that Seprin is Britain's No. 1 branded antibacterial?

PRESCRIBING INFORMATION

Indications Sensitive bacterial infections of the lower respiratory, urinary and genital tracts, sinusitis, otitis media, skin infections, septicaemia, typhoid and paratyphoid fevers, and other infections caused by sensitive organisms.

Dosage Seprin Forte Tablets. Adults and children over 12 years: 1 forte tablet twice daily. Maximum dosage for particularly severe infections 1½ forte tablets twice daily. In acute infections Seprin should be given for a minimum of 5 days or until the patient has been symptom-free for 2 days.

Contra-indications Seprin is contra-indicated in patients with marked liver parenchymal damage, blood dyscrasias or severe renal insufficiency. Seprin should not be given to patients hypersensitive to sulphonamides or co-trimoxazole; should not be given during pregnancy or to neonates.

Precautions In cases of renal impairment a reduced dosage is indicated and an adequate urinary output should be maintained. Regular blood counts are necessary whenever long term therapy is used. Caution is advised in patients with folate deficiency. Care should be taken when giving Seprin to patients receiving oral anticoagulants of the coumarin group, pyrimethamine or sulphonylureas.

Adverse Reactions Occasionally, nausea, vomiting, glossitis and skin rashes may occur with normal doses and, very rarely, haematological reactions.

Presentation Seprin Forte Tablets each contain 160 mg Trimethoprim B.P. and 800 mg Sulphamethoxazole B.P.

PL3/0121



RESEARCH IS OUR ONLY SHAREHOLDER

Further information is available on request.

Wellcome Medical Division

The Wellcome Foundation Ltd, Crewe, Cheshire

Seprin*
CO-TRIMOXAZOLE
Forte
a successful first choice
in chest infections

*Trade Mark.

Just published

TODAY'S TREATMENT/4

The drugs that we use today are increasingly potent, dangerous, and expensive, and every doctor should have some understanding of clinical pharmacology and drug-induced diseases. Both these subjects, which have been badly taught in medical schools, are covered comprehensively in this new book, which consists of articles taken from the *BMJ*. Also included are articles that provide a clear and up-to-the-minute introduction to anaesthetics.

**Price: Inland £4.50;
Overseas US\$20.50*
(Inland £4.00;
Overseas US\$19.00* for
BMA members)**

**including air mail postage*

When ordering BMA members must quote their membership number or the full price will be applicable.

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller



Slowly, surely...



Lopressor® SR
metoprolol tartrate
The 24 hour beta-blocker

Presentation Tablets containing 200mg metoprolol tartrate in a sustained release formulation. *Indications* Hypertension and angina pectoris. *Dosage* Initially one tablet daily. Most patients respond satisfactorily within 14 days to one or two tablets once daily. *Side-effects* Slight gastro-intestinal discomfort and disturbance of sleep pattern. *Precautions* Avoid in patients with cardiac decompensation unless adequately controlled by digitalis and/or diuretics. Care in patients with chronic obstructive pulmonary disease, where dosage of bronchodilator may require adjustment. It may be necessary to adjust the dosage of the hypoglycaemic agent in labile and insulin dependent diabetics. Lopressor SR therapy should be brought to the attention of the anaesthetist prior to general anaesthesia. Occasionally patients taking beta-blocking drugs develop skin rash and/or dry eyes; where such a reaction is otherwise inexplicable, beta-blocker therapy should be gradually withdrawn. Withdrawal of therapy with Lopressor SR should be effected gradually, over 8-10 days. Use in pregnancy only when there are compelling reasons; if used during pregnancy or lactation special attention should be paid to foetus, neonate or infant for undesirable effects of the drug's beta-blocking action. *Contra-indications* Atrioventricular block of second or third degree, heart failure refractory to digitalis, severe bradycardia, cardiogenic shock. *Packs* Lopressor SR tablets in calendar packs of 28 tablets (2 x 14 tablet foils) PL0001/0081 basic NHS price £7.20. ® denotes registered trademark. Full prescribing information is available on request from Geigy Pharmaceuticals, Horsham, West Sussex.

Geigy