

The first Oral broad spectrum anti-fungal.



NEW

Nizoral^{trademark}
(ketoconazole)

Superficial Fungal Infections.

NIZORAL* is a new Oral anti-fungal agent effective against both dermatophyte and yeast infection of the skin and scalp.

Recurrent Vaginal Candidosis.

NIZORAL* offers a simple, effective Oral therapy to aid patient compliance.

Systemic Fungal Infections.

NIZORAL* is a uniquely convenient and well tolerated treatment. It is also effective in prophylaxis for immuno-compromised patients.

PRESCRIBING INFORMATION: Presentation White, flat, half scored uncoated tablets marked 'JANSEN' on the reverse. Each contains ketoconazole 200mg. **Uses** Nizoral* is an orally active antimycotic for the treatment, in adults, of superficial and systemic mycoses including dermatophyte and yeast infection of the skin and hair, yeast infection of the mouth and G.I. tract, recurrent or chronic vaginal candidosis not responding to topical treatment. Also maintenance treatment in systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in 'at risk' patients. In children: systemic mycoses and severe local infection where previous topical treatment has failed. **Side-effects, Precautions, Contra-indications** Contra-indicated in pregnancy. For optimum absorption, Nizoral* should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral*. Nausea, skin rash, occasionally observed. **Dosage** Adults: superficial and systemic mycoses, prophylactic treatment - 200mg once daily increased to 400mg daily if required; vaginal candidosis - 200mg morning and evening for 5 days. Children: 50 or 100mg depending on bodyweight (approximately 3mg/kg). **Product Licence No.** PL 0242/0083. **Basic NHS Cost** 10 tablets £4.90. 30 tablets £14.26 (correct at time of printing).

Further information is available from  Janssen Pharmaceutical Limited, Janssen House, Chapel Street, Marlow, Bucks. SL7 1ET.

*Trademark.

MARITAL PATHOLOGY

Divorce is becoming increasingly common: at current rates about six million people in Britain will divorce before the end of the century. Every doctor and health worker will have to deal with the diverse effects of marital unhappiness. This book discusses clearly and concisely reasons for marital misery, its effects on the health of the family, and how counsellors can help. This book will benefit all who have to deal with these problems.

Price: Inland £1.75
Abroad US\$5.00

Payment must be enclosed with order or a surcharge of 50p will be made for rendering invoices and statements.

Order your copy now

From: The Publishers
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR

or

Darton Longman & Todd Ltd.
89 Lillie Road
London SW6 1UD

MARITAL PATHOLOGY

J DOMINIAN



From the British Medical Journal



Actifed.
Taken around
the world.

Presentation. Actifed is available as tablets and syrup. Each tablet contains 2.5 mg triprolidine hydrochloride and 60 mg pseudoephedrine hydrochloride. Each 5 ml of syrup contains 1.25 mg triprolidine hydrochloride and 30 mg pseudoephedrine hydrochloride.

Indications Symptomatic relief of upper respiratory congestion in the common cold, hay fever, vasomotor and allergic rhinitis, acute sinusitis, otitis barotrauma.

Dosage Three times a day: Adults and children over 12 years: 1 tablet or 10 ml. 6-12 years: 7.5 ml. 1-6 years: 5 ml. 3-12 months: 2.5 ml. **Contra-indications** Actifed is contra-indicated in persons hypersensitive to

pseudoephedrine or triprolidine and in those under treatment with MAO inhibitors and within 2 weeks of stopping such treatment. **Precautions** Although at recommended dosage pseudoephedrine has virtually no pressor effects in normotensive subjects, Actifed should be used with caution in patients with cardiovascular disorders. As with other antihistamine containing preparations, drowsiness may occur. In some patients the action of antihistamines may be potentiated by alcohol. 25 tablets. PL3/5003. 150 ml. PL3/5004.

ActifedTM
Pseudoephedrine Hydrochloride BP
Triprolidine Hydrochloride BP

The decongestant
chosen by NASA


THE FIRST YEAR OF LIFE

Recent research has vastly improved our understanding of a child's development in the first year of life. In particular, assessment of normal progress is now much more systematic, so that deviations from the normal are recognised earlier

Dr. Bernard Valman's illustrated account of current methods of examination, assessment and management of the medical problems of early infancy will prove invaluable to all concerned with small babies.

The articles have now been collected into book form in exactly the same format as they appeared in the BMJ.
Price: Inland £5.00;
Overseas US\$12.50
(Inland £4.50;
Overseas US\$11.25 for BMA members)

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller



TODAY'S TREATMENT/4

The drugs that we use today are increasingly potent, dangerous, and expensive, and every doctor should have some understanding of clinical pharmacology and drug-induced diseases. Both these subjects, which have been badly taught in medical schools, are covered comprehensively in this new book, which consists of articles taken from the *BMJ*. Also included are articles that provide a clear and up-to-the-minute introduction to anaesthetics.

**Price: Inland £4.50;
Overseas US\$20.50*
(Inland £4.00;
Overseas US\$19.00* for
BMA members)**

**including air mail postage*

When ordering BMA members must quote their membership number or the full price will be applicable.

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller



Available NOW
the first edition, in English of the

**Journal of
TRADITIONAL
CHINESE
MEDICINE**

September 1981 Volume 1, Number 1
English Edition

A selection of papers published in this issue follows:

CLINICAL AND EXPERIMENTAL STUDIES ON "SHENGJI (TISSUE-GROWING)" OINTMENT IN INFECTED OPEN FRACTURES • TREATMENT OF GRAND MAL SEIZURES WITH "QUINGYANGSHEN" (ROOT OF *Cynanchum otophyllum*) AND OBSERVATIONS ON EXPERIMENTAL ANIMALS • VOLVULUS OF THE STOMACH SUCCESSFULLY TREATED WITH ACUPUNCTURE REPORT OF 9 CASES • OBSERVATION ON TREATMENT OF POST-OPERATIVE ANAL PAIN WITH EMBEDDED NEEDLE • TREATMENT OF TRICHOMONAL ENTERITIS BY CHINESE MEDICINAL HERBS Report of 16 cases • **EFFECT OF TRADITIONAL MASSAGE ON CLOSURE INSUFFICIENCY OF GLOTTIS IN SINGERS** An analysis of 16 cases.

Published quarterly the subscription rates are:


1981 (2 issues) Inland £15.00; Overseas US\$30.00
1981 (2 issues) by Air Mail US\$34.00
1981-1982 (6 issues) Inland £45.00; Overseas US\$90.00
1981-1982 (6 issues) by Air Mail US\$102.00

The Journal of Traditional Chinese Medicine is published in Beijing, China and solely distributed worldwide (except China) by:

**Professional & Scientific Publications
BMA House, Tavistock Square
London WC1H 9JR, England**

ORDER YOUR SUBSCRIPTION NOW
from Professional & Scientific Publications

Take two



Voltarol[®] diclofenac sodium **for osteoarthritis**

Presentation Tablets of 25 and 50mg diclofenac sodium; suppositories of 100mg diclofenac sodium. **Indications** Rheumatoid arthritis, osteoarthritis, low back pain, ankylosing spondylitis. **Dosage** Tablets: 75-150mg daily in two or three divided doses. Suppositories: one daily, usually administered at night. In more severe cases, combined therapy with tablets is recommended (daily dose should not exceed 150mg). **Contra-indications** Peptic ulceration; patients in whom attacks of asthma, urticaria or acute rhinitis are precipitated by aspirin or other NSAIs. **Precautions** Do not prescribe during pregnancy unless there are compelling reasons. Patients with a history of peptic ulcer, haematemesis or melaena, or with severe hepatic or renal insufficiency, should be kept under close surveillance. Voltarol has been reported to depress salicylate levels and vice versa; the clinical relevance of this is not yet clear. Use suppositories only with caution in patients with painful or irritable ano-rectal conditions. **Side-effects** Transient epigastric pain, nausea and diarrhoea, headache and slight dizziness have been reported, as occasionally have skin rash, peripheral oedema and abnormalities of serum transaminases and (very rarely) peptic ulcer and haematemesis or melaena. Local reactions to suppositories include itching, burning and increased frequency of bowel movement. **NHS price** Tablets 25mg: pack of 100 - £9.00; 50mg: pack of 100 - £17.50; Suppositories: pack of 10 - £2.98. **Product licence numbers** Tablets 25mg PL0001/0036, 50mg PL0001/0082, Suppositories 100mg PL0001/0083. Full prescribing information is available from **Geigy** Pharmaceuticals, Horsham, West Sussex.