

The first Oral broad spectrum anti-fungal.



NEW

Nizoral^{trademark}
(ketoconazole)

Superficial Fungal Infections.

NIZORAL* is a new Oral anti-fungal agent effective against both dermatophyte and yeast infection of the skin and scalp.

Recurrent Vaginal Candidosis.

NIZORAL* offers a simple, effective Oral therapy to aid patient compliance.

Systemic Fungal Infections.

NIZORAL* is a uniquely convenient and well tolerated treatment. It is also effective in prophylaxis for immuno-compromised patients.

PRESCRIBING INFORMATION: Presentation White, flat, half scored uncoated tablets marked 'JANSSEN' on the reverse. Each contains ketoconazole 200mg. **Uses** Nizoral* is an orally active antimycotic for the treatment, in adults, of superficial and systemic mycoses including dermatophyte and yeast infection of the skin and hair, yeast infection of the mouth and G.I. tract, recurrent or chronic vaginal candidosis not responding to topical treatment. Also maintenance treatment in systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in 'at risk' patients. In children: systemic mycoses and severe local infection where previous topical treatment has failed. **Side-effects, Precautions, Contra-indications** Contra-indicated in pregnancy. For optimum absorption, Nizoral* should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral*. Nausea, skin rash, occasionally observed. **Dosage** Adults: superficial and systemic mycoses, prophylactic treatment - 200mg once daily increased to 400mg daily if required: vaginal candidosis - 200mg morning and evening for 5 days. Children: 50 or 100mg depending on bodyweight (approximately 3mg/kg). **Product Licence No.** PL 0242/0083. **Basic NHS Cost** 10 tablets £4.90. 30 tablets £14.26 (correct at time of printing).

Further information is available from  Janssen Pharmaceutical Limited, Janssen House, Chapel Street, Marlow, Bucks. SL7 1ET.

*Trademark

THE FIRST YEAR OF LIFE

Recent research has vastly improved our understanding of a child's development in the first year of life.

In particular, assessment of normal progress is now much more systematic, so that deviations from the normal are recognised earlier

Dr. Bernard Valman's illustrated account of current methods of examination, assessment and management of the medical problems of early infancy will prove invaluable to all concerned with small babies.

The articles have now been collected into book form in exactly the same format as they appeared in the BMJ.

Price: Inland £5.00;

Overseas US\$12.50


(Inland £4.50;

Overseas US\$11.25 for BMA members)

Order your copy now

From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR

or any leading bookseller





Actifed.
Taken around
the world.

Presentation. Actifed is available as tablets and syrup. Each tablet contains 2.5mg triprolidine hydrochloride and 60mg pseudoephedrine hydrochloride. Each 5ml of syrup contains 1.25mg triprolidine hydrochloride and 30mg pseudoephedrine hydrochloride.

Indications Symptomatic relief of upper respiratory congestion in the common cold, hay fever, vasomotor and allergic rhinitis, acute sinusitis, otitis barotrauma.

Dosage Three times a day: Adults and children over 12 years: 1 tablet or 10ml. 6-12 years: 7.5ml. 1-6 years: 5ml. 3-12 months: 2.5ml. **Contra-indications** Actifed is contra-indicated in persons hypersensitive to

pseudoephedrine or triprolidine and in those under treatment with MAO inhibitors and within 2 weeks of stopping such treatment. **Precautions** Although at recommended dosage pseudoephedrine has virtually no pressor effects in normotensive subjects, Actifed should be used with caution in patients with cardiovascular disorders. As with other antihistamine containing preparations, drowsiness may occur. In some patients the action of antihistamines may be potentiated by alcohol. 25 tablets. PL3/5003. 150ml. PL3/5004.

ActifedTM
Pseudoephedrine Hydrochloride BP
Triprolidine Hydrochloride BP

The decongestant
chosen by NASA



Further information is available on request.

ABC OF EAR NOSE AND THROAT

Diseases of the ear, nose and throat account for a substantial proportion of all consultations in general practice; they cause time off work and, particularly in children, a great deal of misery and discomfort.

Mr. Harold Ludman's illustrated articles on the common problems of ENT provide the busy clinician with a practical and reliable guide to their assessment and management.

These articles have been collected together in book form in the same format as they appeared in the BMJ.


Price: Inland £3.50;
Overseas US\$16.00*
(Inland £3.00;
Overseas US\$14.50* for
BMA members)

**including air mail postage*

Order your copy now

From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR

or any leading bookseller



EPIDEMIOLOGY FOR THE UNINITIATED

Do you know the difference between the incidence of a disease and its prevalence? How to set up a valid controlled trial? How to plan and conduct a survey? Many doctors would like to carry out some simple clinical research but find they lack basic information of this kind. The answers were published in 1978-79 in a series of BMJ articles, now collected together in book form—essential reading for anyone contemplating starting a research study.


Price: Inland £2.50;
Abroad US\$6.25

(Concessionary price to BMA members: Inland £2.00;
Abroad US\$5.00.

When ordering BMA members must quote their membership number or the full price will be applicable.)

Payment must be enclosed with order or a surcharge of 30p will be made for rendering invoices and statements.

Order your copy now
From: The Publisher
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR
or any leading bookseller



PROCEDURES IN PRACTICE

An illustrated step-by-step guide for doctors making their first attempts at the procedures commonly used in hospital practice. The articles explain the techniques in detail and give indications and contraindications for a range of tasks from setting up a drip to aspirating bone marrow.

The articles have been collected into book form in the same format as they appeared in the BMJ.


Price: Inland £4.00;
Overseas US\$19.00*
(Inland £3.50;
Overseas US\$17.50* for
BMA members)

**including air mail postage*

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller

PROCEDURES IN PRACTICE

ARTICLES FROM THE BRITISH MEDICAL JOURNAL



Freedom of movement



Voltarol[®]

diclofenac sodium

for arthritis

Presentation Tablets of 25 and 50mg diclofenac sodium; suppositories of 100mg diclofenac sodium. **Indications** Rheumatoid arthritis, osteoarthritis, low back pain, ankylosing spondylitis. **Dosage** Tablets: 75-150mg daily in two or three divided doses. Suppositories: one daily, usually administered at night. In more severe cases, combined therapy with tablets is recommended (daily dose should not exceed 150mg). **Contra-indications** Peptic ulceration; patients in whom attacks of asthma, urticaria or acute rhinitis are precipitated by aspirin or other NSAIs. **Precautions** Do not prescribe during pregnancy unless there are compelling reasons. Patients with a history of peptic ulcer, haematemesis or melaena, or with severe hepatic or renal insufficiency, should be kept under close surveillance. Voltarol has been reported to depress salicylate levels and vice versa; the clinical relevance of this is not yet clear. Use suppositories only with caution in patients with painful or irritable ano-rectal conditions. **Side-effects** Transient epigastric pain, nausea and diarrhoea, headache and slight dizziness have been reported, as occasionally have skin rash, peripheral oedema and abnormalities of serum transaminases and (very rarely) peptic ulcer and haematemesis or melaena. Local reactions to suppositories include itching, burning and increased frequency of bowel movement. **NHS price** Tablets 25mg: pack of 100 - £9.00; 50mg: pack of 100 - £17.50; Suppositories: pack of 10 - £2.98. **Product licence numbers** Tablets 25mg PL0001/0036, 50mg PL0001/0082, Suppositories 100mg PL0001/0083. Full prescribing information is available from Geigy Pharmaceuticals, Horsham, West Sussex.

Geigy