


Just published

STATISTICS IN PRACTICE

No doctor can afford to ignore statistics: most modern medical research uses statistics. This important and authoritative book, which is a collection of articles that have appeared in the BMJ, provides clear information on designing studies, applying statistical techniques, and interpreting studies that use statistics. It can be easily understood by those with no statistical training and should be read by all those who want to keep abreast of new developments.

Price: Inland £7.00
Overseas US\$27.50*
(Inland £6.00;
Overseas US\$25.00* for
BMA members)
**including air mail postage*

Order your copy now
From: The Publisher
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR
or any leading bookseller



EPIDEMIOLOGY FOR THE UNINITIATED

Do you know the difference between the incidence of a disease and its prevalence? How to set up a valid controlled trial? How to plan and conduct a survey? Many doctors would like to carry out some simple clinical research but find they lack basic information of this kind. The answers were published in 1978-79 in a series of BMJ articles, now collected together in book form—essential reading for anyone contemplating starting a research study.

Price: Inland £2.50;
Abroad US\$6.25

(Concessionary price to BMA members: Inland £2.00;
Abroad US\$5.00.


When ordering BMA members must quote their membership number or the full price will be applicable.)


Payment must be enclosed with order or a surcharge of 30p will be made for rendering invoices and statements.

Order your copy now
From: The Publisher
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR
or any leading bookseller

EPIDEMIOLOGY FOR THE UNINITIATED

GEOFFREY HINE AND J. HENRI





Septtrin Assurance

Prescribing Information

Indications Sensitive bacterial infections of the lower respiratory, urinary and genital tracts, sinusitis, otitis media, skin infections, septicaemia, typhoid and paratyphoid fevers, and other infections caused by sensitive organisms.

Dosage Septtrin Forte Tablets. Adults and children over 12 years: 1 forte tablet twice daily. Maximum dosage for particularly severe infections 1½ forte tablets twice daily. In acute infections Septtrin should be given for a minimum of five days or until the patient has been symptom-free for two days.

Contra-indications Septtrin is contra-indicated in

patients with marked liver parenchymal damage, blood dyscrasias or severe renal insufficiency.

Septtrin should not be given to patients hypersensitive to sulphonamides, trimethoprim or co-trimoxazole; should not be given during pregnancy or to neonates.

Precautions In renal impairment a reduced dosage is indicated and an adequate urinary output should be maintained. Regular blood counts are necessary whenever long-term therapy is used. Caution is advised in patients with folate deficiency. Care should be taken when giving Septtrin to patients receiving oral anticoagulants of the coumarin group, pyrimethamine or sulphonylureas.

Adverse Reactions Occasionally, nausea, vomiting, glossitis and skin rashes may occur with normal doses and, very rarely, haematological reactions.

Presentation Septtrin Forte Tablets each contain 160 mg Trimethoprim BP and 800 mg Sulphamethoxazole BP. PL3/0121.

Septtrin* Forte 1b.d. co-trimoxazole

Further information is available on request.
Wellcome Medical Division
The Wellcome Foundation Ltd., Crewe, Cheshire



*Trade Mark

THE FIRST YEAR OF LIFE

Recent research has vastly improved our understanding of a child's development in the first year of life.

In particular, assessment of normal progress is now much more systematic, so that deviations from the normal are recognised earlier

Dr. Bernard Valman's illustrated account of current methods of examination, assessment and management of the medical problems of early infancy will prove invaluable to all concerned with small babies.

The articles have now been collected into book form in exactly the same format as they appeared in the BMJ.

Price: Inland £5.00;

Overseas US\$12.50


(Inland £4.50;

Overseas US\$11.25 for BMA members)

Order your copy now

From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR

or any leading bookseller



The first Oral broad spectrum anti-fungal.



NEW

Nizoral
(ketoconazole)

Superficial Fungal Infections.

NIZORAL* is a new Oral anti-fungal agent effective against both dermatophyte and yeast infection of the skin and scalp.

Recurrent Vaginal Candidosis.

NIZORAL* offers a simple, effective Oral therapy to aid patient compliance.

Systemic Fungal Infections.

NIZORAL* is a uniquely convenient and well tolerated treatment. It is also effective in prophylaxis for immuno-compromised patients.


PRESCRIBING INFORMATION: **Presentation** White, flat, half scored uncoated tablets marked 'JANSSEN' on the reverse. Each contains ketoconazole 200 mg. **Uses** Nizoral* is an orally active antimycotic for the treatment, in adults, of superficial and systemic mycoses including dermatophyte and yeast infection of the skin and hair, yeast infection of the mouth and G.I. tract, recurrent or chronic vaginal candidosis not responding to topical treatment. Also maintenance treatment in systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in 'at risk' patients. In children: systemic mycoses and severe local infection where previous topical treatment has failed. **Side-effects, Precautions, Contra-indications** Contra-indicated in pregnancy. For optimum absorption, Nizoral* should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral*. Nausea, skin rash, occasionally observed. **Dosage** Adults: superficial and systemic mycoses, prophylactic treatment - 200 mg once daily increased to 400 mg daily if required; vaginal candidosis - 200 mg morning and evening for 5 days. Children: 50 or 100 mg depending on bodyweight (approximately 3 mg/kg). **Product Licence No.** PL 0242/0083. **Basic NHS Cost** 10 tablets £4.90. 30 tablets £14.26 (correct at time of printing).

Further information is available from  Janssen Pharmaceutical Limited, Janssen House, Chapel Street, Marlow, Bucks. SL7 1ET.

*Trademark

The logical choice

Select the correct figure from the six numbered ones



Voltarol[®] diclofenac sodium for osteoarthritis

Presentation Tablets of 25 and 50mg diclofenac sodium; suppositories of 100mg diclofenac sodium. **Indications** Rheumatoid arthritis, osteoarthritis, low back pain, ankylosing spondylitis. **Dosage** Tablets: 75-150mg daily in two or three divided doses. Suppositories: one daily, usually administered at night. In more severe cases, combined therapy with tablets is recommended (daily dose should not exceed 150mg). **Contra-indications** Peptic ulceration; patients in whom attacks of asthma, urticaria or acute rhinitis are precipitated by aspirin or other NSAIs. **Precautions** Do not prescribe during pregnancy unless there are compelling reasons. Patients with a history of peptic ulcer, haematemesis or melaena, or with severe hepatic or renal insufficiency, should be kept under close surveillance. Voltarol has been reported to depress salicylate levels and vice versa; the clinical relevance of this is not yet clear. Use suppositories only with caution in patients with painful or irritable ano-rectal conditions. **Side-effects** Transient epigastric pain, nausea and diarrhoea, headache and slight dizziness have been reported, as occasionally have skin rash, peripheral oedema and abnormalities of serum transaminases and (very rarely) peptic ulcer and haematemesis or melaena. Local reactions to suppositories include itching, burning and increased frequency of bowel movement. **NHS price** Tablets 25mg: pack of 100 - £9.00; 50mg: pack of 100 - £17.50; Suppositories: pack of 10 - £2.98. **Product licence numbers** Tablets 25mg PL0001/0036, 50mg PL0001/0082, Suppositories 100mg PL0001/0083. Full prescribing information is available from Geigy Pharmaceuticals, Horsham, West Sussex.

Geigy