The first Oral broad spectrum anti-fungal.



Superficial Fungal Infections.

NIZORAL* is a new Oral anti-fungal agent effective against both dermatophyte and yeast infection of the skin and scalp.

Recurrent Vaginal Candidosis.

NIZORAL* offers a simple, effective Oral therapy to aid patient compliance.

Systemic Fungal Infections.

NIZORAL* is a uniquely convenient and well tolerated treatment. It is also effective in prophylaxis for immunocompromised patients.

PRESCRIBING INFORMATION: Presentation White, flat, half scored uncoated tablets marked 'JANSSEN' on the reverse. Each contains ketoconazole 200 mg. Uses Nizoral' is an orally active antimycotic for the treatment, in adults, of superficial and systemic mycases including demotophyte and yeast infection of the six and hair, yeast infection of the south and G.I. tract, recurrent or chronic vagainal conditions not responding to topical freatment. Also maintenance treatment in systemic mycases and severe local infection where previous topical treatment has failed. Side-effects, Precautions, Centra-indications Contra-indicated in pregnancy. For optimum absorption, Nizoral' should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less then two laws of the Nizoral' Nousea, skin rash, occasionally observed. Desege Adults: superficial and systemic mycases, prophylactic treatment—200 mg once daily increased to 400 mg daily fire frequired: vaginal candidates and conditions. The product Leases Me. PL (242/0083). Resize MRSC Case 10 tablets \$24.90. 30 tablets \$24.90. and of printing of printing in distributions and conditions are described in programment.

TODAY'S TREATMENT/4

The drugs that we use today are increasingly potent, dangerous, and expensive, and every doctor should have some understanding of clinical pharmacology and drug-induced diseases. Both these subjects, which have been badly taught in medical schools, are covered comprehensively in this new book, which consists of articles taken from the *BMJ*. Also included are articles that provide a clear and up-to-the-minute introduction to anaesthetics.

Price: Inland £4.50; Overseas US\$20.50* (Inland £4.00; Overseas US\$19.00* for BMA members) *including air mail postage

When ordering BMA members must quote their membership number or the full price will be applicable.

Order your copy now From: The Publisher, British Medical Journal, BMA House, Tavistock Square, London WC1H 9JR or any leading bookseller





pseudoephedrine hýdrochloride. Each 5 ml of syrup contains 1.25 mg triprolidine hydrochloride and 30mg pseudoephedrine hydrochloride Indications Symptomatic relief of upper respiratory congestion in the common cold, hay fever, vasomotor and allergic rhinitis, acute sinusitis, otitis barotrauma. **Dosage** Three times a day: Adults and children over 12 years: 1 tablet or 10ml. 6-12 years: 7.5ml. 1-6 years: 5ml. 3-12 months: 2.5ml. Contra-indications Actifed is contra-indicated in persons hypersensitive to

treatment. Precautions Although at recommended dosage pseudoephedrine has virtually no pressor effects in normotensive subjects. Actifed should be used with caution in patients with cardiovascular disorders. As with other antihistamine containing preparations, drowsiness may occur. In some patients the action of antihistamines may be potentiated by alcohol. 25 tablets. PL3/5003. 150 ml. PL3/5004.

Pseudoephedrine Hydrochloride BP Triprolidine Hydrochloride BP

The decongestant chosen by NASA



Further information is available on request. Wellcome Medical Division. The Wellcome Foundation Ltd., Crewe, Cheshire Just published

STATISTICS IN PRACTICE


No doctor can afford to ignore statistics: most modern medical research uses statistics. This important and authoritative book, which is a collection of articles that have appeared in the BMJ, provides clear information on designing studies, applying statistical techniques, and interpreting studies that use statistics. It can be easily understood by those with no statistical training and should be read by all those who want to keep abreast of

new developments.


Price: Inland £7.00 Overseas US\$27.50* (Inland £6.00: Overseas US\$25.00* for BMA members) *including air mail postage

Order your copy now

From: The Publisher British Medical Journal BMA House Tavistock Square London WC1H 9JR or any leading bookseller



in the work-up for vascular surgery



Praxilene FORTE 200mg infusion relieves ischaemic rest pain, allows time for detailed assessment and preparation for arterial surgery

Naftidrofuryl (Praxilene) can significantly reduce the patient's local symptoms and improve his mental state and wellbeing — it is these local symptoms which often strongly influence the urgency of the decision to amputate.

Angiology, In Press, 1982

Dosage: I ampoule twice daily by intravenous or intra-arterial infusion in 250 to 500ml saline, dextrose or low molecular weight dextran.

Contra-indications: patients with atrioventricular block or those who show hypersensitivity to the drug.

Precautions; exercise care in patients with severe cardiac insufficiency and cardiac disorders. The possibility of an additive effect occurring between Praxilene FORTE and antiarrhythmic and beta-blocking drugs should be considered. Side-effects: nausea may occasionally be noted.

Not for bolus injection.

PL 3759/0004. Packs 10 x 10ml ampoules £10.95.

Further information and Data Sheet available on request.



Lipha Pharmaceuticals Limited
Old Farm Road, West Drayton, Middlesex UB7 7LD
LIP 84

Proxidene naftidrofuryl INN FORTE buys time

KINOWS INO DAIDINGERS

staphylococcal infection

is a phylocologic form

is after decretic

in there dievare

diam the delence

to man of the body

hielded from efficient

findecapy.

Accelia exerts a potent instructerial effect even in the presence of large actestions of pas, into talch Kapparently accessives in effective accessivations?

Arcidia, which has the diffic to genetrate in a distribution of the distribution of th

- Brain abscess
- Hypopyon
- Endophthalmitis
- Lung abscess
- Endocarditis
- Pneumonia with abscess
- o Empyema
- Renal carbuncle
- Deep wound infection
- Septie arthritis
- Foreign bodies post
- Osteomyelitis
- Districtic gange-new

*#**

Children and the graph of the State of the S

administrated the conjumption with other drugs which such confusion for the nume exceeders pullinary. Produces the code Alice (44.275019

Common C B product

