

# Nizoral

TRADEMARK  
ketoconazole

The first  
oral anti-fungal  
effective against  
all pathogenic  
fungi

dosage:

Vaginal Candidosis:	All other superficial and systemic fungal infections:
1 tablet b.i.d. (with food) for 5 days	1 tablet daily (with food) until complete symptomatic and mycological cure is obtained

**PRESCRIBING INFORMATION**

**Presentation:** White, flat, half scored uncoated tablets marked 'JANSSEN' on the reverse. Each contains ketoconazole 200 mg. **Uses:** Nizoral is an orally active antimycotic for the treatment, in adults, of superficial and systemic mycoses including dermatophyte and yeast infection of the skin and hair, yeast infection of the mouth and G.I. tract, recurrent or chronic vaginal candidosis not responding to topical treatment. Also maintenance treatment in systemic mycosis and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycosis and severe local infection where previous topical treatment has failed. **Side-effects, Precautions, Contra-indications:** Contra-indicated in pregnancy. For optimum absorption, Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub> blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, occasionally observed.

Not all indications are as yet approved in all countries.

Janssen Pharmaceutica   
Beerse, Belgium

*Just published*

# STATISTICS IN PRACTICE

No doctor can afford to ignore statistics: most modern medical research uses statistics. This important and authoritative book, which is a collection of articles that have appeared in the BMJ, provides clear information on designing studies, applying statistical techniques, and interpreting studies that use statistics. It can be easily understood by those with no statistical training and should be read by all those who want to keep abreast of new developments.

Price: Inland £7.00

Overseas US\$27.50\*

(Inland £6.00;

Overseas US\$25.00\* for  
BMA members)

*\*including air mail postage*

**Order your copy now**

From: The Publisher


British Medical Journal

BMA House

Tavistock Square

London WC1H 9JR

*or any leading bookseller*





Actifed.  
Taken around  
the world.

**Presentation.** Actifed is available as tablets and syrup. Each tablet contains 2.5mg triprolidine hydrochloride and 60mg pseudoephedrine hydrochloride. Each 5ml of syrup contains 1.25mg triprolidine hydrochloride and 30mg pseudoephedrine hydrochloride.

**Indications** Symptomatic relief of upper respiratory congestion in the common cold, hay fever, vasomotor and allergic rhinitis, acute sinusitis, otitis barotrauma.

**Dosage** Three times a day: Adults and children over 12 years: 1 tablet or 10ml. 6-12 years: 7.5ml. 1-6 years: 5ml. 3-12 months: 2.5ml. **Contra-indications** Actifed is contra-indicated in persons hypersensitive to

pseudoephedrine or triprolidine and in those under treatment with MAO inhibitors and within 2 weeks of stopping such treatment. **Precautions** Although at recommended dosage pseudoephedrine has virtually no pressor effects in normotensive subjects, Actifed should be used with caution in patients with cardiovascular disorders. As with other antihistamine containing preparations, drowsiness may occur. In some patients the action of antihistamines may be potentiated by alcohol. 25 tablets. PL3/5003. 150ml. PL3/5004.


**Actifed**<sup>TM</sup>  
Pseudoephedrine Hydrochloride BP  
Triprolidine Hydrochloride BP

The decongestant  
chosen by NASA



Further information is available on request.  
Wellcome Medical Division, The Wellcome Foundation Ltd., Crewe, Cheshire

# in the work-up for vascular surgery



Praxilene FORTE 200mg infusion relieves ischaemic rest pain, allows time for detailed assessment and preparation for arterial surgery

Naftidrofuryl (Praxilene) can significantly reduce the patient's local symptoms and improve his mental state and wellbeing – it is these local symptoms which often strongly influence the urgency of the decision to amputate.

Angiology, In Press, 1982

*Dosage:* 1 ampoule twice daily by intravenous or intra-arterial infusion in 250 to 500ml saline, dextrose or low molecular weight dextran.

*Contra-indications:* patients with atrioventricular block or those who show hypersensitivity to the drug.

*Precautions:* exercise care in patients with severe cardiac insufficiency and cardiac disorders. The possibility of an additive effect occurring between Praxilene FORTE and antiarrhythmic and beta-blocking drugs should be considered.

*Side-effects:* nausea may occasionally be noted.

**Not for bolus injection.**

PL 3759/0004. Packs 10 x 10ml ampoules £10.95.

Further information and Data Sheet available on request.




Lipha Pharmaceuticals Limited  
Old Farm Road, West Drayton, Middlesex UB7 7LD

LIP 84

**Praxilene<sup>®</sup>**  
naftidrofuryl INN  
**FORTE**  
**buys time**

**Take two**



# **Voltarol<sup>®</sup>**

**diclofenac sodium**

## **for osteoarthritis**

**Presentation** Tablets of 25 and 50mg diclofenac sodium; suppositories of 100mg diclofenac sodium. **Indications** Rheumatoid arthritis, osteoarthritis, low back pain, ankylosing spondylitis. **Dosage** Tablets: 75-150mg daily in two or three divided doses. Suppositories: one daily, usually administered at night. In more severe cases, combined therapy with tablets is recommended (daily dose should not exceed 150mg). **Contra-indications** Peptic ulceration; patients in whom attacks of asthma, urticaria or acute rhinitis are precipitated by aspirin or other NSAIs. **Precautions** Do not prescribe during pregnancy unless there are compelling reasons. Patients with a history of peptic ulcer, haematemesis or melaena, or with severe hepatic or renal insufficiency, should be kept under close surveillance. Voltarol has been reported to depress salicylate levels and vice versa; the clinical relevance of this is not yet clear. Use suppositories only with caution in patients with painful or irritable ano-rectal conditions. **Side-effects** Transient epigastric pain, nausea and diarrhoea, headache and slight dizziness have been reported, as occasionally have skin rash, peripheral oedema and abnormalities of serum transaminases and (very rarely) peptic ulcer and haematemesis or melaena. Local reactions to suppositories include itching, burning and increased frequency of bowel movement. **NHS price** Tablets 25mg: pack of 100 - £9.00; 50mg: pack of 100 - £17.50; Suppositories: pack of 10 - £2.98. **Product licence numbers** Tablets 25mg PL0001/0036, 50mg PL0001/0082, Suppositories 100mg PL0001/0083. Full prescribing information is available from **Geigy** Pharmaceuticals, Horsham, West Sussex.