

NizoralTM

ketoconazole

The first
oral anti-fungal
effective against
all pathogenic
fungi

dosage:

Vaginal Candidosis:	All other superficial and systemic fungal infections:
1 tablet b.i.d. (with food) for 5 days	1 tablet daily (with food) until complete sympto- matic and mycological cure is obtained

PRESCRIBING INFORMATION

Presentation: White, flat, half scored uncoated tablets marked 'JANSSEN' on the reverse. Each contains ketoconazole 200 mg. **Uses:** Nizoral is an orally active antimycotic for the treatment, in adults, of superficial and systemic mycoses including dermatophyte and yeast infection of the skin and hair, yeast infection of the mouth and G.I. tract, recurrent or chronic vaginal candidosis not responding to topical treatment. Also maintenance treatment in systemic mycosis and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children, systemic mycosis and severe local infection where previous topical treatment has failed. **Side-effects, Precautions, Contra-indications:** Contra-indicated in pregnancy. For optimum absorption, Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, occasionally observed.

Not all indications are as yet
approved in all countries.

Janssen Pharmaceutica 
Beerse, Belgium

STATISTICS IN PRACTICE

No doctor can afford to ignore statistics: most modern medical research uses statistics. This important and authoritative book, which is a collection of articles that have appeared in the BMJ, provides clear information on designing studies, applying statistical techniques, and interpreting studies that use statistics. It can be easily understood by those with no statistical training and should be read by all those who want to keep abreast of new developments.

Price: Inland £7.00

Overseas US\$27.50*

(Inland £6.00;

Overseas US\$25.00* for
BMA members)

**including air mail postage*

Order your copy now

From: The Publisher


British Medical Journal

BMA House

Tavistock Square

London WC1H 9JR

or any leading bookseller





Actifed.
Taken around
the world.

Presentation. Actifed is available as tablets and syrup. Each tablet contains 2.5 mg triprolidine hydrochloride and 60 mg pseudoephedrine hydrochloride. Each 5 ml of syrup contains 1.25 mg triprolidine hydrochloride and 30 mg pseudoephedrine hydrochloride.


Indications Symptomatic relief of upper respiratory congestion in the common cold, hay fever, vasomotor and allergic rhinitis, acute sinusitis, otitis barotrauma.

Dosage Three times a day: Adults and children over 12 years: 1 tablet or 10 ml. 6-12 years: 7.5 ml. 1-6 years: 5 ml. 3-12 months: 2.5 ml. **Contra-indications** Actifed is contra-indicated in persons hypersensitive to

pseudoephedrine or triprolidine and in those under treatment with MAO inhibitors and within 2 weeks of stopping such treatment. **Precautions** Although at recommended dosage pseudoephedrine has virtually no pressor effects in normotensive subjects, Actifed should be used with caution in patients with cardiovascular disorders. As with other antihistamine containing preparations, drowsiness may occur. In some patients the action of antihistamines may be potentiated by alcohol. 25 tablets PL3/5003. 150 ml PL3/5004.

ActifedTM
Pseudoephedrine Hydrochloride BP
Triprolidine Hydrochloride BP

The decongestant
chosen by NASA

 Further information is available on request.
Wellcome Wellcome Medical Division, The Wellcome Foundation Ltd., Crewe, Cheshire

ABC OF HYPERTENSION


Hypertension is a common disorder that is easily detected. But it may be deceptively simple: the measurement of blood pressure, interpretation of the results, and the decisions that follow need to be based on a comprehensive knowledge of recent advances.

These illustrated articles on blood pressure measurement, reduction and management have now been collected together in the same format as they appeared in the BMJ to provide a practical guide for general practitioners and others concerned with hypertension.

**Price: Inland £3.50;
Overseas US\$16.00*
(Inland £3.00;
Overseas US\$14.50* for
BMA members)**

**including air mail postage*

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller



PROCEDURES IN PRACTICE

An illustrated step-by-step guide for doctors making their first attempts at the procedures commonly used in hospital practice. The articles explain the techniques in detail and give indications and contraindications for a range of tasks from setting up a drip to aspirating bone marrow.

The articles have been collected into book form in the same format as they appeared in the BMJ.

Price: Inland £4.00;
Overseas US\$19.00*
(Inland £3.50;
Overseas US\$17.50* for
BMA members)

**including air mail postage*


Order your copy now

From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR

or any leading bookseller

PROCEDURES IN PRACTICE

ARTICLES FROM THE BRITISH MEDICAL JOURNAL



Available NOW
the second issue in English of the

**Journal of
TRADITIONAL
CHINESE
MEDICINE**

December 1981 Volume 1, Number 2

A selection of papers published in this and the first (September 1981) issue follows:

RECENT ADVANCES IN DIAGNOSIS AND TREATMENT OF ACUTE APPENDICITIS COMPLICATED WITH PERITONITIS BY COMBINED TRADITIONAL CHINESE AND WESTERN MEDICINE • CLINICAL STUDIES OF PSORIASIS TREATED WITH ANTIPSORIATIC PRESCRIPTIONS • SPLEEN POINT IN FACE ACUPUNCTURE USED IN FIBROGASTROSCOPY • A REPORT ON THE NON-MUTAGENICITY OF QINGHAOSU BY THE AMES' TEST • TREATMENT OF GRAND MAL SEIZURES WITH "QUINGYANGSHEN" (ROOT OF CYNACHUM OTOPHYLLUM) AND OBSERVATIONS ON EXPERIMENTAL ANIMALS • VOLVULUS OF THE STOMACH SUCCESSFULLY TREATED WITH ACUPUNCTURE Report of 9 cases • TREATMENT OF TRICHOMONAL ENTERITIS BY CHINESE MEDICINAL HERBS Report of 16 cases • EFFECT OF TRADITIONAL MASSAGE ON CLOSURE INSUFFICIENCY OF GLOTTIS IN SINGERS An analysis of 16 cases.

Published quarterly the subscription rates are:

1981 (2 issues) Inland £15.00; Overseas US\$30.00
1981 (2 issues) by Air Mail US\$34.00
1981-1982 (6 issues) Inland £45.00; Overseas US\$90.00
1981-1982 (6 issues) by Air Mail US\$102.00

The Journal of Traditional Chinese Medicine is published in Beijing, China and solely distributed worldwide (except China) by:

**Professional & Scientific Publications
BMA House, Tavistock Square
London WC1H 9JR, England**

**ORDER YOUR SUBSCRIPTION NOW
from Professional & Scientific Publications**

sodium fusidate B.P

"... Fucidin, which has the ability to penetrate in significant amounts into tissues carrying a poor blood supply..."

- **Brain abscess**
- **Hypopyon**
- **Endophthalmitis**
- **Lung abscess**
- **Endocarditis**
- **Pneumonia with abscess**
- **Empyema**
- **Renal carbuncle**
- **Deep wound infection**
- **Septic arthritis**
- **Foreign bodies** } **grafts**
 prostheses
- **Osteomyelitis**
- **Diabetic gangrene**

[illegible]

1. Johnson, R., and Larsen, H.C.A. *Ann. Intern. Med.*, 66, 790, 1964.
2. Crisp, H. M. *Br. Med. J.*, 1, 705, 1965.
3. Rowley, D. G. *J. Bone Joint Surg.*, 52B, 302, 1970.

Ben Laboratories Limited
 100-101, Park Road, Princess Risborough
 Bucks. HP27 9NR