Available NOW the second issue in English of the

Journal of TRADITIONAL CHINESE MEDICINE

December 1981 Volume 1, Number 2

A selection of papers published in this and the first (September 1981) issue follows:

RECENT ADVANCES IN DIAGNOSIS AND TREATMENT OF ACUTE APPENDICITIS
COMPLICATED WITH PERITONITIS BY COMBINED TRADITIONAL CHINESE AND
WESTERN MEDICINE • CLINICAL STUDIES OF PSORIASIS TREATED WITH
ANTIPSORIATIC PRESCRIPTIONS • SPLEEN POINT IN FACE ACUPUNCTURE USED
IN FIBROGASTROSCOPY • A REPORT ON THE NON-MUTAGENICITY OF QINGHAOSU
BY THE AMES' TEST • TREATMENT OF GRAND MAL SEIZURES WITH "QUINGYANGSHEN"
(ROOT OF CYNACHUM OTOPHYLLUM) AND OBSERVATIONS ON EXPERIMENTAL
ANIMALS • VOLVULUS OF THE STOMACH SUCCESSFULLY TREATED WITH
ACUPUNCTURE Report of 9 cases • TREATMENT OF TRICHOMONAL ENTERITIS
BY CHINESE MEDICINAL HERBS Report of 16 cases • EFFECT OF TRADITIONAL
MASSAGE ON CLOSURE INSUFFICIENCY OF GLOTTIS IN SINGERS An analysis of 16 cases.

Published quarterly the subscription rates are:

1981 (2 issues) Inland £15.00; Overseas US\$30.00 1981 (2 issues) by Air Mail US\$34.00 1981-1982 (6 issues) Inland £45.00; Overseas US\$90.00 1981-1982 (6 issues) by Air Mail US\$102.00

The Journal of Traditional Chinese Medicine is published in Beijing, China and solely distributed worldwide (except China) by:

Professional & Scientific Publications BMA House, Tavistock Square London WC1H 9JR, England

ORDER YOUR SUBSCRIPTION NOW from Professional & Scientific Publications

PROCEDURES IN PRACTICE


An illustrated step-by-step guide for doctors making their first attempts at the procedures commonly used in hospital practice. The articles explain the techniques in detail and give indications and contraindications for a range of tasks from setting up a drip to aspirating bone marrow.

The articles have been collected into book form in the same format as they appeared in the BMJ.

Price: Inland £4.00; Overseas US\$19.00* (Inland £3.50; Overseas US\$17.50* for BMA members)

*including air mail postage

Order your copy now From: The Publisher. British Medical Journal. BMA House, Tavistock Square, London WC1H 9JR or any leading bookseller





and 30mg pseudoephedrine hydrochloride.

Indications Symptomatic relief of upper respiratory congestion in the common cold, hay fever, vasomotor and allergic rhinitis, acute sinusitis, otitis barotrauma. **Dosage** Three times a day: Adults and children over 12 years: 1 tablet or 10ml. 6-12 years: 7.5ml. 1-6 years: 5ml. 3-12 months: 2.5ml. Contra-indications Actifed is contra-indicated in persons hypersensitive to

no pressor effects in normotensive subjects, Actifed should be used with caution in patients with cardiovascular disorders. As with other antihistamine containing preparations, drowsiness may occur. In some patients the action of antihistamines may be potentiated by alcohol. 25 tablets. PL3/5003. 150ml. PL3/5004

Triprolidine Hydrochloride BP

The decongestant chosen by NASA



ABC OF EAR NOSE AND THROAT

Diseases of the ear, nose and throat account for a substantial proportion of all consultations in general practice; they cause time off work and, particularly in children, a great deal of misery and discomfort.


Mr. Harold Ludman's illustrated articles on the common problems of ENT provide the busy clinician with a practical and reliable guide to their assessment and management.

These articles have been collected together in book form in the same format as they appeared in the BMJ.

Price: Inland £3.50; Overseas US\$16.00* (Inland £3.00; Overseas US\$14.50* for BMA members)

*including air mail postage

Order your copy now From: The Publisher, British Medical Journal, BMA House, Tavistock Square, London WC1H 9JR or any leading bookseller



STATISTICS by T. D. V. SWINSCOW LSOLJARE ONE

from the British Medical Journal

The statistical testing of data is indispensable in many types of medical investigation and a help on countless occasions in clinical practice. This book provides step-by-step instruction. Subjects covered include standard deviation, χ^2 tests, t tests, non-parametric tests and correlation. Methods specially adapted to pocket calculators.


"The ability to put symbols in sequence may well be one of the greatest advances in man's development.... This excellent and attractive volume will give you an idea into which category an article or projected article fits. The ink you save may be your own. It is superb value.'

O'Donnell, Barry, British Medical Journal, 1977, 1, 451.

". . . this gem of a practical booklet squeezes the essence of biostatistics into 75 well-written, wellorganized pages. Newcomers to the field will like the way obscure concepts are clarified; experts will find the book valuable as a teaching aid."

ADVERTISEMENT

de Jong, Rudolph H., J.A.M.A., 1977, 237, 1874



Seventh edition

PRICE: Inland £2.50 Overseas US\$10.00, including despatch by Air Mail

Payment must be enclosed with order or a surcharge of 50p will be made for rendering invoices and statements

ORDER YOUR COPY NOW

From: The Publisher. British Medical Journal, BMA House, Tavistock Square, London WC1H 9JR

or through any leading bookseller

ketoconazole





Janssen Pharmaceutica 🍒



Freedom of movement



Voltarol[®] for arthritis

Presentation Tablets of 25 and 50mg diclofenac sodium; suppositories of 100mg diclofenac sodium. Indications Rheumatoid arthritis, osteoarthritis, low back pain, ankylosing spondylitis Dosage Tablets: 75-150mg daily in two or three divided doses. Suppositories: one daily, usually administered at night. In more severe cases, combined therapy with tablets is recommended (daily dose should not exceed 150mg). Contra-indications Peptic ulceration; patients in whom attacks of asthma, urticaria or acute rhinitis are precipitated by asprin or other NSAIs. Precautions Do not prescribe during pregnancy unless there are compelling reasons. Patients with a history of peptic ulcer, haematemesis or melaena, or with severe hepatic or renal insufficiency, should be kept under close surveillance. Voltarol has been reported to depress salicylate levels and vice versa; the clinical relevance of this is not yet clear. Use suppositories only with caution in patients with painful or irritable ano-rectal conditions. Side-effects Transient epigastric pain, nausea and diarrhoea, headache and slight dizziness have been reported, as occasionally have skin rash, peripheral oederna and abnormalities of serum transaminases and (very, rarely) peptic ulcer and haematemesis or melaena. Local reactions to suppositories include itching, burning and increased frequency of bowel movement. NHS price Tablets 25mg: pack of 100 – £9.00; 50mg: pack of 100 – £17.50; Suppositories: pack of 10 – £2.98. Product licence numbers Tablets 25mg PL0001/0036, 50mg PL0001/0082, Suppositories 100mg PL0001/0083. Full prescribing information is available from Geigy Pharmaceuticals, Horsham, West Sussex.