

Nizoral

ketconazole

The first
oral anti-fungal
effective against
all pathogenic
fungi

dosage:

Vaginal Candidosis:	All other superficial and systemic fungal infections:
1 tablet b.i.d. (with food) for 5 days	1 tablet daily (with food) until complete symptomatic and mycological cure is obtained

PRESCRIBING INFORMATION

Presentation: White, flat, half scored uncoated tablets marked 'JANSSEN' on the reverse. Each contains ketconazole 200 mg. **Uses:** Nizoral is an orally active antimycotic for the treatment, in adults, of superficial and systemic mycoses including dermatophyte and yeast infection of the skin and hair, yeast infection of the mouth and G.I. tract, recurrent or chronic vaginal candidosis not responding to topical treatment. Also maintenance treatment in systemic mycosis and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycosis and severe local infection where previous topical treatment has failed. **Side-effects, Precautions, Contra-indications:** Contra-indicated in pregnancy. For optimum absorption, Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, occasionally observed.

Not all indications are as yet approved in all countries.

Janssen Pharmaceutica
Beerse, Belgium



TODAY'S TREATMENT/4

The drugs that we use today are increasingly potent, dangerous, and expensive, and every doctor should have some understanding of clinical pharmacology and drug-induced diseases. Both these subjects, which have been badly taught in medical schools, are covered comprehensively in this new book, which consists of articles taken from the *BMJ*. Also included are articles that provide a clear and up-to-the-minute introduction to anaesthetics.

**Price: Inland £4.50;
Overseas US\$20.50*
(Inland £4.00;
Overseas US\$19.00* for
BMA members)**

**including air mail postage*

When ordering BMA members must quote their membership number or the full price will be applicable.

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller





Actifed.
Taken around
the world.

Presentation. Actifed is available as tablets and syrup. Each tablet contains 2.5 mg triprolidine hydrochloride and 60 mg pseudoephedrine hydrochloride. Each 5 ml of syrup contains 1.25 mg triprolidine hydrochloride and 30 mg pseudoephedrine hydrochloride.

Indications Symptomatic relief of upper respiratory congestion in the common cold, hay fever, vasomotor and allergic rhinitis, acute sinusitis, otitis barotrauma.

Dosage Three times a day: Adults and children over 12 years: 1 tablet or 10 ml. 6-12 years: 7.5 ml. 1-6 years: 5 ml. 3-12 months: 2.5 ml. **Contra-indications** Actifed is contra-indicated in persons hypersensitive to

pseudoephedrine or triprolidine and in those under treatment with MAO inhibitors and within 2 weeks of stopping such treatment. **Precautions** Although at recommended dosage pseudoephedrine has virtually no pressor effects in normotensive subjects, Actifed should be used with caution in patients with cardiovascular disorders. As with other antihistamine containing preparations, drowsiness may occur. In some patients the action of antihistamines may be potentiated by alcohol. 25 tablets. PL3/5003. 150 ml. PL3/5004.

ActifedTM
Pseudoephedrine Hydrochloride BP
Triprolidine Hydrochloride BP

The decongestant
chosen by NASA

HOW TO DO IT

Every doctor knows the sinking sensation induced by waiting one's turn for an examination viva and the social anxieties of changing jobs. The misery of first night nerves does not, however, disappear with time: each decade brings new challenges. On these occasions—chairing a committee, giving evidence in court, appearing on television—practical tips can be enormously helpful. The same is true of the many skills not taught in medical school—how to lecture, referee papers, edit, or use a library. All these and many more nuggets of down-to-earth advice have been collected from past issues of the *BMJ* to form a compendium for the novice—whatever his or her age or seniority.

Price: Inland £4.00;
Abroad US\$10.00

(Concessionary price to BMA
members: Inland £3.50;
Abroad US\$8.75.

When ordering BMA members must
quote their membership number or
the full price will be applicable.)

Payment must be enclosed with order or a surcharge
of 30p will be made for rendering invoices and
statements.

Order your copy now
From: The Publisher
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR
or any leading bookseller

HOW TO DO IT

Chair a committee. Be a dictator. Take an examination. Be a student. Give evidence. Organise a meeting. Chair a conference. Use a library. Improve a student. Chair a medical journal. Raise funds. Be an examiner. Give a reference. Survive as an editor. Use slides. Appear on television. Use an auditor. Apply for a research grant. Use an overhead projector. Plan a research project. Write a paper.

PROCEDURES IN PRACTICE

An illustrated step-by-step guide for doctors making their first attempts at the procedures commonly used in hospital practice. The articles explain the techniques in detail and give indications and contraindications for a range of tasks from setting up a drip to aspirating bone marrow.

The articles have been collected into book form in the same format as they appeared in the BMJ.


Price: Inland £4.00;
Overseas US\$19.00*
(Inland £3.50;
Overseas US\$17.50* for
BMA members)

**including air mail postage*

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller

PROCEDURES IN PRACTICE

ARTICLES FROM THE BRITISH MEDICAL JOURNAL



Available NOW
the second issue in English of the

**Journal of
TRADITIONAL
CHINESE
MEDICINE**

December 1981 Volume 1, Number 2

A selection of papers published in this and the first (September 1981) issue follows:

RECENT ADVANCES IN DIAGNOSIS AND TREATMENT OF ACUTE APPENDICITIS COMPLICATED WITH PERITONITIS BY COMBINED TRADITIONAL CHINESE AND WESTERN MEDICINE • CLINICAL STUDIES OF PSORIASIS TREATED WITH ANTIPSORIATIC PRESCRIPTIONS • SPLEEN POINT IN FACE ACUPUNCTURE USED IN FIBROGASTROSCOPY • A REPORT ON THE NON-MUTAGENICITY OF QINGHAOSU BY THE AMES' TEST • TREATMENT OF GRAND MAL SEIZURES WITH "QUINGYANGSHEN" (ROOT OF CYNACHUM OTOPHYLLUM) AND OBSERVATIONS ON EXPERIMENTAL ANIMALS • VOLVULUS OF THE STOMACH SUCCESSFULLY TREATED WITH ACUPUNCTURE Report of 9 cases • TREATMENT OF TRICHOMONAL ENTERITIS BY CHINESE MEDICINAL HERBS Report of 16 cases • EFFECT OF TRADITIONAL MASSAGE ON CLOSURE INSUFFICIENCY OF GLOTTIS IN SINGERS An analysis of 16 cases.

Published quarterly the subscription rates are:

1981 (2 issues) Inland £15.00; Overseas US\$30.00
1981 (2 issues) by Air Mail US\$34.00
1981-1982 (6 issues) Inland £45.00; Overseas US\$90.00
1981-1982 (6 issues) by Air Mail US\$102.00

The Journal of Traditional Chinese Medicine is published in Beijing, China and solely distributed worldwide (except China) by:

**Professional & Scientific Publications
BMA House, Tavistock Square
London WC1H 9JR, England**

**ORDER YOUR SUBSCRIPTION NOW
from Professional & Scientific Publications**



Behind the
gentleness of

Burinex^{*}K

bumetanide and slow release potassium chloride

lies the power of

Burinex

Burinex K
gently effective
for maintenance

Burinex tablets
combine strength with
gentleness for more refractory oedema

Burinex injection
fast powerful action for emergencies

Formulations Burinex Injection: 0.5 mg/ml in 2 ml, 4 ml and 10 ml ampoules. Burinex Tablets: 1 mg and 5 mg. Burinex K: 0.5 mg bumetanide, 7.7 mmol slow release potassium chloride. **Indications** Acute pulmonary oedema and oedema of cardiac, renal or hepatic origins. **Dosages** Burinex Injection: Initially 1-2 mg i.v., if necessary repeated at 20 minute intervals to achieve desired response. Where appropriate higher doses may be given by infusion over 30-60 minutes. Burinex Tablets: Most patients require 1 mg Burinex daily as morning or evening dose. In refractory cases dosage can be increased to achieve the desired response. For high dose treatment 5 mg Burinex should be given initially and increased by 5 mg steps at 12-24 hour intervals until desired response is achieved. Burinex K: Most patients require 2 tablets Burinex K daily. **Contra-indications, Precautions and Side Effects** Contra-indicated in hepatic coma, severe electrolyte depletion and severe progressive renal failure. Hypokalaemia and circulatory collapse may follow inappropriately excessive diuresis. Concurrent digitalis therapy in association with electrolyte disturbances may lead to digitalis toxicity. Concurrent antihypertensive or antidiabetic therapy may require adjustment. Caution should be exercised in the first trimester of pregnancy. Burinex K is contra-indicated in combination with potassium sparing agents. Burinex K should be stopped immediately if signs or symptoms of bowel ulceration appear. Side effects such as skin rashes, muscular cramps, rises in serum uric acid and thrombocytopenia may rarely occur. **Product Licence Numbers:** Burinex Injection: 0043/0060 Burinex Tablets: 0043/0021, 0043/0043 Burinex K: 0043/0027B **Basic N.H.S. Prices** Burinex Injection: 0.5 mg/ml - 5 x 4 ml £3.34 Burinex Tablets: 1 mg - 100 tabs £4.74 Burinex K: 100 tabs £3.24



^{*}Burinex is a trade mark

Leo Laboratories Limited, Longwick Road, Princes Risborough, Aylesbury, Bucks. HP17 9RR