

Nizoral

ketoconazole

antifungal
dermatological
preparation

dosage:

vaginal
candidosis:

2 tablets
once daily
(with food)
for 5 days

all other superficial and
systemic fungal infections:

1 tablet daily
(with food)
until complete sympto-
matic and mycological
cure is obtained

For a complete list of contraindications, see package leaflet.

Uses, side-effects, precautions, contra-indications, see package leaflet.

Side-effects, precautions, contra-indications, see package leaflet.

For a complete list of contraindications, see package leaflet.

For a complete list of contraindications, see package leaflet.

Janssen Pharmaceutica
B-2340 Beerse, Belgium



BRITISH MEDICAL JOURNAL
EUROPEAN LIFE SCIENCES EDITORS ASSOCIATION
EDITING A MEDICAL JOURNAL:
a workshop for editors of medical journals

to be held at the
Royal Station Hotel, Station Road, York YO2 2AA on 4/5 November 1982

Programme

Thursday 4 November 1982

1500	Registration
1515	Introduction, Dr Stephen Lock (Editor, <i>BMJ</i>)
1520	What others are up to: reports from the annual meetings of the Council of Biology Editors and ELSE
1600	Tea
1630	The business aspects of publishing a journal, Mr J Cameron (Editorial Director, John Wiley)
1900 for 1930	Dinner

Friday 5 November 1982

0900-1030	Workshops*
1030	Coffee
1100-1230	Workshops*
1300	Lunch
1400	Problems and pitfalls of being a part-time editor: introduced and chaired by Dr Simon Barley (Editor, <i>Journal of the Royal College of General Practitioners</i>)
1530	Tea

*The workshops will be on: subediting (led by Dr W F Whimster and Ms Maeve O'Connor) and layout (led by Mr Leslie McCrombie)

The fee of £70 (£65 for members of ELSE or the BMA) includes a single room and all meals. The day rate for those wanting to attend only on Friday is £28 (£25 for ELSE or BMA members). The number of places is limited, and applications, including the fee (cheques made payable to the British Medical Journal), should be sent as soon as possible to the Editor, *British Medical Journal*, BMA House, Tavistock Square, London WC1H 9JR. Please would editors enclose a copy of their journal when applying, for possible use in the layout workshop.

Statistics for editors of medical journals

5/6 November 1982

A meeting on statistics for editors of medical journals, aimed at helping them to assess the statistical aspects of papers they receive for publication, will be held at the Royal Station Hotel, Station Road, York YO2 2AA from 1630 on 5 November to 1600 on 6 November.

The meeting will take the form of a series of seminars, conducted by Mr Douglas Altman (Medical Statistician, MRC Clinical Research Centre), which will provide an elementary grounding in statistics for non-statistician editors to help them assess papers and to make the best use of their statistical referees. Numbers will be limited.

The fee of £70 (£65 for members of the BMA or ELSE) includes a single room and all meals. A special rate of £130 (£125 for members of BMA or ELSE) is available for editors attending both the statistics conference and the editing conference on 4/5 November. Applications, including fee (cheques made payable to the British Medical Journal), should be sent as soon as possible to the Editor, *British Medical Journal*, BMA House, Tavistock Square, London WC1H 9JR.



Actifed.
Taken around
the world.

Presentation. Actifed is available as tablets and syrup. Each tablet contains 2.5 mg triprolidine hydrochloride and 60 mg pseudoephedrine hydrochloride. Each 5 ml of syrup contains 1.25 mg triprolidine hydrochloride and 30 mg pseudoephedrine hydrochloride.

Indications Symptomatic relief of upper respiratory congestion in the common cold, hay fever, vasomotor and allergic rhinitis, acute sinusitis, otitis barotrauma.

Dosage Three times a day: Adults and children over 12 years: 1 tablet or 10 ml. 6-12 years: 7.5 ml. 1-6 years: 5 ml. 3-12 months: 2.5 ml. **Contra-indications** Actifed is contra-indicated in persons hypersensitive to

pseudoephedrine or triprolidine and in those under treatment with MAO inhibitors and within 2 weeks of stopping such treatment. **Precautions** Although at recommended dosage pseudoephedrine has virtually no pressor effects in normotensive subjects, Actifed should be used with caution in patients with cardiovascular disorders. As with other antihistamine containing preparations, drowsiness may occur. In some patients the action of antihistamines may be potentiated by alcohol. 25 tablets. PL3/5003. 150 ml. PL3/5004.

ActifedTM
Pseudoephedrine Hydrochloride BP
Triprolidine Hydrochloride BP

The decongestant
chosen by NASA



Further information is available on request.
Wellcome Medical Division, The Wellcome Foundation Ltd., Crewe, Cheshire

Available NOW
the second issue in English of the

**Journal of
TRADITIONAL
CHINESE
MEDICINE**

December 1981 Volume 1, Number 2

A selection of papers published in this and the first (September 1981) issue follows:

RECENT ADVANCES IN DIAGNOSIS AND TREATMENT OF ACUTE APPENDICITIS COMPLICATED WITH PERITONITIS BY COMBINED TRADITIONAL CHINESE AND WESTERN MEDICINE • CLINICAL STUDIES OF PSORIASIS TREATED WITH ANTIPSORIATIC PRESCRIPTIONS • SPLEEN POINT IN FACE ACUPUNCTURE USED IN FIBROGASTROSCOPY • A REPORT ON THE NON-MUTAGENICITY OF QINGHAOSU BY THE AMES' TEST • TREATMENT OF GRAND MAL SEIZURES WITH "QUINGYANGSHEN" (ROOT OF CYNACHUM OTOPHYLLUM) AND OBSERVATIONS ON EXPERIMENTAL ANIMALS • VOLVULUS OF THE STOMACH SUCCESSFULLY TREATED WITH ACUPUNCTURE Report of 9 cases • TREATMENT OF TRICHOMONAL ENTERITIS BY CHINESE MEDICINAL HERBS Report of 16 cases • EFFECT OF TRADITIONAL MASSAGE ON CLOSURE INSUFFICIENCY OF GLOTTIS IN SINGERS An analysis of 16 cases.

Published quarterly the subscription rates are:

1981 (2 issues) Inland £15.00; Overseas US\$30.00

1981 (2 issues) by Air Mail US\$34.00

1981-1982 (6 issues) Inland £45.00; Overseas US\$90.00

1981-1982 (6 issues) by Air Mail US\$102.00

The Journal of Traditional Chinese Medicine is published in Beijing, China and solely distributed worldwide (except China) by:

**Professional & Scientific Publications
BMA House, Tavistock Square
London WC1H 9JR, England**

**ORDER YOUR SUBSCRIPTION NOW
from Professional & Scientific Publications**

In hypertension

TENORMIN

Atenolol 100mg

The only beta-blocker to put it all together in one.

Full 24 hour control

One tablet daily

Wide patient
spectrum

Few CNS
side-effects

Hydrophilic

Possible
advantages
in smokers

Cardioselective

Cardioprotective

Tenormin fits the profile of the ideal beta-blocker for hypertension.

TENORMIN

A unique combination of hydrophilicity
and cardioselectivity

Prescribing Notes:

Dosage: One tablet daily. **Contraindications:** Heart block. Co-administration with verapamil. **Precautions:** Untreated cardiac failure, bradycardia, renal failure, anaesthesia and pregnancy. **Side Effects:** Coldness of extremities and muscular fatigue. Sleep disturbance rarely seen. Rashes and dry eyes have been reported with beta blockers—consider discontinuance if they occur. Cessation of therapy with beta blockers should be gradual. **Pack size and Basic NHS cost:** 'Tenormin' 28's £7.27.
Product Licence Number: 'Tenormin' 0029/0122.

Full prescribing information is available on request to the company



Stuart Pharmaceuticals Limited
Carr House Carrs Road
Cheadle Cheshire SK8 2EG
Tenormin is a trade mark for atenolol.

