


# Pharmacia Diagnostics

## a total approach to atopic allergy

**Prediction:** Early detection of high total IgE identifies the infant at risk enabling early prophylaxis.

**Diagnosis:** Total IgE measured by Phadebas IgE PRIST® and Phadezym IgE PRIST® showing the likelihood of atopic allergy.  
Specific IgE measured by Phadebas RAST® and Phadezym RAST® determines the causative allergens.



**Treatment:** Allergen immunotherapy with standardized, purified and stabilized preparations, providing optimal dosage and predictable response.

**Monitoring:** Measurement of blocking antibodies to insect venoms by Phadebas IgG RAST® shows the response to immunotherapy.

## Pharmacia Diagnostics leads the way to optimal management of atopic allergy

Our considerable expertise is at your service; please refer your allergy related problems to your nearest divisional office.

**Australia:** Pharmacia (South Seas) Pty. Ltd., NORTH RYDE, N.S.W., Tel: 02-888 36 22  
**Austria:** Pharmacia Ges. m.b.H., WIEN, Tel: 222-67 05 07  
**Belgium:** N.V. Pharmacia Beiga S.A., BRUSSELS, Tel: 02-7 36 99 90  
**Canada:** Pharmacia (Canada) Inc., DORVAL, Québec, Tel: 514-684-8881  
**Denmark:** Pharmacia AS, HILLERÖD, Tel: 02-26 52 00  
**Finland:** OY STAR AB, Tel: 931-17 66 11  
**France:** Pharmacia France S.A., BOIS D'ARCY, Tel: 03-043 99 19  
**German Federal Republic:** Deutsche Pharmacia G.m.b.H., FREIBURG, Tel: 0761-41011  
**Great Britain:** Pharmacia (Great Britain) Ltd, HOUNSLOW, Tel: 01-572 7321

**Japan:** Pharmacia (Japan) K.K., TOKYO 150, Tel: 03-406-4995  
**The Netherlands:** Pharmacia Nederland B.V., WOERDEN, Tel: 3480-16464  
**Norway:** Norsk Pharmacia A/S, OSLO, Tel: 02-209735  
**Sweden:** Pharmacia Norden AB, UPPSALA, Tel: 018-15 60 40  
**Switzerland:** Pharmacia-pf(Schweiz) A.G., ZUERICH, Tel: 01-55 60 77  
**USA:** Pharmacia Diagnostics Division of Pharmacia Inc., PISCATAWAY N.J., Tel: 201-457-8000

**Other Countries:** Pharmacia Diagnostics AB, International, UPPSALA, Tel: 018-16 30 00

**Pharmacia Diagnostics AB**  
Uppsala, Sweden

 **Pharmacia  
Diagnostics**

Available NOW  
in English the  
**Journal of  
TRADITIONAL  
CHINESE  
MEDICINE**

June 1982 Volume 2, Number 2

*A selection of papers published in this and the March 1982 issues:*

**THE TRADITIONAL CHINESE METHOD OF CLOSED REDUCTION OF FRACTURES AND DISLOCATIONS WITH A ROLLING ROD** Further Development in Its Use and the Mechanical Principles Involved • **MANIPULATIVE TREATMENT OF 12 CASES OF CERVICAL SPONDYLOSIS WITH TRIGEMINAL NEURALGIA** • **111 CASES OF SNAKE-BITE (AGKISTRODON ACUTIS GUENTHER) TREATED BY COMBINED TRADITIONAL CHINESE AND WESTERN MEDICINE** **KIDNEY REINFORCEMENT REGIMEN IN THE TREATMENT OF BRONCHIAL ASTHMA** Endocrinological and Immunological Studies • **MOXIBUSTION OF POINT TANZHONG AND MASSAGE OF POINT TIANZONG IN 47 CASES ACUTE MASTITIS** • **ANTIMALARIAL EFFICACY AND MODE OF ACTION OF QINGHAOSU AND ITS DERIVATIVES IN EXPERIMENTAL MODELS** • **METABOLISM AND PHARMAKINETICS OF QINGHAOSU AND ITS DERIVATIVES** • **CLINICAL STUDIES ON THE TREATMENT OF MALARIA WITH QINGHAOSU AND ITS DERIVATIVES** • **THERAPEUTIC EFFECT AND MECHANISM OF ATHERAPEUTIC EFFECT AND MECHANISM OF ACUPUNCTURE AT NEIGUAN (P. 6) IN CHRONIC RHEUMATIC HEART DISEASE**

*Published quarterly the subscription rates are:*

1982 (4 issues) Inland £30.00; Overseas US\$60.00


1982 (4 issues) by Air Mail US\$68.00

1981 Volume 1, numbers 1 (first edition) and 2 are still available at  
Inland £12.00; Overseas US\$24.00 (by air mail US\$27.00)

**The Journal of Traditional Chinese Medicine is published in Beijing, China and solely distributed worldwide (except China) by:**

Professional & Scientific Publications  
BMA House, Tavistock Square  
London WC1H 9JR, England

**ORDER YOUR SUBSCRIPTION NOW  
from Professional & Scientific Publications**



## Septtrin Assurance

### Prescribing Information

**Indications** Sensitive bacterial infections of the lower respiratory, urinary and genital tracts, sinusitis, otitis media, skin infections, septicaemia, typhoid and paratyphoid fevers, and other infections caused by sensitive organisms.

**Dosage** Septtrin Forte Tablets. Adults and children over 12 years: 1 forte tablet twice daily. Maximum dosage for particularly severe infections 1½ forte tablets twice daily. In acute infections Septtrin should be given for a minimum of five days or until the patient has been symptom-free for two days.

**Contra-indications** Septtrin is contra-indicated in

patients with marked liver parenchymal damage, blood dyscrasias or severe renal insufficiency.

Septtrin should not be given to patients hypersensitive to sulphonamides, trimethoprim or co-trimoxazole; should not be given during pregnancy or to neonates.

**Precautions** In renal impairment a reduced dosage is indicated and an adequate urinary output should be maintained. Regular blood counts are necessary whenever long-term therapy is used. Caution is advised in patients with folate deficiency. Care should be taken when giving Septtrin to patients receiving oral anticoagulants of the coumarin group, pyrimethamine or sulphonylureas.

**Adverse Reactions** Occasionally, nausea, vomiting, glossitis and skin rashes may occur with normal doses and, very rarely, haematological reactions.

**Presentation** Septtrin Forte Tablets each contain 160mg Trimethoprim BP and 800mg Sulphamethoxazole BP. PL3/0121.

## Septtrin\* Forte 1b.d. co-trimoxazole

Further information is available on request.  
Wellcome Medical Division  
The Wellcome Foundation Ltd., Crewe, Cheshire



\*Trade Mark

# PROCEDURES IN PRACTICE

An illustrated step-by-step guide for doctors making their first attempts at the procedures commonly used in hospital practice. The articles explain the techniques in detail and give indications and contraindications for a range of tasks from setting up a drip to aspirating bone marrow.

The articles have been collected into book form in the same format as they appeared in the BMJ.

Price: Inland £4.00;  
Overseas US\$19.00\*  
(Inland £3.50;  
Overseas US\$17.50\* for  
BMA members)

*\*including air mail postage*


**Order your copy now**

From: The Publisher,  
British Medical Journal,  
BMA House,  
Tavistock Square,  
London WC1H 9JR

*or any leading bookseller*

## PROCEDURES IN PRACTICE

ARTICLES FROM THE BRITISH MEDICAL JOURNAL



# THE FIRST YEAR OF LIFE

Recent research has vastly improved our understanding of a child's development in the first year of life.


In particular, assessment of normal progress is now much more systematic, so that deviations from the normal are recognised earlier

Dr. Bernard Valman's illustrated account of current methods of examination, assessment and management of the medical problems of early infancy will prove invaluable to all concerned with small babies.

The articles have now been collected into book form in exactly the same format as they appeared in the BMJ.

Price: Inland £5.00;  
Overseas US\$12.50  
(Inland £4.50;  
Overseas US\$11.25 for  
BMA members)

**Order your copy now**  
From: The Publisher,  
British Medical Journal,  
BMA House,  
Tavistock Square,  
London WC1H 9JR  
*or any leading bookseller*



# Nizoral

TRADENAME

ketoconazole

The first  
oral anti-fungal  
effective against  
all pathogenic  
fungi

dosage:

vaginal candidosis:	all other superficial and systemic fungal infections:
2 tablets once daily (with food) for 5 days	1 tablet daily (with food) until complete sympto- matic and mycological cure is obtained

**PRESCRIBING INFORMATION**

**Presentation:** white, flat, half scored uncoated tablets marked "Janssen" on one side and 1/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and GI tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children, systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub>-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped.

Not all indications are as yet  
approved in all countries.

Janssen Pharmaceutica  
B-2340 Beerse, Belgium





# fucidin<sup>\*</sup>

sodium fusidate B.P.


## knows no barriers!

For the treatment of staphylococcal infection

“... (staphylococci) form focal, often necrotic lesions, where they are protected from the defence mechanism of the body and shielded from efficient chemotherapy.”

“... Fucidin exerts a potent antibacterial effect even in the presence of large collections of pus, into which it apparently penetrates in effective concentrations.”

“... Fucidin, which has the ability to penetrate in significant amounts into tissues carrying a poor blood supply.”



- Brain abscess
- Hypopyon
- Endophthalmitis
- Lung abscess
- Endocarditis
- Pneumonia with abscess
- Empyema
- Renal carbuncle
- Deep wound infection
- Septic arthritis
- Foreign bodies, <sup>grafts</sup> prostheses
- Osteomyelitis
- Diabetic gangrene

**Dosage:**  
Adults: 500mg (2 capsules) 4 times daily with food.  
Children: 20-40mg/kg body weight daily in 4 equal doses with food.  
The standard dosage may be doubled for severe staphylococcal infections.  
**Adverse reactions and precautions:**  
Occasionally mild diarrhoea occurs in patients receiving high doses of Fucidin. In some cases this may be relieved by giving the capsules with food. Fucidin should be discontinued if severe diarrhoea occurs. Fucidin should be discontinued if severe allergic reactions occur. Fucidin should be discontinued if severe skin reactions occur. Fucidin should be discontinued if severe liver reactions occur. Fucidin should be discontinued if severe kidney reactions occur. Fucidin should be discontinued if severe blood reactions occur. Fucidin should be discontinued if severe other reactions occur.

antibacterial in combination with other drugs which may interfere with the same sensitivity patterns.  
Fucidin is a registered trademark of LEO Laboratories.  
Each capsule contains 500mg sodium fusidate B.P.

**References:**  
1. J. Antimicrob. Chemother. 1970, 1, 1-10.  
2. J. Antimicrob. Chemother. 1970, 1, 1-10.  
3. J. Antimicrob. Chemother. 1970, 1, 1-10.  
**LEO Laboratories Limited**  
Loughborough, Leics.  
L12 2BQ, England  
\*Fucidin is a trademark