## ketoconazole

# therapy infec

Vaginal candidosis:	In all dermatological and in systemic fungal infections:
2 tablets once daily (with food) for 5 days	1 tablet daily (with food) until complete sympto- matic and mycological cure is obtained

Not all indications are as yet approved in all countries.

PRESCRIBING INFORMATION
Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and k/200 on the reverse Each tablet contains 200 mg ketoconazole. Uses: Nizoral is an orally active antimycotic for the treatment in adults, of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed. Side-effects, precautions, contra-indicated in pregnancy. For maximal absorption. Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub> blockers) should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub> blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped.



Janssen Pharmaceutica B-2340 Beerse, Belgium

## HYPERTENSION


Hypertension is a common disorder that is easily detected. But it may be deceptively simple: the measurement of blood pressure, interpretation of the results, and the decisions that follow need to be based on a comprehensive knowledge of recent advances.

These illustrated articles on blood pressure measurement, reduction and management have now been collected together in the same format as they appeared in

the BMJ to provide a practical guide for general practitioners and others concerned with hypertension.

Price: Inland £3.50; Overseas US\$16.00\* (Inland £3.00; Overseas US\$14.50\* for BMA members) \*including air mail postage

Order your copy now From: The Publisher, British Medical Journal, BMA House, Tavistock Square, London WC1H 9JR or any leading bookseller



### HOW TO DO IT


Every doctor knows the sinking sensation induced by waiting one's turn for an examination viva and the social anxieties of changing jobs. The misery of first night nerves does not, however, disappear with time: each decade brings new challenges. On these occasions—chairing a committee, giving evidence in court, appearing on television—practical tips can be enormously helpful. The same is true of the many skills not taught in medical school—how to lecture, referee papers, edit, or use a library. All these and many more nuggets of down-to-earth advice have been collected from past issues of the *BMJ* to form a compendium for the novice—whatever his or her age or seniority.

Price: Inland £4.00;
Abroad US\$10.00
(Concessionary price to BMA members: Inland £3.50;
Abroad US\$8.75.

When ordering BMA members must quote their membership number or the full price will be applicable.)

Payment must be enclosed with order or a surcharge of 30p will be made for rendering invoices and statements.

Order your copy now
From: The Publisher
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR
or any leading bookseller



### ENOWSIODDAMES

For the treatment of

staphylococcal infection

- (... (staphylocooti) form focal, often negrotic lesions, where they are protected from the defence mechanism of the body and shielded from efficient chemotherapy.
- Fucidin exerts a potent antibacterial effect even in the presence of large collections of pus, into which (Capparently penetrates in effective concentrations.)
- Fucidia, which has the audity to penetrate in significant assemble into tissues carrying a poor blood supply.

- Brain abscess
- Ilypeigyon
- Endophthalmitis
- Lungabacesa
- Endocestitie
- Phousions with abscess
- Empyéma
- e-Renal carbuncle
- Deep wound infection
- Septic arthetts
- · Poreign bodies
- Osteonyellis
- a Diniselle gangiane

### Oral danage:

Children 20 - 6 - 74 (2) Open Ground Georgian (September 20 - 1900) (1900) (1900) (1900) Annie Ground Georgian (1900) (19

elected to the second

stational (1) is provided that \$20 (00) by