

Nizoral

TRADEMARK

ketoconazole

Oral therapy for all common fungal infections

Vaginal candidosis:	In all dermatological and in systemic fungal infections:
2 tablets once daily (with food) for 5 days	1 tablet daily (with food) until complete symptomatic and mycological cure is obtained

Not all indications are as yet approved in all countries.

PRESCRIBING INFORMATION

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and k/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults, of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.

Side-effects, precautions, contra-indications: contra-indicated in pregnancy. For maximal absorption, Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped.



Janssen Pharmaceutica
B-2340 Beerse, Belgium

TODAY'S TREATMENT/4

The drugs that we use today are increasingly potent, dangerous, and expensive, and every doctor should have some understanding of clinical pharmacology and drug-induced diseases. Both these subjects, which have been badly taught in medical schools, are covered comprehensively in this new book, which consists of articles taken from the *BMJ*. Also included are articles that provide a clear and up-to-the-minute introduction to anaesthetics.

Price: Inland £4.50;
Overseas US\$20.50*
(Inland £4.00;
Overseas US\$19.00* for
BMA members)

**including air mail postage*

When ordering BMA members must quote their membership number or the full price will be applicable.

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller



ABC OF HYPERTENSION

Hypertension is a common disorder that is easily detected. But it may be deceptively simple: the measurement of blood pressure, interpretation of the results, and the decisions that follow need to be based on a comprehensive knowledge of recent advances.

These illustrated articles on blood pressure measurement, reduction and management have now been collected together in the same format as they appeared in the BMJ to provide a practical guide for general practitioners and others concerned with hypertension.


Price: Inland £3.50;
Overseas US\$16.00*
(Inland £3.00;
Overseas US\$14.50* for
BMA members)

**including air mail postage*

Order your copy now
From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR
or any leading bookseller

ABC OF HYPERTENSION

ARTICLES FROM THE BRITISH MEDICAL JOURNAL



Night and day... you are the one



"NIGHT AND DAY" from GAY DIVORCE
Music & Words by Cole Porter
© 1932 Harris Inc. (Warner Bros.)
British Publisher: Chappell Music Ltd.

Voltarol[®]

for osteoarthritis

Presentation Tablets of 25 and 50mg diclofenac sodium; suppositories of 100mg diclofenac sodium. **Indications** Rheumatoid arthritis, osteoarthritis, low back pain and other acute musculo-skeletal disorders, ankylosing spondylitis, acute gout. **Dosage** Tablets: 75-150mg daily in two or three divided doses. Suppositories: one daily, usually at night. In more severe cases, combined therapy with tablets is recommended (daily dose should not exceed 150mg). **Side-effects** Transient epigastric pain, nausea and diarrhoea, headache and slight dizziness have been reported, as occasionally have skin rash, peripheral oedema and abnormalities of serum transaminases and (very rarely) peptic ulcer and haematemesis or melaena. Local reactions to suppositories include itching, burning and increased frequency of bowel movement. **Precautions** Do not prescribe during pregnancy unless there are compelling reasons. Patients with a history of peptic ulcer, haematemesis or melaena, or with severe hepatic or renal insufficiency, should be kept under close surveillance. Voltarol has been reported to depress salicylate levels and vice versa: the clinical relevance of this is not yet clear. Use suppositories only with caution in patients with painful or irritable ano-rectal conditions. **Contra-indications** Peptic ulceration; patients in whom attacks of asthma, urticaria or acute rhinitis are precipitated by aspirin or other NSAIs. **Packs** Tablets 25mg (PL0001/0936) in packs of 100, basic NHS price £9.54; tablets 50mg (PL0001/0082) in packs of 100, £18.55; suppositories 100mg (PL0001/0083) in packs of 10, £3.16. Full prescribing information is available from Geigy Pharmaceuticals, Horsham, West Sussex.

Geigy