



Nizoral

TRACEMARK

ketoconazole

ral therapy for all common fungal infections

Vaginal candidosis:	In all dermatological and in systemic fungal infections:
2 tablets once daily (with food) for 5 days	1 tablet daily (with food) until complete sympto- matic and mycological cure is obtained

Not all indications are as yet
approved in all countries.

PRESCRIBING INFORMATION

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and k/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults, of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.

Side-effects, precautions, contra-indications: contra-indicated in pregnancy. For maximal absorption, Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped.



Janssen Pharmaceutica
B-2340 Beerse, Belgium

STATISTICS AT SQUARE ONE

by T. D. V. SWINSCOW

from the British Medical Journal


The statistical testing of data is indispensable in many types of medical investigation and a help on countless occasions in clinical practice. This book provides step-by-step instruction. Subjects covered include standard deviation, χ^2 tests, t tests, non-parametric tests and correlation. Methods specially adapted to pocket calculators.

"The ability to put symbols in sequence may well be one of the greatest advances in man's development. . . . This excellent and attractive volume will give you an idea into which category an article or projected article fits. The ink you save may be your own. It is superb value."

O'Donnell, Barry, *British Medical Journal*, 1977, 1, 451.

"... this gem of a practical booklet squeezes the essence of biostatistics into 75 well-written, well-organized pages. Newcomers to the field will like the way obscure concepts are clarified; experts will find the book valuable as a teaching aid."

de Jong, Rudolph H., *J.A.M.A.*, 1977, 237, 1874



Seventh edition

**PRICE: Inland £2.50
Overseas US\$10.00,
including despatch by
Air Mail**

*Payment must be enclosed with
order or a surcharge will
be made for rendering invoices
and statements*

ORDER YOUR COPY NOW

From: The Publisher,
British Medical Journal,
BMA House,
Tavistock Square,
London WC1H 9JR

or through any leading bookseller

18th Edition of

BAILEY AND LOVE'S SHORT PRACTICE OF SURGERY

Revised by,

A. J. HARDING RAINS, M.S., F.R.C.S.(Eng.)
and H. DAVID RITCHIE, M.A., Ch.M.,
F.R.C.S.(Ed.), F.R.C.S.(Eng.)

1370 pages 1531 illustrations £23.00 net.
ISBN 0 7186 0450 4

H. K. LEWIS & Co. LTD.
136 GOWER STREET LONDON WC1E 6BS
Tel: 01-387 4282

THE ROYAL COLLEGE OF SURGEONS OF EDINBURGH SPECIALTY FELLOWSHIP IN CARDIOTHORACIC SURGERY – FRCSEd (C/Th)

A diet of the Specialty Fellowship Examination in Cardiothoracic Surgery will be held on 8 November 1983.

Candidates who should normally hold a Diploma of Fellowship of a Surgical College or an equivalent Diploma are required to have three years' post-Fellowship experience in Cardiothoracic Surgery of which one year must have been completed in an approved centre in the United Kingdom. Candidates must submit written evidence of their experience in the specialty including their operative experience.

The application form, examination calendar and Regulations are available on request from the:

**Examinations Secretary,
The Royal College of Surgeons of Edinburgh,
Nicolson Street, Edinburgh EH8 9DW**

Applications for entry must be received by 23 September 1983. Fee: £130.00.

TRY A DIFFERENT PRESCRIPTION

Depressed by your present car?
Nervous about choosing the right replacement?
Call First Front, London's leading BMW centre.
We'll prescribe the ideal tonic.



First Front

289/299 KENNINGTON LANE LONDON SE11 5QY
Tel. 01-582 6000 TELEX: 296957

**THE PROFESSIONAL'S CAR
THE PROFESSIONAL DEALER**

ABC OF HYPERTENSION

Hypertension is a common disorder that is easily detected. But it may be deceptively simple: the measurement of blood pressure, interpretation of the results, and the decisions that follow need to be based on a comprehensive knowledge of recent advances.


These illustrated articles on blood pressure measurement, reduction and management have now been collected together in the same format as they appeared in the BMJ to provide a practical guide for general practitioners and others concerned with hypertension.

Price: Inland £3.50; Overseas US\$16.00*
(Inland £3.00; Overseas US\$14.50* for
BMA members)

**including air mail postage*

Order your copy now

From: The Publisher, British Medical Journal,
BMA House, Tavistock Square, London WC1H 9JR
or any leading bookseller



Burinex

bumetanide

gentle
with
hidden
strength

Prescribing Information:— **Indications** Oedema of renal, cardiac or hepatic origin. **Dosage** Most patients require 1 mg Burinex daily given as morning or evening dose. In refractory cases dosage can be increased to achieve the desired response. For high dose treatment 5 mg Burinex should be given initially and increased by 5 mg steps at 12-24 hour intervals until desired response is achieved. **Contra-indications, Precautions and Side Effects** Contra-indicated in hepatic coma, severe electrolyte depletion and severe progressive

renal failure. Hypovolaemia and circulatory collapse may follow inappropriately excessive diuresis. Electrolyte disturbances resulting in digitalis toxicity may occur. Concurrent antihypertensive or antidiabetic therapy may require adjustment. Caution should be exercised in first trimester of pregnancy. Side effects such as skin rashes, muscular cramps, rises in serum uric acid and thrombocytopenia may rarely occur. **Product Licence Number:** 1 mg tablets 0043/0021. **Basic N.H.S. Price:** £5.60 per 100.



Leo Laboratories Limited
Longwick Road, Princes Risborough
Aylesbury, Bucks

* Burinex is a trade mark