Nizoral™

ketoconazole

oral therapy for all common fungal infections

Vaginal candidosis:	In all dermatological and in systemic fungal infections:
2 tablets once daily (with food) for 5 days	1 tablet daily (with food) until complete symptomatic and mycological cure is obtained

Not all indications are as yet approved in all countries.

PRESCRIBING INFORMATION

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and k/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses: Nizoral** is an orally active antimycotic for the treatment in adults, of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption, **Nizoral** should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H_2 blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after **Nizoral**. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped.

Janssen Pharmaceutica
B-2340 Beerse, Belgium

ABC OF HYPERTENSION

Hypertension is a common disorder that is easily detected. But it may be deceptively simple: the measurement of blood pressure, interpretation of the results, and the decisions that follow need to be based on a comprehensive knowledge of recent advances.

These illustrated articles on blood pressure measurement, reduction and management have now been collected together in the same format as they appeared in the BMJ to provide a practical guide for general practitioners and others concerned with hypertension.

Price: Inland £3.50; Overseas US$16.00*
(Inland £3.00; Overseas US$14.50* for BMA members)

**including air mail postage*

Order your copy now
From: The Publisher, British Medical Journal, BMA House, Tavistock Square, London WC1H 9JR
or any leading bookseller

ABC OF HYPERTENSION

ARTICLES FROM THE BRITISH MEDICAL JOURNAL



AN INTERNATIONAL FORUM FOR MEDICAL RESEARCH AND OPINION



The New England Journal of Medicine

- For over 170 years, the *Journal* has been one of the world's most respected, most quoted sources of new medical knowledge — an international forum for quality research and opinion.
- The *Journal* presents original articles and interpretive reviews of a variety of developments in the major aspects of medicine: its science, its art and practice, and its position in today's society.
- Whatever your medical discipline or the stage of your career, you should subscribe to the *Journal*.

. . . . . .

The New England Journal of Medicine

13 Colina Road, Tottenham
London, England N15 3JA U.K.

All payments remitted directly to NEJM must be in pounds sterling drawn on a London bank or in US dollars drawn on a US bank. **NO RISK GUARANTEE: You may cancel at any time and receive a full refund on all unmailed issues.**

Name _____

Address _____

City _____

Country _____

Postal Code _____

Subscription rates for *52 weekly issues* per annum (please circle):

	Pounds Sterling	U.S. Dollars
PHYSICIANS AND ALL OTHERS		
One year	£ 41	$ 75
Two years	79	145
Three years	117.50	215
POST-GRADUATE PHYSICIANS IN SPECIALTY TRAINING		
One year	£ 30	$ 55
Two years	57	105
Three years	84.50	155
MEDICAL STUDENTS		
Per single year	£ 24.50	$ 45

☐ My remittance is enclosed.* ☐ Please invoice me in advance.
☐ New subscriber. ☐ Current subscriber (please enclose address label).

Professional Specialty: _____

*Please allow 4-6 weeks for shipment of first issue *from the time your payment is received at the NEJM.* Rates are subject to change without notice.

3SBJ1

IN HYPERTENSION AND ANGINA

puts it all together...

Hydrophilic

Reduced risk of drug interaction

Wide range of patients

Few side effects



Cardioselective

Effective anti-anginal

Effective anti-hypertensive

Full 24 hour protection

Increased work performance

...in one tablet daily

TENORMIN

fits the profile of the ideal beta blocker in hypertension and angina

'Tenormin' Prescribing notes:

Presentation: 'Tenormin' tablets containing atenolol 100mg are round, bi-convex, orange and film coated. **Uses:** Management of hypertension and angina pectoris. **Dosage:** Hypertension: One tablet daily. Angina: 100mg daily in single or divided doses. **Contraindications:** Heart block. Co-administration with verapamil. **Precautions:** Untreated cardiac failure, bradycardia, renal failure, anaesthesia and pregnancy. Clonidine withdrawal. **Side Effects:** Coldness of extremities and muscular fatigue. Sleep disturbance rarely seen. Rashes and dry eyes have been reported with beta blockers – consider discontinuance if they occur. Cessation of therapy with beta blockers should be gradual. **Pack size and Basic NHS cost:** 'Tenormin' 28's £7.05. **Product Licence Number:** 'Tenormin' 0029/0122.

Full precribing information is available on request to the Company

Stuart Pharmaceuticals Ltd
Carr House, Carrs Road
Cheadle, Cheshire SK8 2EG