

# Nizoral

TRADEMARK

ketoconazole

## Oral therapy for all common fungal infections

Vaginal candidosis:	In all dermatological and in systemic fungal infections:
2 tablets once daily (with food) for 5 days	1 tablet daily (with food) until complete symptomatic and mycological cure is obtained

Not all indications are as yet approved in all countries.

#### **PRESCRIBING INFORMATION**


**Presentation:** white, flat, half scored uncoated tablets marked "Janssen" on one side and k/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults, of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.

**Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption, Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub> blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped.



**Janssen Pharmaceutica**  
B-2340 Beerse, Belgium

# An Investment in Neuroscience



Dopaminergic neurons in the rat substantia nigra.  
Quantification of dopamine content by coded colours - red colours code the highest dopamine content.


Understanding the role of neuronal plasticity  
as restorative mechanism in nervous system disorders.  
Trying to link progress in neuroscience research  
to human therapeutic benefit.



**FIDIA Research Laboratories**  
P.O. Box 110 · Abano Terme · ITALY · 35031

# IN HYPERTENSION AND ANGINA

## puts it all together...



## ...in one tablet daily

### TENORMIN

**fits the profile of the ideal beta blocker  
in hypertension and angina**

**'Tenormin' Prescribing notes:**

**Presentation:** 'Tenormin' tablets containing atenolol 100mg are round, bi-convex, orange and film coated. **Uses:** Management of hypertension and angina pectoris. **Dosage:** Hypertension: One tablet daily. Angina: 100mg daily in single or divided doses. **Contraindications:** Heart block. Co-administration with verapamil. **Precautions:** Untreated cardiac failure, bradycardia, renal failure, anaesthesia and pregnancy. Clonidine withdrawal. **Side Effects:** Coldness of extremities and muscular fatigue. Sleep disturbance rarely seen. Rashes and dry eyes have been reported with beta blockers—consider discontinuance if they occur. Cessation of therapy with beta blockers should be gradual. **Pack size and Basic NHS cost:** 'Tenormin' 28s £7.05. **Product Licence Number:** 'Tenormin' 0029/0122.

Full prescribing information is available on request to the Company



**Stuart Pharmaceuticals Ltd**  
Carr House, Carrs Road  
Cheadle, Cheshire SK8 2EG

'Tenormin' is a trademark.

