# ketoconazole

# therc

Vaginal candidosis:	In all dermatological and in systemic fungal infections:
2 tablets once daily (with food) for 5 days	1 tablet daily (with food) until complete sympto- matic and mycological cure is obtained

Not all indications are as yet approved in all countries.

PRESCRIBING INFORMATION

PRESCRIBING INFORMATION
Presentation: white, flat, half scored uncoated tablets
marked "Janssen" on one side and k/200 on the reverse.
Each tablet contains 200 mg ketoconazole. Uses: Nizoral is
an orally active antimycotic for the treatment in adults,
of vaginal candidosis, superficial and systemic mycoses
including dermatophyte and yeast infections of the skin,
hair and nails, yeast infections of the mouth and G.I. tract.
Also maintenance treatment of systemic mycoses and
chronic mucocutaneous candidosis and prophylaxis in
at risk" patients in children, systemic mycoses and severe. chronic mucocutaneous candidosis and prophylaxis in at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.

Side-effects, precautions, contra-indications: contra-indicated in pregnancy. For maximal absorption. Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub> blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis treatment with ketoconazole should be stopped.



Janssen Pharmaceutica B-2340 Beerse, Belgium

# STATISTICS by T. D. V. SWINSCOW AT SQUARE ONE

### from the British Medical Journal


The statistical testing of data is indispensable in many types of medical investigation and a help on countless occasions in clinical practice. This book provides step-by-step instruction. Subjects covered include standard deviation,  $\chi^2$  tests, t tests, non-parametric tests and correlation. Methods specially adapted to pocket calculators.

"The ability to put symbols in sequence may well be one of the greatest advances in man's development... This excellent and attractive volume will give you an idea into which category an article or projected article fits. The ink you save may be your own. It is superb value."

O'Donnell, Barry, British Medical Journal, 1977, 1, 451.

"... this gem of a practical booklet squeezes the essence of biostatistics into 75 well-written, well-organized pages. Newcomers to the field will like the way obscure concepts are clarified; experts will find the book valuable as a teaching aid."

de Jong, Rudolph H., J.A.M.A., 1977, 237, 1874



## Eighth edition

PRICE: Inland £3.00
Overseas US\$11.25\*
(Inland £2.50; Overseas
US\$10.00\* to BMA
members)
\*including despatch by
Air Mail


Payment must be enclosed with order

### **ORDER YOUR COPY NOW**

From: The Publisher, British Medical Journal, BMA House, Tavistock Square, London WC1H 9JR

or through any leading bookseller

### AN INTERNATIONAL FORUM FOR MEDICAL RESEARCH AND OPINION



### The New England Journal of Medicine

- For over 170 years, the Journal has been one of the world's most respected, most quoted sources of new medical knowledge — an international forum for quality research and opinion.
- The Journal presents original articles and interpretive reviews of a variety of developments in the major aspects of medicine: its science, its art and practice, and its position in today's society.
- Whatever your medical discipline or the stage of your career, you should subscribe to the Journal.

### The New England Journal of Medicine 13 Colina Road, Tottenham London, England N15 3JA U.K.

All payments remitted directly to NEJM must be in pounds sterling drawn on a London bank or in US dollars drawn on a US bank. NO RISK GUARANTEE: You may cancel at any time and receive a full refund on all unmailed issues.

Name

Address		
City		
Country		
Postal Code	•	
Subscription rates for 52 we (please co		r annum
PHYSICIANS AND ALL OTHE	Pounds RS Sterling	
One year	£ 41	\$ 75
Two years	79	145
Three years	117.50	215
POST-GRADUATE PHYSICIANS	IN SPECIALTY	TRAINING
One year	£ 30	<b>\$</b> 55
Two years	57	105
Three years	84.50	155
MEDICAL STUDENTS		
Per single year	£ 24.50	<b>\$ 45</b>
☐ My remittance is enclosed.* ☐ New subscriber.	Please invoice me in advance.  □ Current subscriber (please enclose address label).	
Professional Specialty:		
*Please allow 4-6 weeks for shipment payment is received at the NEJM. Ra		

# ABC OF HYPERTENSION

Hypertension is a common disorder that is easily detected. But it may be deceptively simple: the measurement of blood pressure, interpretation of the results, and the decisions that follow need to be based on a comprehensive knowledge of recent advances.

These illustrated articles on blood pressure measurement, reduction and management have now been collected together in the same format as they appeared in the BMJ to provide a practical guide for general practitioners and others concerned with hypertension.

Price: Inland £3.50; Overseas US\$16.00\* (Inland £3.00; Overseas US\$14.50\* for BMA members)


\*including air mail postage

Order your copy now


From: The Publisher, British Medical Journal, BMA House, Tavistock Square, London WC1H 9JR or any leading bookseller

# ABC OF HYPERTENSION

ARTICLES FROM THE BRITISH MEDICAL JOURNAL



3SBJ1



Prescribing Information:—Indications Oedema of renal, cardiac or hepatic origin. Dosage Most patients require 1 mg Burinex daily given as morning or evening dose. In refractory cases dosage can be increased to achieve the desired response. For high dose treatment 5 mg Burinex should be given initially and increased by 5 mg steps at 12-24 hour intervals until desired response is achieved. Contraindicated in hepatic coma, severe electrolyte depletion and severe progressive

renal failure. Hypovolaemia and circulatory collapse may follow inappropriately excessive diuresis. Electrolyte disturbances resulting in digitalis toxicity may occur. Concurrent antihypertensive or antidiabetic therapy may require adjustment. Caution should be exercised in first trimester of pregnancy. Side effects such as skin rashes, muscular cramps, rises in serum uric acid and thrombocytopenia may rarely occur. Product Licence Number: 1 mg tablets 0043/0021. Basic N.H.S. Price: £5.60 per 100.



Leo Laboratories Limited Longwick Road, Princes Risborough Aylesbury, Bucks

Burinex is a trade mark