

# Nizoral

TRADE MARK

ketoconazole

## Oral therapy for all common fungal infections

Vaginal candidosis:	In all dermatological and in systemic fungal infections:
2 tablets once daily (with food) for 5 days	1 tablet daily (with food) until complete symptomatic and mycological cure is obtained

Not all indications are as yet approved in all countries.

#### PRESCRIBING INFORMATION

**Presentation:** white, flat, half scored uncoated tablets marked "Janssen" on one side and k/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults, of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.

**Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption, Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub> blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped.



Janssen Pharmaceutica  
B-2340 Beerse, Belgium

# HOW TO DO IT

Every doctor knows the sinking sensation induced by waiting one's turn for an examination viva and the social anxieties of changing jobs. The misery of first night nerves does not, however, disappear with time: each decade brings new challenges. On these occasions—chairing a committee, giving evidence in court, appearing on television—practical tips can be enormously helpful. The same is true of the many skills not taught in medical school—how to lecture, referee papers, edit, or use a library. All these and many more nuggets of down-to-earth advice have been collected from past issues of the *BMJ* to form a compendium for the novice—whatever his or her age or seniority.

Price: Inland £4.00

Overseas US\$10.00\*

(Inland £3.50; Overseas US\$8.75\*  
for BMA members)

\*including air mail postage

Payment must be enclosed with order

**Order your copy now**

From: The Publisher,  
British Medical Journal  
BMA House  
Tavistock Square  
London WC1H 9JR  
or any leading bookseller

## HOW TO DO IT

Chair a committee.  
Be a dictator.  
Take an examination.  
Be a reporter.  
Organise an international conference.  
Improve a student.  
Chair a medical journal.  
Interviewed.  
Survive as an examiner.  
Construct an auditor.  
Attend an inquest.  
Examine on television.  
Research grant.  
Plan a research project.  
Write a paper.  
Use a library.  
Give a reference.  
Take a clinical programme.  
Apply slides.  
Use an overhead projector.  
Use a research project.

"... a tribute to their excellence that on rereading they did not pall. It will make a useful present for one's favourite houseman or registrar."  
*NZ Med J* 1980; 91: 73

"It is a joy to read... All in all, this book can be highly recommended."  
*Med J Aust* 1980; 1: 680

# THE ESSENTIAL CORRESPONDENCE COURSE FOR THE MRCP

**This new course for PART I and PART II has been completely rewritten and updated with new clinical slides, X-rays and ECG's included.**

**Write for full details to:**

**J. ARNOLD  
(Correspondence Courses)  
19 VILLAGE WAY,  
LONDON SE21 7AN**

## ABC OF HYPERTENSION

Hypertension is a common disorder that is easily detected. But it may be deceptively simple: the measurement of blood pressure, interpretation of the results, and the decisions that follow need to be based on a comprehensive knowledge of recent advances.

These illustrated articles on blood pressure measurement, reduction and management have now been collected together in the same format as they appeared in the BMJ to provide a practical guide for general practitioners and others concerned with hypertension.

Price: Inland £3.50; Overseas US\$16.00\*  
(Inland £3.00; Overseas US\$14.50\* for  
BMA members)

*\*including air mail postage*

**Order your copy now**

From: The Publisher, British Medical Journal,  
BMA House, Tavistock Square, London WC1H 9JR  
or any leading bookseller

## How Will You Stay Informed When The Next "AIDS"-Like Health Emergency Strikes?

Researchers at the United States Centers for Disease Control first communicated their findings on AIDS in its *Morbidity and Mortality Weekly Report*. *MMWR* is a unique source of critical health information which impacts the world.

*MMWR* features up-to-date information on worldwide epidemiological and public-health developments. This concise, easy to read bulletin provides you with the most current guidelines concerning treatment of transmittable and reportable diseases.

Avoid getting lost in the void between immediate lay press coverage and the delayed formal reporting of scientific articles. *MMWR* is a vital communication link with the medical community.

From AIDS to Toxic Shock Syndrome, from Legionnaire's Disease to Swine Flu, *MMWR* can make the difference for you and your patients when *timely, accurate* information is essential.

*Morbidity and Mortality Weekly Report* is published by the United States Centers for Disease Control in Atlanta, Georgia. The Massachusetts Medical Society, publisher of *The New England Journal of Medicine*, is reprinting and distributing this publication as a public service on a non-profit basis.

### ORDER FORM

**Morbidity and Mortality Weekly Report  
MMS Publications, C.S.P.O. Box 9120  
Waltham, Massachusetts 02254 USA**

All payments must be remitted in U.S. dollars drawn on a U.S. bank. Payments in any other form will be returned, causing delays. You may write to us for the name of a subscription agency representative in your country if you wish to remit in your own currency.

Name \_\_\_\_\_

Address \_\_\_\_\_

City \_\_\_\_\_

Country \_\_\_\_\_ Postal Code \_\_\_\_\_

Professional Specialty: \_\_\_\_\_

*MMWR* is available on a one year basis only. ALL ORDERS MUST BE PREPAID. Make checks payable to: MMS Publications.

### SUBSCRIPTION RATES — U.S. Funds On A U.S. Bank Only

In addition to the weekly publication, *MMWR* expects to produce 4 Quarterly Surveillance Reports and between 2 and 4 Supplements during the course of a yearly subscription, which are also available from MMS Publications for a small additional fee.

(check one please):

International Air Mail: ☐ 52 issues & Annual — \$75  
☐ with Supplements — \$85 (11f)

Please allow 4-6 weeks from the time your order is received until the first mailing. NO RISK GUARANTEE: You may cancel at any time and receive a full refund on all unmailed issues.

S3B01



# Burinex<sup>®</sup>

bumetanide

gentle  
with  
hidden  
strength

**Prescribing Information:**—**Indications** Oedema of renal, cardiac or hepatic origin. **Dosage** Most patients require 1 mg burinex daily given as morning or evening dose. In refractory cases dosage can be increased to achieve the desired response. For high dose treatment 5 mg Burinex should be given initially and increased by 5 mg steps at 12-24 hour intervals until desired response is achieved. **Contra-indications, Precautions and Side Effects** Contra-indicated in hepatic coma, severe electrolyte depletion and severe progressive

renal failure. Hypovolaemia and circulatory collapse may follow inappropriately excessive diuresis. Electrolyte disturbances resulting in digitalis toxicity may occur. Concurrent antihypertensive or antidiabetic therapy may require adjustment. Caution should be exercised in first trimester of pregnancy. Side effects such as skin rashes, muscular cramps, rises in serum uric acid and thrombocytopenia may rarely occur. **Product Licence Number:** 1 mg tablets 0043/0021. **Basic N.H.S. Price:** £5.80 per 100.



Leo Laboratories Limited  
Longwick Road, Princes Risborough  
Aylesbury, Bucks

\* Burinex is a trade mark