



The modern approach:

TRADEMARK

Nizoral

ketoconazole

**just 2 oral tablets daily
with a meal
for 5 days.**

adults have some problem
with *Candida* vaginitis

Reluctance to apply topical treatment

the patient's dislike of topical application
leads to inadequate therapy

Prescribing Information

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole.

Uses: Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophytes and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I.-tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.

Side-effects, precautions, contra-indications:

contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritis may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.



JANSSEN
PHARMACEUTICA
B-2340 Beerse, Belgium

AVIATION MEDICINE

Richard M Harding & F John Mills


This authoritative and highly readable account of the little known specialty of aviation medicine will interest both general readers and those interested in it as a career. Comprehensive information on the management of airline passengers with particular medical conditions before, during, and after flight will help doctors faced with deciding whether or not their patient is fit to fly and with coping with a medical emergency in the air while themselves passengers on board an aircraft. The chapters on the problems of altitude, acceleration, the function of the special senses in flight, special forms of flight, and aviation psychology describe clearly and precisely the physiological and behavioural effects that flight imposes on man.

Price: Inland £5.50;
Overseas £7.50/USA\$13.00
(Inland £5.00;
Overseas £7.00/USA\$12.00 to
BMA members)

Despatched by air overseas

Payment must be enclosed with order

Order your copy now
From: The Publisher
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR
or any leading bookseller



Just published

ABC OF SEXUALLY TRANSMITTED DISEASES


MICHAEL W ADLER

AIDS and genital herpes are only the more dramatic signs of a general increase in the number and range of sexually transmitted diseases. Yet these conditions are not seen only in sexually transmitted disease clinics and doctors need to be aware that common clinical problems such as urinary tract infections, rashes, and pelvic pain may have a sexual origin. Professor Michael Adler's ABC describes the presentation, diagnosis, and management of these conditions, emphasising the need to take a broad clinical view of patients and their problems.

Price: Inland £4.00;
Overseas £6.00/USA\$10.50
(BMA members: Inland £3.50;
Overseas £5.50/USA\$9.50)
Despatched by air overseas

Payment must be enclosed with order

Order your copy now
From: The Publisher
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR
or any leading bookseller




**Who's afraid of
going upstairs?**

Your angina patients

Transiderm®-Nitro
glyceryl trinitrate

**One patch a day for the prophylaxis
of angina attacks**

Prescribing Notes: **Presentation** Transiderm-Nitro 5 and 10 are transdermal drug delivery systems, comprising respectively 10cm² and 20cm² self-adhesive, pink-coloured patches, containing a drug reservoir of glyceryl trinitrate. **Indication** Prophylactic treatment of attacks of angina pectoris, as monotherapy or in combination with other anti-anginal agents. **Dosage** One Transiderm-Nitro 5 patch is to be applied every 24 hours. If a higher dose is required a Transiderm-Nitro 10 patch may be substituted. It is recommended that the patch is applied to the lateral chest wall. The patch should be removed after 24 hours, and the replacement patch applied to a new area of skin. Allow several days to elapse before applying a fresh patch to the same area of skin. If acute attacks of angina pectoris occur, rapidly acting nitrate preparations may be required. Efficacy and tolerability beyond 28 days' therapy have yet to be established. **Side-effects** Headache may occur and usually regresses after a few days. Reflex tachycardia can be controlled by concomitant treatment with a beta-blocker. Postural hypotension, nausea and dizziness occur rarely. Allergic skin reactions, a local mild itching or burning sensation may occasionally occur. Upon removal of the patch, any slight reddening of the skin will usually disappear in a few hours. **Precautions** In recent myocardial infarction or acute heart failure, Transiderm-Nitro should be employed only under careful clinical surveillance. As with all anti-anginal nitrate preparations, withdrawal of treatment should be gradual, by replacement with decreasing doses of long-acting oral nitrates. **Contra-contraindications** Hypersensitivity to nitrates, severe hypotension, marked anaemia, increased intraocular or intracranial pressure. **Packs** Boxes of 30 patches. Transiderm-Nitro 5 (PL0001/0094), basic NHS price 64.4p per day. Transiderm-Nitro 10 (PL0001/0095), basic NHS price 70.9p per day. *denotes registered trademark. Full prescribing information is available on request from Geigy Pharmaceuticals, Horsham, West Sussex.



Geigy