


Nizoral^{TRADEMARK}

(ketoconazole) tablets

over 3 million
prescriptions
world-wide:

the new beginning
in antifungal therapy.



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal in-

fections can now be treated effectively and conveniently. Typically, in *Candida vaginitis*, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.

 **JANSSEN**
PHARMACEUTICA

the drug discovery company

Prescribing Information

Presentation: white, flat, half-scored, uncoated tablets marked Janssen on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole.

Uses: Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children, systemic mycoses and severe local infections where previous topical treatment has failed.

Side-effects, precautions, contra-indications: contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 3 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild, asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynecomastia has been reported.

ABC OF SEXUALLY TRANSMITTED DISEASES


MICHAEL W ADLER

AIDS and genital herpes are only the more dramatic signs of a general increase in the number and range of sexually transmitted diseases. Yet these conditions are not seen only in sexually transmitted disease clinics and doctors need to be aware that common clinical problems such as urinary tract infections, rashes, and pelvic pain may have a sexual origin. Professor Michael Adler's ABC describes the presentation, diagnosis, and management of these conditions, emphasising the need to take a broad clinical view of patients and their problems.

Price: Inland £4.00;
Overseas £6.00/USA\$10.50
(BMA members: Inland £3.50;
Overseas £5.50/USA\$9.50)
Despatched by air overseas


Payment must be enclosed with order

Order your copy now
From: The Publisher
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR
or any leading bookseller



IN HYPERTENSION AND ANGINA

...in one tablet daily



...in one tablet daily

TENORMIN

fits the profile of the ideal beta blocker
in hypertension and angina

'Tenormin' Prescribing notes:

Presentation: 'Tenormin' tablets containing atenolol 100 mg are round, bi-convex, orange and film coated. **Uses:** Management of hypertension and angina pectoris. **Dosage:** Hypertension: One tablet daily. Angina: 100 mg daily in single or divided doses. **Contraindications:** Heart block. Co-administration with verapamil. **Precautions:** Untreated cardiac failure, bradycardia, renal failure, anaesthesia and pregnancy. Clonidine withdrawal. **Side Effects:** Coldness of extremities and muscular fatigue. Sleep disturbance rarely seen. Rashes and dry eyes have been reported with beta blockers—consider discontinuance if they occur. Cessation of therapy with beta blockers should be gradual. **Pack size and Basic NHS cost:** 'Tenormin' 28's £6.98 **Product Licence Number:** 'Tenormin' 0029/0122.

Full prescribing information is available on request to the Company



Stuart Pharmaceuticals Ltd
Carr House, Carrs Road
Cheadle, Cheshire SK8 2EG

'Tenormin' is a trademark :

