


# Nizoral

(ketoconazole) tablets

over 3 million  
prescriptions  
world-wide:

the new beginning  
in antifungal therapy.



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal in-

fections can now be treated effectively and conveniently. Typically, in *Candida vaginitis*, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.

 **JANSSEN**  
PHARMACEUTICA

the drug discovery company

**Prescribing Information**

**Presentation:** white, flat, half scored uncoated tablets marked Janssen on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole.

**Uses:** Nizoral is an orally active antifungal for the treatment in adults of vaginal candidosis, superficial and systemic mycoses, including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in at risk patients. In children, systemic mycoses and severe local infections where previous topical treatment has failed.

**Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption, Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub>-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynecomastia has been reported.




# SURGE

tiaprofenic acid

live life

# When a patient comes to you with flu, maybe you should recommend Flora.

Perhaps the best time to start your patients thinking more seriously about their health is when they come to you with some more routine complaint.



It is now widely accepted that changes in diet and lifestyle can dramatically decrease the risk of Coronary Heart Disease developing.

So the best time to recommend Flora is before, not after, any symptoms of CHD present themselves.

Flora is high in essential polyunsaturated fats (*cis-cis* linoleic), because it is made with pure sunflower oil.

Whenever an opportunity arises to recommend ways in which your patients could reduce the CHD risk factors, recommend they change to Flora, too.

**Flora. High in essential polyunsaturates.**



# THE FIRST YEAR OF LIFE

Second edition

H B Valman

Bernard Valman's articles on the first year of life, first published in the *BMJ* in 1979 and then reproduced as a book, have proved invaluable to many doctors and others concerned in the care of children. The second edition of *The First Year of Life* includes numerous small changes to bring the book up to date and incorporates a much fuller section on whooping cough. The section on contraindications to immunisation has also been revised to take account of the latest guidance.

Price: Inland £5.50

Overseas £8.00/USA\$14.00

(Inland £5.00; Overseas £7.50/USA\$13.00 to BMA members)

Despatched by air overseas

Payment must be enclosed with order

Order your copy now


From: The Publisher

British Medical Journal, BMA House,  
Tavistock Square, London WC1H 9JR  
or any leading bookseller

## THE FIRST YEAR OF LIFE

H B VALMAN

ARTICLES FROM THE BRITISH MEDICAL JOURNAL



## THE MRCP CORRESPONDENCE COURSE IN INTERNAL MEDICINE

- \* **Continually updated**
- \* **New clinical slides**
- \* **Includes X-rays and ECG's**
- \* **Essential for Part I and Part II**

Write for full information to:

**J. ARNOLD Dept 5**  
**(Correspondence Courses)**  
**19, VILLAGE WAY,**  
**LONDON SE21 7AN**

## ALCOHOL PROBLEMS

In recent years alcohol problems have increased dramatically and the thinking on them has undergone a revolution. *Alcohol Problems* brings together two series of articles published in the *BMJ*—the ABC of Alcohol, with its emphasis on straightforward advice for the clinician, and *Alcohol and Alcoholism*, Dr Richard Smith's more discursive survey of current thinking and controversies. Together they cover both the clinical aspects of managing alcohol problems and the social and political factors that surround them.

Price: Inland £5.50

Overseas £8.00/USA\$14.00

(Inland £5.00; Overseas £7.50/USA\$13.00 to BMA members)

Despatched by air overseas

Payment must be enclosed with order

Order your copy now

From: The Publisher

British Medical Journal, BMA House,  
Tavistock Square, London WC1H 9JR  
or any leading bookseller

# THE CARDIOPROTECTIVE CORONARY THERAPEUTIC AGENT ADALAT®

THE MODERN COMPREHENSIVE DRUG THERAPY OF CORONARY HEART DISEASE.



**Indications:** Treatment and prophylaxis of acute and chronic coronary insufficiency (especially angina pectoris, postinfarction syndrome). **Side effects:** Adalat is well tolerated. Side effects occur rarely. They are more common at the beginning of treatment, usually transient, and, in most cases, mild. Occasionally there can be headache, facial flush, heat sensation, dizziness, nausea and tiredness. Extensive biochemical studies did not reveal any abnormal values which could have been ascribed to the drug. **Contraindication:** Pregnancy. **Precaution:** Adalat can enhance or supplement the action of hypotensive preparations and  $\beta$ -blocking agents. Therapy with cardiac glycosides can be initiated or continued in association with Adalat treatment. Incompatibility with other drugs has not been found. **Dosage and administration:** In longterm treatment the capsule is usually swallowed whole with a little liquid. The average daily dose is 3 x 1 capsule, in special cases up to 3 x 2 capsules. In these patients the interval between the individual doses should not be less than two hours. In case of imminent anginal attack, a rapid onset of action is achieved by biting the capsule. **Composition:** One capsule contains 10 mg 1,4-dihydro-2,6-dimethyl-4-(o-nitrophenyl)-pyridine-3,5-dicarboxylic acid dimethyl ester (nifedipine). **Presentation:** Box of 30 or 50 capsules.

Bayer Germany

THE CARDIOPROTECTIVE CORONARY THERAPEUTIC AGENT



E 4/043

