

Nizoral^{TRADEMARK}

(ketoconazole) (tablets)

over 3 million prescriptions world-wide:

the new beginning in antifungal therapy.



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but problematic fungal infections

can now be treated effectively and conveniently. Typically, in recurrent *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



the drug discovery company

Prescribing Information

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole.

Uses: Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.

Side-effects, precautions, contra-indications: contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynecomastia has been reported.



Heceta lighthouse, Oregon, USA

Prescribing Information

Presentation Vials containing 500mg, 1g or 2g of cefotaxime sodium. **Indications** Infections before identification of the organism. Infections caused by bacteria of established sensitivity, including chest infections, septicaemia, urinary tract infections, soft tissue infections, obstetric and gynaecological infections, bone and joint infections, meningitis, gonorrhoea. **Dosage** Claforan is administered i.m. or i.v. **Adults:** Usually 2-6g daily (see full prescribing information). For infections caused by sensitive *Pseudomonas* spp., doses of more than 6g daily are usually required. **Children:** 100-150mg/kg/day in 2 to 4 divided doses. Up to 200mg/kg/day may be given in very severe infections. **Neonates:** 50mg/kg body weight daily in 2 to 4 equally divided doses. In cases of severe infection, divided daily doses of 150-200mg/kg have been given. **Dosage in renal impairment** Reduced dosage is only required in severe renal failure (GFR < 5ml/min = serum creatinine approx. 751 µmol/l) when, after an initial loading dose of 1g, the daily dose is halved without change in frequency of dosing. **Contra-indications** Known allergy to cephalosporins. **Precautions** Cephalosporin antibiotics may usually be given safely to patients who

are hypersensitive to penicillins, although cross reactions have been reported. Special care is indicated in patients who have had an anaphylactic response to penicillin. Patients with severe renal dysfunction — see previous. Cephalosporin antibiotics at high dosage should be given with caution to patients receiving aminoglycoside antibiotics or potent diuretics such as frusemide. At recommended doses, enhancement of nephrotoxicity is unlikely with Claforan. A false-positive reaction to glucose may occur with reducing substances. Claforan should not be mixed in the syringe with aminoglycoside antibiotics. The safety of Claforan in human pregnancy has not been established. **Side effects** Adverse reactions are rare and generally mild and transient, but include diarrhoea (pseudomembranous colitis has been rarely reported), candidiasis, rashes, fever, eosinophilia, leukopenia, transient rises in liver transaminase and alkaline phosphatase, transient pain at the site of injection and phlebitis. **Product licence holder and number** Roussel Laboratories Ltd., Broadwater Park, North Orbital Road, Uxbridge, Middlesex UB9 5HP. 0108/0074. **Package quantities and basic N.H.S. price** Vials of 500mg, 1g and 2g in packs of 10. One gram vial £4.95. **Date of preparation** November 1984.

ROUSSEL



35 Boulevard des Invalides,
75007 Paris, Cedex 07, France

cefotaxime
Claforan
Roussel Laboratories Ltd.

Publication date 21 May 1985

APPROPRIATE TECHNOLOGY

Many of the difficulties of Third World medicine—the effects of poverty, overpopulation, and political unrest—are beyond the individual doctor's control; but what the western trained doctor practising in the developing world can and must do is to adapt his practice to compensate for the absence of sophisticated equipment, trained staff, and expensive drugs. It is wasteful, for instance, to buy elaborate machinery if there is no reliable electricity supply to run it off, or order expensive orthopaedic aids from abroad when, with a bit of ingenuity, effective ones can be made from local materials.


Written by doctors and technicians with first hand experience of the problems, *Appropriate Technology* is a guide to the sensible choice and use of resources, including the purchase and maintenance of operating theatre and laboratory equipment and specialist instruments for cardiology and gastroenterology. It sets out the practicalities of immunisation, rehydration, and transfusion; anaesthetics; obstetric care and child health; sanitation; and the purchase and distribution of drugs, as well as describing workable methods of health education and the use of teaching aids; of maintaining medical records; and of conducting epidemiological surveys and research at low cost.

The British Medical Journal has subsidised this title to make this information widely available at as low a cost as possible.

Price: Inland £2.50;
Overseas £5.00/USA \$8.00
(Inland £2.00;
Overseas £4.50/USA \$7.00
to BMA members), including
postage by air overseas

Please quote membership number

Payment must be enclosed with order



Order your copy now from: The Publishing Manager, BRITISH MEDICAL JOURNAL, BMA House, Tavistock Square, London WC1H 9JR, or any leading bookseller.

H. K. LEWIS of GOWER STREET, LONDON

BRITAIN'S SPECIALIST MEDICAL BOOK CENTRE



MEDICAL BOOKSHOP

Catalogues sent post free on request
(Please state particular interests)

AND LIBRARY

Annual Subscription from £10.50
(available in U.K. only)
(Reduced rates for multiple subscription)
Prospectus post free on request

H. K. LEWIS & Co. Ltd., 136 GOWER STREET, LONDON WC1E 6BS
Telephone: 01-387 4282 Telegrams: Publicavit, London W.C.1

19th Edition

BAILEY & LOVE'S SHORT PRACTICE OF SURGERY

Revised by

A. J. HARDING RAINS, MS, FRCS

and

H. DAVID RITCHIE, MS, ChM, FRCS

with specialist contributors

1300 pages

1223 illustrations

£28.00 net.

ISBN 0 7186 0477 6

H. K. LEWIS & CO. LTD

136 GOWER STREET, LONDON WC1E 6BS
01-387-4282

Now Available

Medicine and Psychiatry

A practical approach

Edited by **Francis Creed** and **Jeremy Pfeffer**

"...an impressive collection of information on the interaction of mind and body..."

Hospital Update

"...deserves to be read..."

The Lancet

1985/Paperback/488pp/0 272 79820 7/£14.95

**NEW
IN
PAPERBACK**

Atheroma and Thrombosis

Edited by **V V Kakkar**

A comprehensive review by leading authorities providing a wealth of up-to-date information from a broad variety of clinical and basic studies.

1985/Cased/350pp/0 272 79816 9/£25.00

Proceedings of the European Dialysis and Transplant Association 21

Edited by **A M Davison** and **P J Guillou**

Detailed statistical reports on dialysis and transplantation together with comprehensive reviews of current practice will prove invaluable to those interested in renal diseases.

1985/Cased/1200pp/0 272 79804 5/£45.00

Evaluation and Treatment of Chronic Pain

Edited by **Gerald Aronoff**

An essential reference for all physicians called upon to evaluate and treat chronic pain patients.

1985/Cased/576pp/0 8067 0151 X/£50.00

Imaging Anatomy of the Head and Spine

H N Schnitzlein and **F R Murtagh**

This superb colour atlas presents a correlative display of horizontal, coronal and sagittal sections of the head and spine with a wide variety of MRI scans, CT scans and conventional radiographs.

1985/Cased/336pp/0 8067 1881 1/£149.95

All prices subject to change without notice
Available from your bookseller, or in case of difficulty from
Order Department, Pitman Publishing Ltd,
128 Long Acre, London WC2E 9AN

Pitman

EMPLOYING STAFF

NORMAN ELLIS

ARTICLES FROM THE BRITISH MEDICAL JOURNAL

In the past 20 years employment law has become more complex, and most of the new laws are relevant to general practitioners and other employers in small businesses. "Ignorance is no defence", and employers breaching the law may pay heavily for their ignorance. This book brings together for the first time the principles of employment law relevant to small businesses. The problems covered include: writing the employment contract; dismissal; recruiting and training staff; sick pay; maternity rights; and all other relevant topics. The information is comprehensive, reliable, and clearly presented, and the topic is now thought so important that questions on it are included in the examination to become a member of the Royal College of General Practitioners.

Price: Inland £5.00; Overseas £7.00/USA\$10.50

(Inland £4.50; Overseas £6.50/USA\$9.50 to BMA members, please quote membership number).

Including postage, by air overseas

Payment must be enclosed with order

Order your copy now

From: The Publishing Manager, British Medical Journal,
BMA House, Tavistock Square, London WC1H 9JR
or any leading bookseller

EMPLOYING STAFF
NORMAN ELLIS

HOW TO DO IT

SECOND EDITION

The first edition of HOW TO DO IT proved a useful and popular guide to those things a doctor needs to know but is rarely taught: how to take an examination, how to interview and be interviewed, how to plan and write up research, how to behave at an inquest. In the second edition the original chapters have been expanded and updated, and there are several more chapters on new challenges—choosing a computer, flying, holding a press conference—and some on older ones not included in the first edition—assessing a job, dealing with a publisher.

Price: Inland £6.50; Abroad £9.00/USA\$14.50
BMA members (please quote membership number):
Inland £6.00; Abroad £8.50/USA\$13.50
including postage, by air abroad

Payment must be enclosed with order

Order from: The Publishing Manager
British Medical Journal, BMA House, Tavistock Square
London WC1H 9JR or any leading bookseller

THE ESSENTIAL CORRESPONDENCE COURSE FOR THE M.R.C.P.

This new course for PART I and PART II has been completely re-written and updated with new clinical slides, X-rays and ECG's included.

**Write for full information to:
J. ARNOLD Dept 1
(Correspondence Courses)
19, VILLAGE WAY,
LONDON SE21 7AN**

AS YOU WERE

VE DAY—A MEDICAL RETROSPECT

Elston Grey-Turner, the much loved former secretary of the BMA who died in 1984, was often referred to by colleagues as "The Colonel" in tribute to his military service in the RAMC and the Territorial Army. As a tribute to his memory the *BMJ* commissioned a collection of reminiscences by doctors of their experiences in the second world war and their feelings as it came to an end. For some, VE Day was a time for celebration, but others were too busy to notice or, as prisoners of war, did not even know that it had happened. The exigencies of war brought enormous advances in surgery and medicine—particularly in the use of blood transfusion and penicillin—while in the civilian hospitals newly qualified doctors and medical students took on responsibilities that are almost unimaginable today.

Often moving, occasionally horrific, sometimes hilarious, these highly personal memories reflect the many aspects of war from a medical viewpoint.

This book is published as a memorial to Elston Grey-Turner, profits from its sale going to BMA charities

Price: Inland £6.00;
Abroad £8.75/USA\$12.50
BMA members: Inland £5.50
Abroad £8.25/USA\$11.50
including postage, by air abroad
Please quote membership number

Payment must be enclosed with order

Order from
The Publishing Manager
British Medical Journal
BMA House, Tavistock Square
London WC1H 9JR
or any leading bookseller

THE CARDIOPROTECTIVE CORONARY THERAPEUTIC AGENT ADALAT:

THE MODERN COMPREHENSIVE DRUG THERAPY OF CORONARY HEART DISEASE.



Indications: Treatment and prophylaxis of acute and chronic coronary insufficiency (especially angina pectoris, postinfarction syndrome). **Side effects:** Adalat is well tolerated. Side effects occur rarely. They are more common at the beginning of treatment, usually transient, and, in most cases, mild. Occasionally there can be headache, facial flush, heat sensation, dizziness, nausea and tiredness. Extensive biochemical studies did not reveal any abnormal values which could have been ascribed to the drug. **Contra-indication:** Pregnancy. **Precaution:** Adalat can enhance or supplement the action of hypotensive preparations and β -blocking agents. Therapy with cardiac glycosides can be initiated or continued in association with Adalat treatment. Incompatibility with other drugs has not been found. **Dosage and administration:** In longterm treatment the capsule is usually swallowed whole with a little liquid. The average daily dose is 3 x 1 capsule, in special cases up to 3 x 2 capsules. In these patients the interval between the individual doses should not be less than two hours. In case of imminent anginal attack, a rapid onset of action is achieved by biting the capsule. **Composition:** One capsule contains 10 mg 1,4-dihydro-2,6-dimethyl-4-(o-nitrophenyl)-pyridine-3,5-dicarboxylic acid dimethyl ester (nifedipine). **Presentation:** Box of 30 or 50 capsules.

Bayer Germany

THE CARDIOPROTECTIVE CORONARY THERAPEUTIC AGENT



E 4/043

