

TRADEMARK

Nizoral

(ketoconazole)

(tablets)

over 3 million prescriptions world-wide:

**the new beginning
in antifungal therapy.**



Doctors and patients around the world are discovering the modern simplicity of **Nizoral** oral therapy. Common but problematic fungal infections can now be treated effectively *and* conveniently. Typically, in recurrent *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.

 **JANSSEN**
PHARMACEUTICA

the drug discovery company

Prescribing Information
Presentation: white, flat, half scored uncoated tablets marked Janssen on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole.
Uses: **Nizoral** is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and GI-tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in 'at risk' patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.
Side-effects, precautions, contraindications & contra-indicated in pregnancy. For maximal absorption **Nizoral** should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after **Nizoral**. Nausea, skin rash, headache and pruritis may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.

NEW FROM PROUS SCIENCE

Journal of Serology

First issue: September, 1985

International quarterly journal specialising in the serology of infectious diseases, connective tissue and autoimmune disorders.

CALLING FOR ORIGINAL PAPERS on the currently available serodiagnostic tests or describing new techniques of serodiagnostic potential. Sections include reviews on serological tests applicable to the clinical laboratory and recent developments in biotechnology applied to examining the serology of these diseases.

For further details please contact: The Editor, Dr. J.P. Burnie, Dept. of Medical Microbiology, The London Hospital, Turner Street, London E1 2AD.


THE MRCP CORRESPONDENCE COURSE IN INTERNAL MEDICINE

- *Continually updated
- *New clinical slides
- *Includes X-rays and ECG's
- *Essential for Part I and Part II

Write for full information to:
J. ARNOLD Dept 5
(Correspondence Courses)
19, VILLAGE WAY,
LONDON SE21 7AN



Medical Books from Butterworths



Case Presentations in Heart Disease

Alan Mackintosh

This concise and informative book involves the reader directly in the clinical analysis and the decision making about patients suffering from heart disease.

176pp 210 x 130mm 0 407 00541 2 Softcover £6.95

Case Presentations in Gastrointestinal Disease

W D W Reese, S Hughes, J L Shaffer and G R Barclay

A practical approach to clinical gastroenterology containing 60 case presentations of disorders affecting the oesophagus, gastrointestinal tract, liver, biliary tree and pancreas.

224pp 210 x 130mm 0 407 00542 0 Softcover £6.95

Disease, Laboratory Data and Diagnosis

V Schwarz and G M Addison

This collection of 20 carefully chosen case presentations teaches the student how to interpret the case history, clinical examination and laboratory findings to achieve an accurate diagnosis.

160pp 210 x 130mm 0 407 00540 4 Softcover £4.95

The Epilepsies

BIMR Neurology Volume 5

Roger J Porter and Paolo L Morselli

Written by leading authorities in their own field, this volume contains the latest developments in both scientific background and advances in medical management.

488pp 234 x 165mm 0 407 02298 8 Hardcover £36.00

Epilepsies of Childhood

Second Edition

Niall V O'Donohue

Professor O'Donohue's account of childhood epilepsies has been thoroughly updated for this second edition. Particular attention has been paid to new diagnostic techniques and drug treatments.

288pp 234 x 165mm 0 407 00272 3 Hardcover £27.50

Management of Neurological Disorders

Second Edition

Bryan Ashworth and Michael Saunders

This second edition meets the needs of non-specialists faced with the task of managing neurological patients. The emphasis remains on practical guidance to the management of common problems, offering a perspective of total care.

340pp 216 x 138mm 0 407 00310 X Softcover £14.95

Paediatric Neurology

Second Edition

Ingrid Gamstorp

The new edition of this well known text takes into account the widespread use of new investigative techniques and the introduction of many new drugs for almost every neurological condition.

416pp 234 x 165mm 0 407 00263 4 Hardcover £35.00



Butterworths

Borough Green, Sevenoaks, Kent TN15 8PH

ADALAT® RETARD: ANTIHYPERTENSIVE WITH CARDIOPROTECTIVE EFFECT

Long term therapy with Adalat retard

- reduction in peripheral vascular resistance
- reduction in high blood pressure (reliable reduction in pathologically high blood pressure)

- permanent relief of the heart
- improvement of the myocardial oxygen balance
- protection of cell against unphysiological noxious Ca^{2+} concentrations in case of a pathological ion overflow.

THE MODERN COMPREHENSIVE DRUG THERAPY OF CARDIOVASCULAR DISEASE: ADALAT RETARD.

 Composition: 1 tablet Adalat® retard contains 20 mg Nifedipine. Indications: Coronary heart disease. Early and long-term treatment of coronary heart disease (in particular coronary insufficiency, angina pectoris, post-infarction syndrome). All forms of hypertension. Contraindication: Pregnancy. Side effects: Side effects occur only rarely and, if at all, at the beginning of treatment; they are usually mild and transient. Occasional symptoms may be headache, facial flush and heat sensation, leg oedema, nausea, dizziness, tiredness and skin reactions. As is the case with other vasoactive substances, chest pain may very occasionally occur after administration. In this event Adalat should be discontinued if the pains are considered to have been caused by the medicament. Preparation: Adalat retard is administered in the morning. The treatment should be continued until the side effects have been taken into account. Treatment with cardiac glycosides can be begun or continued during nifedipine treatment. A combination treatment of Adalat retard with beta blockers or a saluretic is possible (see S.M.L.). • Dosage: Treatment should be adjusted individually to the degree of severity of the disease and to the patient's responsiveness. A daily dose of 2 x 1 tablet of Adalat retard (2 x 20 mg) is recommended. In individual cases an increase in the daily dose to 2 x 2 tablets of Adalat retard (2 x 40 mg) may be necessary. The recommended interval between tablet intake is about 12 hours and should not be shorter than 4 hours. Generally, the tablet is swallowed unchewed with a little liquid, independent of meals. Note: In the presence of coronary spasms (Prinzmetal angina, and the least in the presence of a history of coronary heart disease) the risk of imminent myocardial attack or acute hypertensive crisis which all require a quick onset of effect. Adalat capsules which are marked by rapid action should be taken chewed. When the patient's condition has improved, a change-over to Adalat retard tablets may be attempted. Presentation: Tablets of 20 mg Nifedipine. Box of 30 tablets, box of 50 tablets, box of 100 tablets, hospital-size pack.

Bayer Germany

E 4/042

