


Nizoral^{TRADEMARK}

(ketoconazole) (tablets)

over 3 million
prescriptions
world-wide:

the new beginning
in antifungal therapy.



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but problematic fungal infections

can now be treated effectively and conveniently. Typically, in recurrent *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



JANSSEN
PHARMACEUTICA


the drug discovery company

Prescribing Information

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole.

Uses: Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.

Side-effects, precautions, contra-indications: contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy, if a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.



active life

SARGAN
300

tiaprofenic acid

pediatric suspension

NEW DRUGS


In the past few years the number of important new drugs and our understanding of pharmacology have continued to increase. Reliable and unbiased information on the therapeutic use of these agents is, however, not always readily available. Articles recently published in the *BMJ* on entirely new groups of drugs – H₂ receptor antagonists, calcium antagonists, captopril – and on new members of groups of drugs already available – beta-blockers, tranquillisers, hypnotics, diuretics – fill this gap and are now collected together in book form. Busy practitioners will find that this comprehensive review allows them to make a more rational choice of treatment.

Price: Inland £6.50;
Overseas £7.50/USA\$13.00
(Inland £6.00;
Overseas £7.00/USA\$12.00 to
BMA members)

Despatched by air overseas

Payment must be enclosed with order

Order your copy now
From: The Publisher
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR
or any leading bookseller



ABC OF COMPUTING

A J ASBURY

Although computers are being widely used in medicine, their possibilities and limitations are still not clear to many potential users. This book, aimed at the non-expert, describes some of the uses of computers in medicine; because most doctors' involvement will be indirect, liaising with computer experts rather than designing systems themselves, the book concentrates on concepts rather than detailed descriptions of how computers work. It provides a useful introduction for the doctor who wants to know how computers can contribute to his practice of medicine.

Price: Inland £5.75;
Overseas £8.00/USA\$14.00
(Inland £5.25;
Overseas £7.50/USA\$13.00
to BMA members) *including postage,
by air overseas*

Please quote membership number

Payment must be enclosed with order



"This book is worthwhile reading for anyone who is wondering what computers might be able to do for them in their own situation."

NZ med j 1984; 97: 463

"This book provides a useful, quick guide to medical computing in a user-friendly two-column package . . . It contains sound advice on acquiring and developing a system, and throughout it is "them" not "us" who are the programmers."

World Medicine 1984; 19: 26

Order your copy now from: The Publishing Manager, BRITISH MEDICAL JOURNAL, BMA House, Tavistock Square, London WC1H 9JR, *or any leading bookseller.*

WHAT'S A GOOD DOCTOR WITHOUT GOOD DRUGS ?

A high-contrast, black and white photograph of a hand holding a glass vial. The hand is positioned on the left, with fingers wrapped around the neck of the vial. The vial is tilted, and its body is visible, showing a dark liquid inside. A label on the vial features the Janssen logo and the text 'JANSSEN PHARMACEUTICA' and 'the Johnson & Johnson company'. The background is dark, making the hand and vial stand out.

Good doctors treat patients, not just diseases or symptoms. Diagnosis is their key to selecting the most appropriate treatment, always weighing risks against benefits.

And when it comes to drug treatment, they like to rely on the best products from innovative drug research.

They know that ever more effective and safer drugs are indispensable to progress in medicine.