


Nizoral^{TRADEMARK}

(ketoconazole) (tablets)

over 3 million prescriptions world-wide:

the new beginning in antifungal therapy.



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but problematic fungal infections

can now be treated effectively and conveniently. Typically, in recurrent *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



JANSSEN
PHARMACEUTICA

the drug discovery company

Prescribing information

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole.

Uses: Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.

Side-effects, precautions, contra-indications: contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂ blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.

NO-SPA[®]

spasmolytic (Drotaverine)

tablets,
ampoules

Potent antispasmodic for relaxation
of smooth muscles

No-Spa is
—more potent than papaverine
—non toxic
—well absorbed from gastro
intestinal tract


No-Spa
—acts directly on hyper-tonic
smooth muscles
—does not possess anticholinergic
property

No-Spa has
—rapid onset of action
—minimal minor side effect
—high tolerance

Indications
—smooth muscle spasm colics of any origin
for short and long term therapy
—of gastrointestinal biliary and urogenital
spastic conditions, dysmenorrhoea
peripheral vascular disorders, coxarthrosis
coronary insufficiency, angina pectoris

Produced by Chinoin Pharmaceutical
and Chemical Works Ltd — Budapest — Hungary

Exported by
Medimpex Hungarian Trading Company
for Pharmaceutical Products
Budapest - Hungary



WHAT'S A GOOD DOCTOR WITHOUT GOOD DRUGS ?

Good doctors treat patients, not just diseases or symptoms. Diagnosis is their key to selecting the most appropriate treatment, always weighing risks against benefits.

And when it comes to drug treatment, they like to rely on the best products from innovative drug research.

They know that ever more effective and safer drugs are indispensable to progress in medicine.

