

# Nizoral<sup>®</sup>

(ketoconazole) (tablets)

over 3 million  
prescriptions  
world-wide:

the new beginning  
in antifungal therapy.



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but problematic fungal infections

can now be treated effectively and conveniently. Typically, in recurrent *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.

 **JANSSEN**  
PHARMACEUTICA

the drug discovery company

Prescribing information:  
Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole.  
Uses: Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.  
Side-effects, precautions, contra-indications: contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anticholinergic drugs, antacids, H<sub>2</sub>-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.

# Toborol<sup>®</sup>

non-cardioselective  
beta-blocker

tablets  
(CLORANOLOL)



**Cloranolol** is a non-cardio-selective beta-adrenergic receptor blocking agent. It is 3 to 10 times more active than propranolol. It does not possess any sympathomimetic action. It exerts a membrane-stabilizing "quinidine-like" effect. It is highly effective in the treatment of hypertensive angina pectoris, different irregularities of the cardiac rhythm, essential circulatory hyperkinetic hypertrophic obstructive cardiomyopathy and hyperthyroidism.

Produced by  
**CHEMICAL WORKS of  
GEDEON RICHTER Ltd.**  
Budapest-Hungary

Exported by  
**MEDIMPEX**  
Budapest-Hungary



# The Cardio-Protectors

**TENORMIN FOR MODERATE HYPERTENSION**  
**TENORMIN LS FOR MILD AND OLDER HYPERTENSIVES**

## 'Tenormin' and 'Tenormin' LS Prescribing Notes

**PRESENTATION:** 'Tenormin' tablets containing atenolol 100mg are round, bi-convex, orange and film coated. 'Tenormin' LS tablets containing atenolol 50mg are round, bi-convex, orange and film coated.

**USES:** Management of Hypertension.

**DOSAGE:** 'Tenormin' 100mg orally once a day and 'Tenormin' LS orally once a day; some patients may respond adequately to 'Tenormin' low strength.

**CONTRA-INDICATIONS:** Heart Block.

**PRECAUTIONS:** Untreated cardiac failure, bradycardia, renal failure, anaesthesia, pregnancy. Caution in patients with chronic obstructive airways disease or asthma. 'Tenormin' modifies the tachycardia of hypoglycaemia. Co-administration of verapamil or Class I antidysrhythmic agents. Withdrawal of clonidine. Withdrawal of beta-blocking drugs should be gradual in patients with ischaemic heart disease.

**SIDE EFFECTS:** Coldness of extremities, bradycardia and muscular fatigue may occur. Sleep disturbance rarely seen. Rashes and dry eyes have been reported with beta-blockers—consider discontinuance if they occur.

**PACK SIZE AND BASIC NHS COST:** 'Tenormin' 28's £6.98; 'Tenormin' LS 28's £4.88.

**PRODUCT LICENCE NUMBER:** 'Tenormin' 29/122; 'Tenormin' LS 29/86.

'Tenormin', 'Tenormin' LS are trademarks.



Full prescribing information is available on request to the Company.  
**Stuart Pharmaceuticals Limited,**  
 Stuart House, 50 Alderley Road, Wilmslow, Cheshire SK9 1RE

