


# Nizoral<sup>TRADEMARK</sup>

(ketoconazole) (tablets)

# over 3 million prescriptions world-wide:

## the new beginning in antifungal therapy.



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but problematic fungal infections

can now be treated effectively *and* conveniently. Typically, in recurrent *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.

Full prescribing information available on request:



**JANSSEN**  
PHARMACEUTICA

*the drug discovery company*

B-2340 Beerse, Belgium

#### Prescribing Information

**Presentation:** white, flat, half scored uncoated tablets marked 'Janssen' on one side and 'K/200' on the reverse. Each tablet contains 200 mg ketoconazole.

**Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in 'at risk' patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed.

**Side effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub>-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy: if a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.



Free to live a more active life

# SURCA 300

tiaprofenic acid

POTENCY IN ARTHRITIS

# If you needed an H<sub>2</sub> antagonist... which would you prefer to take?

**Zantac**  
RANITIDINE

**PRESCRIBING INFORMATION: DOSAGE AND ADMINISTRATION:** THE USUAL ADULT DOSE IS ONE 150 MG TABLET TWICE DAILY OR A SINGLE DOSE OF 300MG AT NIGHT TIME FOR DUODENAL ULCER. IT IS NOT NECESSARY TO TIME THE DOSE IN RELATION TO MEALS. ZANTAC DISPERSIBLE TABLETS SHOULD BE PLACED IN HALF A GLASS OF WATER (MINIMUM 75ML) AND STIRRED UNTIL DISPERSED BEFORE SWALLOWING. IN MOST CASES OF DUODENAL ULCER AND BENIGN GASTRIC ULCER, HEALING WILL OCCUR IN FOUR WEEKS. PATIENTS WITH A HISTORY OF RECURRENT ULCER MAY HAVE AN EXTENDED COURSE OF ONE 150MG TABLET DAILY AT BEDTIME. FOR REFLUX OESOPHAGITIS THE RECOMMENDED COURSE FOR ADULTS IS ONE 150MG TABLET TWICE DAILY FOR UP TO EIGHT WEEKS. **SIDE EFFECTS:** HEADACHES, DIZZINESS AND SKIN RASHES HAVE BEEN REPORTED. ANAPHYLACTOID REACTIONS HAVE BEEN SEEN RARELY. **PRECAUTIONS:** WHERE GASTRIC ULCER IS SUSPECTED, THE POSSIBILITY OF MALIGNANCY SHOULD BE EXCLUDED BEFORE THERAPY IS INSTITUTED. PATIENTS RECEIVING PROLONGED TREATMENT SHOULD BE EXAMINED PERIODICALLY. DOSAGE SHOULD BE REDUCED IN THE PRESENCE OF SEVERE RENAL IMPAIRMENT (SEE DATA SHEET). AS WITH ALL DRUGS, ZANTAC SHOULD BE USED DURING PREGNANCY AND NURSING ONLY IF STRICTLY NECESSARY. **CONTRA-INDICATIONS:** THERE ARE NO KNOWN CONTRA-INDICATIONS TO THE USE OF ZANTAC. **BASIC NHS COST** (EXCLUSIVE OF VAT): (60 X 150MG TABLETS £27.43, 30 X 300MG TABLETS £27.43, 60 X 150MG DISPERSIBLE TABLETS £28.80. **PRODUCT LICENCE NUMBERS:** 4/0279 (150MG), 4/0302 (300MG), 4/0298 (DISPERSIBLE).

ZANTAC IS A GLAXO TRADE MARK.  
FURTHER INFORMATION IS AVAILABLE ON REQUEST FROM:  
GLAXO LABORATORIES LIMITED, GREENFORD, MIDDLESEX UB6 0HE.


**Glaxo**



*Essential facts on plastic surgery*

# ABC OF PLASTIC AND RECONSTRUCTIVE SURGERY

DAI DAVIES

*Just published*

Spectacular advances in the techniques of plastic and reconstructive surgery mean that surgeons can now reposition amputated limbs and restore function to lacerated hands; they also play a vital part in multidisciplinary teams treating patients with head and neck cancer. *ABC of Plastic and Reconstructive Surgery* brings the non-specialist up to date on these advances and explains what can be done about congenital abnormalities and the disfiguring effects of trauma or disease. Fully illustrated to show the surgical procedures used and the results that can be achieved, it includes chapters on cleft lip and palate; burns; scars; benign and malignant skin conditions; cosmetic and reconstructive surgery of the breast; and cranifacial and microvascular surgery. A series from the *British Medical Journal*, now collected in one convenient volume with specially improved reproduction of the illustrations.

Price: Inland £6.50; Abroad £9.00; USA \$15.00

BMA members: Inland £6.00; Abroad £8.50; USA \$14.00 (please quote membership number)

*Prices include postage, by air abroad. Payment must be enclosed with order.*

**Order your copy from:** The Publishing Manager, BRITISH MEDICAL JOURNAL, BMA House, Tavistock Square, London WC1H 9JR, or any leading bookseller.

# A SENSE OF ASHER

## A SENSE OF ASHER

*A new miscellany*



### ***Why are medical journals so dull?***

Richard Asher, who asked the question, was never dull: good sense, pungent wit, and lively humour were his hallmarks, while his writings on clinical matters, with their combination of lucidity, sympathy, and insight, remain models for all aspiring medical authors. *A Sense of Asher*, a selection of his writings chosen and introduced by Ruth Holland, was first published in a Keynes Press limited edition which quickly sold out. This paperback version, now in its third reprinting, contains the complete text of the original, which includes

- ★ **Apriority**
- ★ **The physical basis of mental illness**
- ★ **A case of health**
- ★ **The use of statistics in medicine**
- ★ **Intracranial and extracranial computers**
- ★ **Why are medical journals so dull?**

"A marvellous gift . . . for a doctor's bedside table" (Selwyn Taylor, *British Medical Journal*)

"The richness of the store is shown by the excellence of this collection" (Sir Douglas Black, *Medical History*)

"Enjoyable light reading for anyone in medicine" (Paul Beeson, *New England Journal of Medicine*)

Price: Inland £7.00; Abroad £8.50; USA \$14.00  
 BMA members: Inland £6.50; Abroad £8.00; USA \$13.00  
 (please quote membership number.) *Prices include postage,  
 by air abroad. Payment must be enclosed with order.*

**Order from:** The Publishing Manager, British Medical Journal, BMA House, Tavistock Square, London WC1H 9JR, *or any leading bookseller*

## THE ESSENTIAL CORRESPONDENCE COURSE FOR THE M.R.C.P.

**This new course for PART  
I and PART II has been  
completely re-written and  
updated with new clinical  
slides, X-rays and ECG's  
included.**

**Write for full information to:  
J. ARNOLD Dept 1  
(Correspondence Courses)  
19, VILLAGE WAY,  
LONDON SE21 7AN**

## ABC OF POISONING

JOHN HENRY  
GLYN VOLANS  
with contributions  
from others

### Part 1: Drugs

Whether accidental or self induced, poisoning raises many problems of diagnosis and management. In this first part of a comprehensive manual Dr John Henry, Dr Glyn Volans and their colleagues deal mainly with poisoning from drugs, but also discuss the management of poisoning in general, giving practical information on diagnosis, laboratory tests, and means of elimination. The book also contains a complete up to date list of all drugs used in the treatment of poisoning.

Price: Inland £5.50; Overseas £8.50/USA \$11.50  
(Inland £5.00; Overseas £8.00/USA \$10.50 to  
BMA members)

*Despatched by air overseas*

**Payment must be enclosed with order**

**Order your copy now**

From: The Publishing Manager, British Medical Journal,  
BMA House, Tavistock Square, London WC1H 9JR,  
or any leading bookseller

## APPROPRIATE TECHNOLOGY

Many of the difficulties of Third World medicine—the effects of poverty, starvation, overpopulation, and political unrest—are beyond the individual doctor's control; but what the Western trained doctor practising in the developing world can and must do is to adapt his practice to compensate for the absence of sophisticated equipment, trained staff, and expensive drugs. Written by doctors and technicians with first hand experience of the problems, *Appropriate Technology* is a guide to the sensible choice and use of resources.

*The British Medical Journal has subsidised this title to make this information widely available at as low cost as possible.*

Price: Inland £2.50; Abroad £5.00/USA \$8.00  
BMA members (please quote membership number):  
Inland £2.00; Abroad £4.50/USA \$7.00


*Despatched by air overseas*

**Payment must be enclosed with order**

**Order your copy now**

From: The Publishing Manager, British Medical Journal,  
BMA House, Tavistock Square, London WC1H 9JR  
or any leading bookseller

## APPROPRIATE TECHNOLOGY





# THE CARDIOPROTECTIVE CORONARY THERAPEUTIC AGENT ADALAT:

## THE MODERN COMPREHENSIVE DRUG THERAPY OF CORONARY HEART DISEASE.



**Indications:** Treatment and prophylaxis of acute and chronic coronary insufficiency (especially angina pectoris, postinfarction syndrome). **Side effects:** Adalat is well tolerated. Side effects occur rarely. They are more common at the beginning of treatment, usually transient, and, in most cases, mild. Occasionally there can be headache, facial flush, heat sensation, dizziness, nausea and tiredness. Extensive biochemical studies did not reveal any abnormal values which could have been ascribed to the drug. **Contraindication:** Pregnancy. **Precaution:** Adalat can enhance or supplement the action of hypotensive preparations and  $\beta$ -blocking agents. Therapy with cardiac glycosides can be initiated or continued in association with Adalat treatment. Incompatibility with other drugs has not been found. **Dosage and administration:** In longterm treatment the capsule is usually swallowed whole with a little liquid. The average daily dose is 3 x 1 capsule, in special cases up to 3 x 2 capsules. In these patients the interval between the individual doses should not be less than two hours. In case of imminent anginal attack, a rapid onset of action is achieved by biting the capsule. **Composition:** One capsule contains 10 mg 1,4-dihydro-2,6-dimethyl-4-(o-nitrophenyl)-pyridine-3,5-dicarboxylic acid dimethyl ester (nifedipine). **Presentation:** Box of 30 or 50 capsules.

Bayer Germany

THE CARDIOPROTECTIVE  
CORONARY THERAPEUTIC AGENT



E 4/043

