

# Nizoral<sup>TRADEMARK</sup>

(ketoconazole) tablets

over 5 million  
prescriptions  
world-wide.

**Oral medication  
in antifungal therapy:**



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal infections can now be treated effectively and elegantly.

Typically, in *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



**JANSSEN  
PHARMACEUTICA**

the drug discovery company

#### Prescribing Information

**Presentation:** white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub>-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long-term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances gynaecomastia has been reported.

# Unique Heme Compound NORMOSANG

heme arginate (25 mg heme/ml  
infusion concentrate)



## For the treatment of acute hepatic porphyrias



Packing: 4 x 10 ml ampoules

- quickly efficacious
- stable for at least two years at +2°C to +8°C
- easily administered, well tolerated
- no risk for AIDS or hepatitis B



Huhtamäki Oy-Medica Pharmaceutical Company Ltd.  
Teollisuuskatu 23-25  
SF-00510 HELSINKI, FINLAND  
Tel. (+358 0) 750 281  
Telex 121720 medic sf

Registered in Finland. Available by special request in other countries.



# GOOD DRUGS DON'T GROW ON TREES

*In the prehistoric days of medicine  
some drugs could be found  
growing on trees.  
Today that's a bit different.*

*The therapeutic drugs of our times  
are being unmasked from nature  
rather more scientifically.*

*By thorough bio-medical research  
into the fundamental processes of life.  
By studying exactly why and how  
these processes sometimes go wrong.*

*And so nature shows man where and how  
he can intervene and defend himself.  
With precision drugs that will selectively  
and safely correct what went wrong.  
Today's drugs grow from man's scientific  
knowledge of nature —  
but no longer on trees.*