

Nizoral^{TRADEMARK}

(ketoconazole) tablets

over 5 million prescriptions world-wide.

Oral medication in antifungal therapy:



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal infections can now be treated effectively and elegantly.

Typically, in *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



JANSSEN
PHARMACEUTICA

the drug discovery company

Prescribing Information

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I.-tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long-term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.



INSTITUTE OF MEDICAL ETHICS

Tavistock House North, Tavistock Square, London WC1H 9LG
Tel: 01-387 8132

The Governing Body of the Institute has decided, as a major step in its development, to offer individual membership.

We invite you now to join the Institute. You will receive twelve monthly issues of our bulletin which will give you a unique service of information on medico-moral issues for a subscription of £30.

ABOUT THE INSTITUTE

The Institute is the only major national resource in medical ethics. It is non-partisan and independent. It has grown from a sure base in the London Medical Group (1963) and the network of autonomous provincial medical groups that provide lectures and symposia in most of the university teaching hospitals.

The Institute also publishes the quarterly *Journal of Medical Ethics*, founded in 1975, that has become a highly respected forum for multi-disciplinary analysis, and for which there is a separate subscription.

More recently, the Institute has pioneered research by multi-disciplinary working groups. Reports have been published on the following subjects: the ethics of resource allocation in health care; dilemmas of dying; and the ethics of clinical research investigations on children.

Medical ethics now concerns us all. The rapid developments in medicine and the powerful impact of television and other media are bringing widespread discussion and interest. These factors must not outstrip moral codes. That is our task.

WHO WE ARE

President

Sir Douglas Black

Vice Presidents

A S Duncan *Emeritus Professor of Medical Education,*

Emeritus Dean of Faculty of Medicine, University of Edinburgh

G R Dunstan *Emeritus Professor of Moral & Social Theology,*

University of London

R B Welbourn *Emeritus Professor of Surgical Endocrinology,*

University of London

Director Prebendary E F Shotton

Deputy Director Dr Richard Nicholson *Bulletin Editor*

Scottish Director Dr Kenneth Boyd

Associate Directors

Brendan Callaghan SJ

Dr Don Evans

Dr Raanan Gillon *Journal Editor*

Dr Patrick Magee

Chairman, Finance Committee

The Earl of Limerick

Treasurer Allan Chappell

Governing Body

R B Welbourn, *Chairman*

R D Catterall *Consultant Venereologist*

Anthony Clare *Professor of Psychological Medicine,*

St Bartholomew's HMC

Prof Sir Cyril Clarke *Past President, Royal Coll of Physicians*

Colin Currie *Consultant Geriatrician, Edinburgh*

Sir John Donne *Chairman NHS Training Authority*

Harold Edwards *Consultant Physician*

Prof C M Fletcher *Clinical Epidemiologist*

Prof T W Glenister *Dean, Charing X & Westminster MS*

Roger Higgs *General Practitioner*

Prof Bryan Jennett *Dean, Faculty of Medicine Glasgow*

Prof Ian Kennedy *King's Coll Centre of Medical Law & Ethics*

Hon David Layton *IME Publications Ltd*

Stephen Lock *Editor of British Medical Journal*

John Mahoney SJ *Moral Theologian, Heythrop College*

Thomas Oppé *Professor of Paediatrics, St Mary's HMS*

Sir Desmond Pond *Past President Royal Coll of Psychiatrists*

A R Peacocke *Ian Ramsey Centre, Oxford*

Robert Sells *Consultant Renal Surgeon, Liverpool*

Paul Sieghart *Council of Science and Society*

Richard West *Dean, St. George's HMS*

Jennifer Wilson-Barnett *Reader in Nursing Studies, Chelsea College*

Applications for membership should be made to IME Publications Limited, 151 Great Portland Street, London W1N 5PB. This will give membership for twelve months and is best done by completing the form below.

I wish to become a member of the Institute of Medical Ethics and receive 12 monthly issues of the IME Bulletin. Please invoice me for £30 ☐ \$60 (overseas) ☐

Name Title/Position

Address

.....

ABC OF SEXUALLY TRANSMITTED DISEASES

MICHAEL W ADLER


AIDS and genital herpes are only the more dramatic signs of a general increase in the number and range of sexually transmitted diseases. Yet these conditions are not seen only in sexually transmitted disease clinics and doctors need to be aware that common clinical problems such as vaginal discharge, rashes, and pelvic pain may have a sexual origin.

Professor Michael Adler's ABC describes the presentation, diagnosis and management of these conditions, emphasising the need to take a broad clinical view of patients and their problems.

Price: Inland £4.00;
Overseas £6.00/USA\$10.50
(BMA members: Inland £3.50;
Overseas £5.50/USA\$9.50)
Despatched by air overseas

Please quote membership number

Payment must be enclosed with order



"This book is recommended for anyone wishing to bring himself up to date with an increasingly expanding speciality . . ."

J roy nav med Serv 1984; 70: 187-188

"... Professor Adler has successfully provided a clear concise and up to date textbook on sexually transmitted disease . . . His approach, refreshingly problem orientated, deals with the patient rather than the disease, and emphasises the psychological impact on the patient of a diagnosis to which much stigma is still attached."


Scott Med J 1985; 30: 69

BOOKS FROM THE BMJ

Order from The Publishing Manager, British Medical Journal, BMA House, Tavistock Square,
London WC1H 9JR or any leading bookseller

Unique Heme Compound NORMOSANG

heme arginate (25 mg heme/ml
infusion concentrate)



For the treatment of acute hepatic porphyrias



- quickly efficacious
- stable for at least two years at +2°C to +8°C
- easily administered, well tolerated
- no risk for AIDS or hepatitis B



Packing: 4 x 10 ml ampoules

Huhtamäki Oy-Medica Pharmaceutical Company Ltd.
Teollisuuskatu 23-25
SF-00510 HELSINKI, FINLAND
Tel. (+358 0) 750 281
Telex 121720 medic sf


Registered in Finland. Available by special request in other countries..

one tablet a day

Tenormin

atenolol

the most cardioselective,
hydrophilic beta-blocker available



CARDIOSELECTIVITY MEANS

- Wide patient selection — may be preferable in asthmatics, diabetics and patients who smoke and drink coffee

HYDROPHILICITY MEANS

- Easy, predictable dosing
- Low incidence of CNS side effects
- Low risk of drug interactions



Full information is available from:
Imperial Chemical Industries PLC
Pharmaceuticals Division
Alderley Park, Macclesfield
Cheshire, England

'Tenormin' is a trade mark for atenolol.

Prescribing notes **Contraindications:** Heart block. Co-administration with verapamil. **Precautions:** Untreated cardiac failure, bradycardia, renal failure, anaesthesia and pregnancy. Withdrawal of clonidine. **Side effects:** Coldness of extremities and muscular fatigue may occur. Sleep disturbance rarely seen. Rashes and dry eyes have been reported with beta-blockers — consider discontinuance if they occur. Withdrawal of beta-blocker should be gradual.